

CONJUGACY CLASSES OF LINEAR ACTIONS IN THE PLANE CREMONA GROUP

IVAN CHELTISOV, YURI TSCHINKEL, AND ZHIJIA ZHANG

ABSTRACT. We classify regular generically free actions of finite groups on the projective plane, up to conjugation in the Cremona group.

1. INTRODUCTION

One of the most intensely studied objects in algebraic geometry is the plane Cremona group

$$\mathrm{Cr}_2 = \mathrm{Bir}(\mathbb{P}^2),$$

the group of birational automorphisms of the plane. Recall that, over an algebraically closed field of characteristic zero,

$$\mathrm{Cr}_2 := \langle \mathrm{PGL}_3, \iota \rangle,$$

where

$$(1.1) \quad \begin{aligned} \iota : \quad \mathbb{P}^2 &\dashrightarrow \mathbb{P}^2, \\ (x, y, z) &\mapsto \left(\frac{1}{x}, \frac{1}{y}, \frac{1}{z} \right) \end{aligned}$$

is the standard Cremona involution. Conjugacy classes of finite subgroups of PGL_3 have been described in [Bli18]: there are three types, recalled in Section 3:

- intransitive,
- transitive but imprimitive,
- primitive.

There exist isomorphic non-conjugated *finite subgroups* of PGL_3 which are conjugated in Cr_2 , see Sections 3 and 6. However, finite subgroups of different types are not conjugated in Cr_2 .

A century later, in the seminal paper by Dolgachev and Iskovskikh [DI09], came the classification of finite subgroups of Cr_2 , i.e., finite groups that can act regularly and generically freely on rational surfaces. However, the classification of *subgroups* as well as of *actions*, up to

conjugation in Cr_2 , remained an open problem. For finite subgroups of PGL_3 , this was explicitly asked in [DI09, Section 9]:

Find the conjugacy classes in Cr_2 of actions of finite subgroups of PGL_3 . For example, there are two actions of the subgroup of PGL_3 isomorphic to $\mathfrak{A}_5$ and four actions of $\mathfrak{A}_6$ which differ by outer automorphisms of the groups. Are they conjugated in Cr_2 ?

Since then, it was shown in [Che14] that the two $\mathfrak{A}_5$ -actions are conjugated in Cr_2 , while the four $\mathfrak{A}_6$ -actions are not. A similar problem for the remaining primitive actions follows from [Sak19], see Section 4 for an explicit solution. Furthermore, conjugacy classes of actions of intransitive *abelian* subgroups of PGL_3 have been described in [RY02]; in particular, actions of finite cyclic subgroups are always conjugated in Cr_2 .

In this paper, we completely settle the Dolgachev–Iskovskikh problem for the remaining finite subgroups of PGL_3 . Our main new tool to distinguish the actions up to conjugation in Cr_2 is the formalism of Burnside invariants, introduced in [KT22a] and recalled in Section 2. These fail to distinguish primitive actions [TYZ24, Proposition 7.1 and Example 7.2], and are inconclusive for some imprimitive actions [TYZ24], but those are accessible via the equivariant Sarkisov program. However, the new invariants are decisive in the case of intransitive actions. One of the main results of this paper, proved in Section 6, is:

Theorem 1.1. *Nonabelian intransitive actions $\phi_1, \phi_2 : G \hookrightarrow \mathrm{PGL}_3$ are conjugated in Cr_2 if and only if*

$$[\mathbb{P}^2 \hookrightarrow \phi_1(G)] = [\mathbb{P}^2 \hookrightarrow \phi_2(G)] \in \mathrm{Burn}_2^{\mathrm{inc}}(G).$$

We also obtain an explicit geometric description of this condition, see Theorem 6.1.

Our second main result, proved in Section 5, is:

Theorem 1.2. *Distinct imprimitive actions $\phi_1, \phi_2 : G \hookrightarrow \mathrm{PGL}_3$ are conjugated in Cr_2 if and only if they are conjugated by the standard Cremona involution.*

Together with existing results concerning primitive actions, this settles the Dolgachev–Iskovskikh problem, quoted above [DI09, Section 9]. The proof proceeds via classification of groups and actions, followed by a detailed analysis of equivariant Sarkisov links. As a byproduct, we compute the normalizer of G in the Cremona group in many cases, and answer a question posed by L. Katzarkov:

Theorem 1.3. *Let $G \subset \mathrm{PGL}_3$ be a finite subgroup. Then the normalizer of G in Cr_2 is finite if and only if G is transitive and not isomorphic to any of the following groups:*

$$\mathfrak{A}_4, \quad C_3^2, \quad C_3 \rtimes \mathfrak{S}_3, \quad C_7 \rtimes C_3, \quad C_3^2 \rtimes C_4.$$

Recall that Cr_2 is generated by involutions, over any perfect field [LS24]. Thus, it is reasonable to expect that the normalizer in Cr_2 of any finite subgroup $G \subset \mathrm{Cr}_2$ is also generated by involutions. However, this is not always the case: if $G \subset \mathrm{PGL}_3$ is isomorphic to the *Hessian group*, then G is unique in Cr_2 , up to conjugation [DI09], coincides with its normalizer in Cr_2 by [Sak19], but is not generated by reflections. One can check that normalizers of other transitive subgroups in PGL_3 are generated by reflection.

Keeping in mind results in [CLdC13, Lon16, Zim18], one could ask: When is the normalizer in Cr_2 of a finite subgroup $G \subset \mathrm{Cr}_2$ simple? Such G would have to be simple and coincide with their normalizer. By [Che14], there are exactly two such nonabelian finite subgroups in Cr_2 (up to conjugation), both contained in PGL_3 , namely, $\mathfrak{A}_6$ and $\mathrm{PSL}_2(\mathbb{F}_7)$. Using the classification of elements of prime order in Cr_2 , completed in [BB00, dF04, BB04, Bla07], one can also show that a prime order cyclic subgroup $G \subset \mathrm{Cr}_2$ coincides with its normalizer if and only if G is generated by a Geiser or Bertini involution, and the fixed curve of the involution does not admit nontrivial automorphisms. Such subgroups are conjugated in Cr_2 if and only if the fixed curves of their generating involutions are isomorphic.

Acknowledgments: Ivan Cheltsov has been supported by the Leverhulme Trust grant RPG-2021-229, EPSRC grant EP/Y033485/1, and Simons Collaboration grant *Moduli of varieties*. Yuri Tschinkel has been supported by NSF grant 2301983. This work started during the workshop “Birational Geometry and Number Theory” held at ICMS (Edinburgh) in November 2024, has been developed at CIRM (Luminy) during of a semester-long *Morlet Chair* program, and was finished at the Simons Foundation (New York) in July 2025.

2. BASIC NOTIONS IN EQUIVARIANT GEOMETRY

Automorphisms and birational automorphisms. Throughout, k is an algebraically closed field of characteristic zero. Let X be a smooth projective irreducible variety over k and

$$\mathrm{Aut}(X) \subseteq \mathrm{Bir}(X),$$

the group of regular, respectively, birational, automorphisms of X . Let G be a finite group admitting an embedding

$$\phi : G \hookrightarrow \text{Aut}(X),$$

we refer to this as a choice of a G -action on X , and call X a G -variety. For $G \subseteq \text{Aut}(X)$, let

$$\text{Aut}^G(X) \subseteq \text{Bir}^G(X),$$

be the normalizers of G in the respective groups. This group-theoretic definition has the following geometric interpretation: $\text{Aut}^G(X)$ is the group of all G -biregular self-maps of X , i.e., all $\varphi \in \text{Aut}(X)$ such that there exists a $\psi \in \text{Aut}(G)$ and a commutative diagram

$$\begin{array}{ccc} X & \xrightarrow{\varphi} & X \\ g \downarrow & & \downarrow \psi(g) \\ X & \xrightarrow{\varphi} & X \end{array}$$

for all $g \in G$. Similarly, $\text{Bir}^G(X)$ is the group of all G -birational self-maps, defined by the same diagram, with $\varphi \in \text{Bir}(X)$. There are natural homomorphisms

$$(2.1) \quad \begin{array}{ccc} \text{Aut}^G(X) & \hookrightarrow & \text{Bir}^G(X) \\ & \searrow \alpha & \swarrow \beta \\ & \text{Aut}(G) & \end{array}$$

and we denote the kernels, i.e., centralizers of the respective groups, by

$$\text{ZAut}^G(X) \subseteq \text{ZBir}^G(X).$$

A birational map $X \dashrightarrow Y$ between G -varieties is called

- G -equivariant, if it commutes with the action of G ,
- G -birational, if it commutes with the action of G , up to a fixed automorphism $\psi \in \text{Aut}(G)$.

Conjugacy of groups. Given a G -variety X and embeddings

$$\phi_1, \phi_2 : G \hookrightarrow \text{Aut}(X)$$

one has the following general

Problem A: When are $\phi_1(G)$ and $\phi_2(G)$ conjugated in $\text{Aut}(X)$?

Problem B: When are $\phi_1(G)$ and $\phi_2(G)$ conjugated in $\text{Bir}(X)$?

In the first case, this means that there is a G -biregular self-map $\varphi : X \rightarrow X$ conjugating $\phi_1(G)$ and $\phi_2(G)$. In the second case, φ is G -birational.

Conjugacy of actions. Consider actions $\phi_1, \phi_2 : G \hookrightarrow \text{Aut}(X)$ on a G -variety X as above. They are conjugated in $\text{Aut}(X)$ if there exists a $\varphi \in \text{Aut}(X)$ such that

$$\phi_2 = \varphi \circ \phi_1 \circ \varphi^{-1}.$$

In this case, we don't distinguish these two actions! In the following, when we refer to action, we always mean up to conjugation in $\text{Aut}(X)$.

Clearly, if ϕ_1 and ϕ_2 give the same action on X , then the groups $\phi_1(G)$ and $\phi_2(G)$ are conjugated in $\text{Aut}(X)$. However, the converse does not always hold.

Problem AA: Assuming that the groups $\phi_1(G)$ and $\phi_2(G)$ are conjugated in $\text{Aut}(X)$, when are the actions ϕ_1, ϕ_2 the same?

We can consider a similar problem in $\text{Bir}(X)$: we say that the actions ϕ_1 and ϕ_2 are conjugated in $\text{Bir}(X)$ if there exists a $\varphi \in \text{Bir}(X)$ such that

$$\phi_2 = \varphi \circ \phi_1 \circ \varphi^{-1}.$$

As before, if ϕ_1 and ϕ_2 are conjugated in $\text{Bir}(X)$, then the groups $\phi_1(G)$ and $\phi_2(G)$ are conjugated in $\text{Bir}(X)$.

Problem BA: Assuming that the groups $\phi_1(G)$ and $\phi_2(G)$ are conjugated in $\text{Bir}(X)$, when are the actions ϕ_1, ϕ_2 conjugated in $\text{Bir}(X)$?

To solve these problems, we can precompose one of the actions with a suitable G -biregular self-map $\varphi \in \text{Aut}(X)$, respectively, with a G -birational map $\varphi \in \text{Bir}(X)$, to obtain

$$\phi_1(G) = \phi_2(G),$$

as subgroups of $\text{Aut}(X)$. Then $\phi_2 = \phi_1 \circ \psi$, for some $\psi \in \text{Aut}(G)$, and the actions are conjugated in $\text{Aut}(X)$ if and only if ψ lies in the image

of the homomorphism α from diagram (2.1). We have a diagram

$$\begin{array}{ccccc} \phi_1(G) = \phi_2(G) & \hookrightarrow & \mathrm{Aut}^G(X) & \hookrightarrow & \mathrm{Bir}^G(X) \\ & & \downarrow \alpha & \swarrow \beta & \\ \mathrm{Inn}(G) & \hookrightarrow & \mathrm{Aut}(G) & \twoheadrightarrow & \mathrm{Out}(G), \end{array}$$

where $\mathrm{Inn}(G)$ and $\mathrm{Out}(G)$ are inner and outer automorphisms of G . Clearly, if $\psi \in \mathrm{Inn}(G)$, the actions are conjugated in $\mathrm{Aut}(G)$. Therefore, Problem AA reduces to

Problem IA: Determine the image of

$$\bar{\alpha} : \mathrm{Aut}^G(X) \rightarrow \mathrm{Out}(G).$$

Similarly, Problem BA reduces to

Problem IB: Determine the image of

$$\bar{\beta} : \mathrm{Bir}^G(X) \rightarrow \mathrm{Out}(G).$$

Using this terminology, we can compute the number of G -actions with a fixed image in $\mathrm{Aut}(X)$ as

$$\frac{|\mathrm{Out}(G)|}{|\mathrm{image}(\bar{\alpha})|}.$$

Similarly, the number of actions with fixed image in $\mathrm{Aut}(X)$, up to conjugation in $\mathrm{Bir}(X)$, is given by

$$\frac{|\mathrm{Out}(G)|}{|\mathrm{image}(\bar{\beta})|}.$$

Problem IA is purely group-theoretical, and its solution can be automated with computer algebra packages such as **Magma**. On the other hand, Problem IB is of birational nature. If we knew the generators of $\mathrm{Bir}^G(X)$ then we could compute the image of $\bar{\beta}$. In practice, this is rarely feasible. However, in some situations, the solution is easy, for example, when

- $\mathrm{Out}(G)$ is trivial, e.g., $G = \mathfrak{S}_n$, with $n \neq 6$,
- $\mathrm{Aut}(X) = \mathrm{Bir}(X)$, so that Problems IA and IB coincide, e.g., if X is a curve or if the canonical class K_X is ample,
- $\mathrm{Aut}^G(X) = \mathrm{Bir}^G(X)$,

- one can identify sufficiently many elements in $\text{Bir}^G(X)$ to prove the surjectivity of $\bar{\beta}$.

Example 2.1. There is a unique $\mathfrak{A}_5 \subset \text{PGL}_3 \simeq \text{Aut}(\mathbb{P}^2)$, up to conjugation. We have $\text{Out}(\mathfrak{A}_5) \simeq C_2$ and

$$\text{Aut}^{\mathfrak{A}_5}(\mathbb{P}^2) \simeq \mathfrak{A}_5.$$

As there are no nontrivial homomorphisms $\mathfrak{A}_5 \rightarrow C_2$, we conclude that there are two $\mathfrak{A}_5$ -actions on $\mathbb{P}^2$. However,

$$\text{Bir}^{\mathfrak{A}_5}(\mathbb{P}^2) \simeq \mathfrak{S}_5,$$

and the homomorphism $\bar{\beta}$ is surjective [Che14]. It follows that both $\mathfrak{A}_5$ -actions on $\mathbb{P}^2$ are conjugated in Cr_2 ; a realization of this conjugation is in [CS16, Lemma 6.3.3], see also Example 2.6.

Additional examples with surjective $\bar{\beta}$ will be given in Section 3.

Equivariant Sarkisov program for rational surfaces. Let X be a smooth rational G -surface. The equivariant Minimal Model Program implies that X is G -birational to a G -surface S such that

- S is a smooth del Pezzo surface with $\text{rkPic}(S)^G = 1$, or
- S admits a G -equivariant conic bundle

$$\pi : S \rightarrow \mathbb{P}^1$$

with $\text{rkPic}(S)^G = 2$.

Such G -birational models are called *G -Mori fiber spaces* (in dimension 2). Any G -birational map between such G -Mori fiber spaces can be factorized into a sequence of elementary G -birational maps, called *G -Sarkisov links*. We recall their classification, following the classification of Sarkisov links between 2-dimensional Mori fiber spaces over nonclosed ground fields in [Isk96]. There are three basic types:

- **DP-DP** (del Pezzo to del Pezzo):

$$S \xleftarrow{\sigma} \tilde{S} \xrightarrow{\sigma'} S',$$

where

- σ is a blowup of a G -orbit Σ such that $|\Sigma| < K_S^2$ and $\tilde{S}$ is a del Pezzo surface,
- S' is a (smooth) del Pezzo G -surface with $\text{rkPic}(S')^G = 1$,
- σ' is a blowup of a G -orbit Σ' such that $|\Sigma'| < K_{S'}^2$;

- **DP-CB** (del Pezzo to conic bundle / and its inverse):

$$S \xleftarrow{\sigma} \tilde{S} \xrightarrow{\tilde{\pi}} \mathbb{P}^1$$

where

- σ is a blowup of a G -orbit Σ such that $|\Sigma| < K_S^2$ and $\tilde{S}$ is a del Pezzo surface,
- $\tilde{\pi}$ is a G -equivariant conic bundle;
- **CB-CB** (conic bundle to conic bundle):

$$\begin{array}{ccccc} S & \xleftarrow{\sigma} & \tilde{S} & \xrightarrow{\sigma'} & S' \\ \pi \downarrow & & & & \downarrow \pi' \\ \mathbb{P}^1 & \xlongequal{\quad} & & & \mathbb{P}^1 \end{array}$$

where

- π' is a G -equivariant conic bundle, with $K_{S'}^2 = K_{\tilde{S}}^2$,
- σ is a blowup of a G -orbit Σ such that no points of Σ are contained in the singular fibers of π and every smooth fiber of π contains at most one point of Σ ,
- σ' is the blowdown of the strict transforms of the fibers of π that contain points of Σ .

Example 2.2. We list G -Sarkisov links that start from $S = \mathbb{P}^2$:

- **DP-DP** with $r := |\Sigma| \in \{2, 3, 5, 6, 7, 8\}$
 - $r = 2$: $S' = \mathbb{P}^1 \times \mathbb{P}^1$ and σ' blows down the strict transform of the line passing through Σ ,
 - $r = 3$: $S' = \mathbb{P}^2$ and σ' blows down the strict transforms of lines passing through pairs of points in Σ ,
 - $r = 5$: S' is a del Pezzo surface of degree 5 and σ' blows down the strict transform of the conic through Σ ,
 - $r = 6$: $S' = \mathbb{P}^2$ and σ' blows down strict transforms of conics passing through 5 points of Σ ,
 - $r = 7$: $\tilde{S}$ is a del Pezzo surface of degree 2 and $\sigma' = \sigma \circ \tau$, where τ is the involution of the anticanonical double cover $\tilde{S} \rightarrow \mathbb{P}^2$,
 - $r = 8$: $\tilde{S}$ is a del Pezzo surface of degree 1 and $\sigma' = \sigma \circ \tau$, where τ is the involution of the double cover $\tilde{S} \rightarrow \mathbb{P}(1, 1, 2)$ given by the linear system $|-2K_{\tilde{S}}|$.
- **DP-CB** with $r = |\Sigma| \in \{1, 4\}$
 - $r = 1$: $\tilde{S}$ is the Hirzebruch surface $\mathbb{F}_1$ and $\tilde{\pi}$ a $\mathbb{P}^1$ -bundle,
 - $r = 4$: $\tilde{S}$ is a del Pezzo surface of degree 5 and $\tilde{\pi}$ a conic bundle with 3 singular fibers.

Remark 2.3. Let S be a smooth del Pezzo G -surface of degree ≥ 2 such that $\mathrm{rkPic}(S)^G = 1$. Suppose that there exists a G -Sarkisov link

$$S \xleftarrow{\sigma} \tilde{S} \xrightarrow{\sigma'} S'$$

such that σ is a blowup of a G -orbit Σ with

$$|\Sigma| \in \{K_S^2 - 1, K_S^2 - 2\},$$

where S' is a (smooth) del Pezzo G -surface with $\mathrm{rkPic}(S)^G = 1$, and σ' is a blowup of a G -orbit Σ' . Then $\tilde{S}$ is a smooth del Pezzo surface of degree 1 or 2, and $\mathrm{Aut}(\tilde{S})$ contains an involution τ that centralizes the action of the group G . The involution τ is known as *Bertini* involution (when $K_{\tilde{S}}^2 = 1$) or *Geiser* involution (when $K_{\tilde{S}}^2 = 2$). This implies that S and S' are G -birational, in particular, we have $|\Sigma'| = |\Sigma|$. Hence, we may assume that $\sigma' \circ \beta = \gamma$ for

$$\gamma = \sigma \circ \tau \circ \sigma^{-1}.$$

Note that $\gamma \in \mathrm{Bir}^G(S)$, and its image $\beta(\gamma) \in \mathrm{Aut}(G)$ is trivial, because γ centralizes G . In the following, we will say that γ is a Bertini or Geiser (birational) involution of the surface S , respectively.

A more detailed description of two-dimensional G -Sarkisov links is in [DI09, Section 7]. For instance, if $S \rightarrow \mathbb{P}^1$ is a G -conic bundle with $\mathrm{rkPic}(S)^G = 2$, then $7 \neq (-K_S)^2 \leq 8$ and the following assertions hold:

- if $(-K_S)^2 \leq 1$, then S is not G -birational to a del Pezzo surface, see [DI09, Section 8],
- if $(-K_S)^2 = 3, 5, 6$, then S is a del Pezzo surface,
- if $(-K_S)^2 = 8$, then S is G -birational to a del Pezzo surface.

If $(-K_S)^2 = 4$, there are examples such that S is a del Pezzo surface, and there are examples such that S is not G -birational to a smooth del Pezzo surface, see [DI09, Section 8]. It was an open question, posed in [DI09, Section 9] whether or not S is G -birational to a smooth del Pezzo surface when $(-K_2)^2 = 2$. Even though this is not directly related to the main topic of this paper, we include an example that answers this question:

Example 2.4. Let $\bar{S} \subset \mathbb{P}(1_x, 1_y, 3_z, 3_w)$ be given by

$$zw = x^6 + y^6,$$

let $\eta: S \rightarrow \bar{S}$ be the minimal resolution of singularities, and let G be the subgroup in $\mathrm{Aut}(\bar{S})$ generated by

$$\sigma_{z,w}: (z, w) \mapsto (w, z), \quad \sigma_{x,y}: (x, y) \mapsto (x, y),$$

$$\theta_{12} : (x, y, z, w) \mapsto (\zeta_{12}x, \zeta_{12}^{-1}y, -z, w).$$

Then $(-K_S)^2 = 2$, $\text{rkPic}(S)^G = 2$, $G \simeq (C_2)^2 \cdot \mathfrak{D}_6$, and we have the following G -equivariant commutative diagram:

$$\begin{array}{ccc} & S & \\ \eta \swarrow & & \searrow \pi \\ \bar{S} & \xrightarrow{\xi} & \mathbb{P}_{x,y}^1 \end{array}$$

where π is a conic bundle, and ξ is given by $(x, y, z, t) \mapsto (x, y)$. By the classification of G -Sarkisov links, S is not G -birational to a smooth del Pezzo surface unless there exists the G -equivariant commutative diagram:

$$\begin{array}{ccc} S & \xrightarrow{\quad} & S' \\ \pi \downarrow & & \downarrow \pi' \\ \mathbb{P}^1 & \xlongequal{\quad} & \mathbb{P}^1 \end{array}$$

where S' is a del Pezzo surface of degree 2 with $\text{rkPic}(S')^G = 2$, and π' is a G -conic bundle. Using the classification of actions on del Pezzo surfaces of degree 2 in [DI09], we exclude the latter possibility, so that S is not G -birational to a smooth del Pezzo surface.

Now, we suppose that S is a del Pezzo surface with $\text{rkPic}(S)^G = 1$. Such S is called:

- G -rationally super-rigid if there are no G -Sarkisov links starting from S ,
- G -rationally rigid if every G -Sarkisov link that starts from S ends at a del Pezzo surface which is G -biregular to S ,
- G -solid if S is not G -birational to a G -conic bundle.

If S is G -rationally super-rigid then $\text{Aut}^G(S) = \text{Bir}^G(S)$, and the number of G -actions on S with fixed image in $\text{Aut}(S)$ is the same, whether we consider it up to conjugation in $\text{Aut}(S)$ or in $\text{Bir}(S)$.

Example 2.5. Let $S = \mathbb{P}^2$ and $G = \mathfrak{A}_6$. Then S does not have G -orbits Σ of length $|\Sigma| < K_S^2 = 9$. By Example 2.2, S is G -rationally super-rigid.

Example 2.6. Let $S = \mathbb{P}^2$ and $G = \mathfrak{A}_5$. Then S contains a *unique* G -orbit Σ of length $|\Sigma| < K_S^2 = 9$, this orbit has length 6 and the points of Σ are in general linear position and are not contained in a conic. By Example 2.2, $S' = \mathbb{P}^2$, and is G -biregular to S , thus S is G -rationally rigid.

Burnside formalism. This formalism, introduced in [KT22a] and applied in, e.g., [HKT21], [KT22b], [TYZ24], allows to distinguish birational types of actions of finite groups in many new situations. It takes into account information from the *stabilizer stratification* on a *standard* birational model for the action. This is a model X such that

- there is a Zariski open $U \subset X$ on which the G -action is free,
- the complement $X \setminus U$ is a normal crossings divisor such that for each irreducible component D_α in $X \setminus U = \cup_{\alpha \in \mathcal{A}} D_\alpha$, and all $g \in G$, we have that

$$g(D_\alpha) = D_\alpha \quad \text{or} \quad g(D_\alpha) \cap D_\alpha = \emptyset.$$

On such a model, generic stabilizers of all subvarieties are abelian.

We describe the simplest version of the general theory in the case of surfaces: given a standard model as above, a G -orbit of irreducible components of $X \setminus U$, and a choice of a representative D_α of this orbit, we have a symbol

$$(H_\alpha, Z_\alpha \curvearrowright k(D_\alpha), (b_\alpha)),$$

where

- H_α is the (cyclic) generic stabilizer of D_α ,
- $Z_\alpha \subseteq Z_G(H_\alpha)/H_\alpha$, a subgroup of the centralizer modulo stabilizer, acting generically freely on D_α , and
- $b_\alpha \in H_\alpha^\vee$ is the weight of H_α in the normal bundle of D_α .

Such symbols are considered up to G -conjugation.

A symbol is called *incompressible* if D_α is a curve of genus ≥ 1 , or if Z_α is not cyclic. The class

$$[X \curvearrowright G] := \sum_{\alpha \in \mathcal{A}/G} (H_\alpha, Z_\alpha \curvearrowright k(D_\alpha), (b_\alpha))$$

in the free abelian group

$$\text{Burn}_2^{\text{inc}}(G),$$

spanned by incompressible symbols (up to conjugation), is an equivariant birational invariant of the G -action.

In our application to Theorem 6.1, we have $D_\alpha = \mathbb{P}^1$, and Z_α is one of the

$$\mathfrak{D}_n, \quad \mathfrak{A}_4, \quad \mathfrak{S}_4, \quad \mathfrak{A}_5.$$

The following example from [KT22b, Section 10] and [TYZ24, Section 7] illustrates this.

Example 2.7. For $t \geq 2$ and odd $n \geq 5$, consider $G = C_t \times \mathfrak{D}_n$. Let χ be a primitive character of C_t and ψ a primitive character of $C_n \subset \mathfrak{D}_n$. Let V_ψ be the 2-dimensional faithful representation of $\mathfrak{D}_n$ induced from ψ , and $V_{\chi,\psi} = \chi \otimes V_\psi$. Then G acts generically freely on

$$\mathbb{P}^2 = \mathbb{P}(V) = \mathbb{P}(\mathbf{1} \oplus V_{\chi,\psi}).$$

The class

$$[\mathbb{P}(V) \hookrightarrow G] \in \text{Burn}_2^{\text{inc}}(G)$$

equals

$$(C_t, \mathfrak{D}_n \curvearrowright k(\mathbb{P}^1), (\chi)) + (C_t, \mathfrak{D}_n \curvearrowright k(\mathbb{P}^1), (-\chi)).$$

In particular, if $\chi \neq \pm\chi'$ or $\psi \neq \pm\psi'$, the corresponding actions on $\mathbb{P}(V)$ and $\mathbb{P}(V')$ are not birational.

3. CLASSIFICATION OF FINITE SUBGROUPS IN PGL_3

Finite subgroups $G \subset \text{PGL}_3$ have been classified by Blichfeldt [Bli18]. These arise via projectivizations $\mathbb{P}(V)$ of faithful 3-dimensional representations V of central extensions $\tilde{G}$ of G . We recall the relevant terminology: a G -action on $\mathbb{P}^2 = \mathbb{P}(V)$ is called:

- (I) *intransitive*: if G fixes a point in $\mathbb{P}^2$;
- (T) *transitive but imprimitive*: if G does not fix a point in $\mathbb{P}^2$ but there exists a G -orbit of length 3;
- (P) *primitive*: neither of the above.

The following groups give rise to primitive actions:

$$\mathfrak{A}_5, \quad \mathfrak{A}_6, \quad \text{PSL}_2(\mathbb{F}_7), \quad \text{ASL}_2(\mathbb{F}_3), \quad \text{PSU}_3(\mathbb{F}_2), \quad C_3^2 \rtimes C_4.$$

Each of these is unique in PGL_3 , up to conjugation. Explicit generators are given in [YY93]. The group $\text{ASL}_2(\mathbb{F}_3)$ is known as the *Hessian group* — after a suitable coordinate change, it leaves invariant the Hesse pencil:

$$x^3 + y^3 + z^3 + \lambda xyz = 0, \quad \lambda \in \mathbb{P}^1,$$

in $\mathbb{P}_{x,y,z}^2$. The action on the curves of this pencil gives an exact sequence of groups

$$1 \rightarrow C_3^2 \rtimes C_2 \rightarrow \text{ASL}_2(\mathbb{F}_3) \rightarrow \mathfrak{A}_4 \rightarrow 1,$$

where $C_3^2 \rtimes C_2 \subset \text{PGL}_3$ is the subgroup that leaves invariant a general curve in the pencil, and $C_3^2 \subset C_3^2 \rtimes C_2$ acts on a general elliptic curve in the pencil via translations by 3-torsions. This subgroup is transitive but not primitive, and $\text{ASL}_2(\mathbb{F}_3)$ can also be defined as its normalizer

in PGL_3 . Up to conjugation, the three primitive subgroups $C_3^2 \rtimes C_4$, $\mathrm{PSU}_3(\mathbb{F}_2)$, $\mathrm{ASL}_2(\mathbb{F}_3)$ are nested as follows:

$$C_3^2 \rtimes C_4 \subset \mathrm{PSU}_3(\mathbb{F}_2) \subset \mathrm{ASL}_2(\mathbb{F}_3).$$

We describe groups giving rise to actions of type **(T)**. In this case, G has an orbit of length 3, consisting of points in linear general position. Changing coordinates, we may assume that these points are

$$[1 : 0, 0], \quad [0 : 1 : 0], \quad [0 : 0 : 1].$$

The G -action on these points gives rise to the exact sequence

$$1 \rightarrow T \rightarrow G \xrightarrow{\nu} \mathfrak{S}_3,$$

which turns out to be split. The kernel of ν consists of diagonal automorphisms. The image is either C_3 or $\mathfrak{S}_3$.

Proposition 3.1 ([DI09, Theorem 4.7]). *Let $G \subset \mathrm{PGL}_3 \simeq \mathrm{Aut}(\mathbb{P}^2)$ be a finite subgroup giving rise to an action of type **(T)**. Then, up to conjugation in PGL_3 , one of the following holds:*

- $G \simeq C_n^2 \rtimes C_3$ is generated by

$$\mathrm{diag}(\zeta_n, 1, 1), \quad \sigma_{123}.$$

- $G \simeq C_n^2 \rtimes \mathfrak{S}_3$ is generated by

$$\mathrm{diag}(\zeta_n, 1, 1), \quad \sigma_{123}, \quad \sigma_{12},$$

- $G \simeq (C_n \times C_{n/r}) \rtimes C_3$ is generated by

$$\mathrm{diag}(\zeta_n^r, 1, 1), \quad \mathrm{diag}(\zeta_n^s, \zeta_n, 1), \quad \sigma_{123},$$

where $r \geq 1$, $r \mid n$, and $s^2 - s + 1 \equiv 0 \pmod{r}$,

- $G \simeq (C_n \times C_{n/3}) \rtimes \mathfrak{S}_3$ is generated by

$$\mathrm{diag}(\zeta_n^3, 1, 1), \quad \mathrm{diag}(\zeta_n^2, \zeta_n, 1), \quad \sigma_{123}, \quad \sigma_{12},$$

with $3 \mid n$,

where

$$\sigma_{123} : (x, y, z) \rightarrow (y, z, x), \quad \sigma_{12} : (x, y, z) \rightarrow (y, x, z),$$

on $\mathbb{P}_{x,y,z}^2$. Each of these subgroups is unique up to conjugation in PGL_3 .

Proof. The classification is achieved in [DI09]. To show the uniqueness of each subgroup in PGL_3 , it suffices to show that the two choices of s give rise to the same subgroup in the third case $G \simeq (C_n \times C_{n/r}) \rtimes C_3$.

Let G_1, G_2 be groups corresponding to different choices of roots s_1, s_2 , for fixed n, r , in the third case. Since $s_1 + s_2 = 1 \pmod{r}$, we have

$$\sigma_{12} \cdot \mathrm{diag}(\zeta_n^{s_1}, \zeta_n, 1) \cdot \sigma_{12} = \mathrm{diag}(\zeta_n^{s_1}, 1, \zeta_n).$$

Up to multiplying by powers of $\text{diag}(\zeta_n^r, 1, 1)$, we have

$$\text{diag}(\zeta_n^{s_2}, \zeta_n, 1) \in \sigma_{12} G_1 \sigma_{12}.$$

It follows that

$$\sigma_{12} \cdot G_1 \cdot \sigma_{12} = G_2.$$

□

We turn to actions of type **(I)**, following [DI09, Section 4.2]. The existence of a fixed point implies that we have an embedding $G \hookrightarrow \mathbf{GL}_2$, inducing an action on $\mathbb{P}^1$, via the projection $\mathbf{GL}_2 \rightarrow \mathbf{PGL}_2$; we denote by $\bar{G}$ the image of G , and by C_m its kernel, a cyclic subgroup of order m . We write

$$\chi_r : (x_1, x_2, x_3) \mapsto (x_1, \zeta_r x_2, \zeta_r x_3),$$

for a primitive root of unity ζ_r of order $r \geq 1$. We have:

Proposition 3.2 ([NvdPT08]). *Let $G \subset \mathbf{PGL}_3 \simeq \text{Aut}(\mathbb{P}^2)$ be a finite subgroup giving rise to an action of type **(I)**. Then, up to conjugation in $\mathbf{PGL}_3$, one of the following holds:*

- $\bar{G} \simeq C_n$. G is generated by

$$\chi_r, (x_1, \zeta_n^r x_2, \zeta_n^m x_3), \quad r, m \in \mathbb{Z}_{\geq 0}.$$

- $\bar{G} \simeq \mathfrak{D}_n$:

- n odd. G is generated by

$$\chi_r, (x_1, \zeta_n x_2, \zeta_n^{-1} x_3), (x_1, \zeta_{2^m} x_3, \zeta_{2^m} x_2), \quad r, m \in \mathbb{Z}_{\geq 0}.$$

- n even. G is generated by

$$\chi_r, (x_1, \zeta_{2n} x_2, \zeta_{2n}^{-1} x_3), (x_1, \zeta_4 x_3, \zeta_4 x_2), \quad r \in \mathbb{Z}_{\geq 0}.$$

- n even. G is generated by

$$\chi_r, (x_1, \zeta_{2^{m+1}} \zeta_{2n} x_2, \zeta_{2^{m+1}} \zeta_{2n}^{-1} x_3), (x_1, \zeta_4 x_3, \zeta_4 x_2), \quad r \in \mathbb{Z}_{\geq 0}, m \in \mathbb{Z}_{\geq 1}.$$

- n even, $n \neq 2$. G is generated by

$$\chi_r, (x_1, \zeta_{2n} x_2, \zeta_{2n}^{-1} x_3), (x_1, \zeta_{2^{m+1}} \zeta_4 x_3, \zeta_{2^{m+1}} \zeta_4 x_2), \quad r \in \mathbb{Z}_{\geq 0}, m \in \mathbb{Z}_{\geq 1}.$$

- $\bar{G} \simeq \mathfrak{A}_4$.

- G is generated by

$$\chi_r, (\zeta_3^2 x_2, \zeta_3 x_3 - x_2), (x_1, x_3, -x_2), \quad r \in \mathbb{Z}_{\geq 0},$$

- or G is generated by

$$\chi_r, (\zeta_3^{m+1} \zeta_3^2 x_2, \zeta_3^{m+1} (\zeta_3 x_3 - x_2)), (x_1, x_3, -x_2), \quad r, m \in \mathbb{Z}_{\geq 0}.$$

- $\bar{G} \simeq \mathfrak{S}_4$. G is generated by

$$\begin{aligned} \chi_r, (x_1, \zeta_{2^{m+1}}\zeta_4(-2x_2 + s_1x_3), \zeta_{2^{m+1}}\zeta_4(s_2x_2 + 2x_3)), \\ (x_1, -s_2x_2 - x_3, s_1x_2 + (s_1 + 1)x_3), \end{aligned}$$

where

$$s_1 = \zeta_8^3 + \zeta_8 - 1, \quad s_2 = \zeta_8^3 + \zeta_8 + 1, \quad r, m \in \mathbb{Z}_{\geq 0}.$$

- $\bar{G} \simeq \mathfrak{A}_5$. G is generated by

$$\chi_r, (x_1, (\zeta_5^3 + \zeta_5^4)x_2 - (\zeta_5^4 + 1)x_3, \zeta_5^3x_2 + (\zeta_5^2 + \zeta_5)x_3), (x_1, \zeta_5^2x_3, -\zeta_5^3x_2),$$

for $r \in \mathbb{Z}_{\geq 0}$.

Moreover, different subgroups in the list are not conjugated in PGL_3 .

From this description, we obtain:

Corollary 3.3. *If a group G gives rise to actions of different classes among (I), (T), and (P), then $G \simeq C_3^2$, the actions are of type (I) and (T), and there are two subgroups in PGL_3 isomorphic to G , up to conjugation, generated by*

$$\begin{aligned} \langle \mathrm{diag}(1, \zeta_3, 1), \mathrm{diag}(1, 1, \zeta_3) \rangle \quad \text{of Type (I),} \\ \langle \mathrm{diag}(1, \zeta_3, \zeta_3^2), (x_3, x_1, x_2) \rangle \quad \text{of Type (T).} \end{aligned}$$

Proof. We start by describing the center $Z(G)$ if G is in class

- (I): if G is nonabelian, the quotient $G/Z(G)$ is a noncyclic subgroup of PGL_2 , and $Z(G) \supset C_2$ unless $G = C_r \times \mathfrak{D}_n$, with n, r odd.
- (T): the center $Z(G) = 1, C_3$, or C_3^2 .
- (P): the center $Z(G) = 1$, unless $G = C_3^2 \rtimes C_4$, where $Z(G) = C_2$ and $G/Z(G) = C_3 \rtimes \mathfrak{S}_3$.

We conclude that no group giving rise to actions in type (P) also gives rise to an action of type (I) or (T). Assume that G gives rise to actions of both types (I) and (T). If G is nonabelian, we know that $G = C_3 \times \mathfrak{D}_n$ with n odd. But no group giving rise to actions of type (T) is isomorphic to this group. It follows that G is abelian and $G = Z(G)$. So $|G|$ is a power of 3. It is not hard to see that the only possibility is $G = C_3^2$ and it has two conjugacy classes in PGL_3 , as is explained above. $\square$

Remark 3.4. Finite isomorphic subgroups of $\mathrm{PGL}_2 \simeq \mathrm{Aut}(\mathbb{P}^1)$ are conjugated, see, e.g., [Bea10, Proposition 4.1]. Corollary 3.3 shows that this does not hold for PGL_3 .

We proceed to give a description of conjugacy classes of finite subgroups in PGL_3 .

Corollary 3.5. *Let G_1 and G_2 be isomorphic finite nonabelian subgroups of PGL_3 . Then G_1 and G_2 are conjugated in PGL_3 unless*

$$G_1 = \langle \mathrm{diag}(\zeta_n, 1, 1), (x_1, \zeta_3^2 x_2, \zeta_3 x_3 - x_2), (x_1, x_3, -x_2) \rangle,$$

$$G_2 = \langle \mathrm{diag}(\zeta_n, 1, 1), (x_1, x_2, \zeta_3^2 x_3 - \zeta_3 x_2), (x_1, x_3, -x_2) \rangle,$$

for $n \geq 1$. In this case, G_1 and G_2 are not conjugated in PGL_3 , and

$$G_1 \simeq G_2 \simeq \begin{cases} C_n \rtimes \mathfrak{A}_4 & \text{when } n \text{ is even,} \\ C_{2n} \rtimes \mathfrak{A}_4 & \text{when } n \text{ is odd.} \end{cases}$$

Proof. By Corollary 3.3, we know that nonabelian groups giving rise to actions of different types among **(I)**, **(T)**, and **(P)** are not isomorphic. Each subgroup in type **(T)** and **(P)** is unique in PGL_3 , up to conjugation. It follows that G_1 and G_2 give rise to actions of type **(I)**. Using Proposition 3.2, we check that the only possibility is when $\bar{G} = G/Z(G) \simeq \mathfrak{A}_4$, with the generators given above.

To show G_1 and G_2 are not conjugated in PGL_3 , we note that when $n = 1$, the generic stabilizer of the line $\{x_2 = (2\zeta_3 + 1)x_3\} \subset \mathbb{P}_{x_1, x_2, x_3}^2$ is trivial under the action induced by G_1 , but is C_3 under that induced by G_2 . $\square$

4. PRIMITIVE ACTIONS

Equivariant birational geometry of *primitive* actions was essentially settled, via equivariant MMP, in [Sak19]; the classification of conjugacy of actions was not addressed. We proceed to fill this gap:

- super-rigid, in particular, the actions are not conjugated in Cr_2 :
 - $\mathfrak{A}_6$: four actions on $\mathbb{P}^2$ [Che14, Theorem B.9],
 - $\mathrm{PSL}_2(\mathbb{F}_7)$: two actions on $\mathbb{P}^2$ [Che14, Theorem B.8],
 - $\mathrm{PSU}_2(\mathbb{F}_3)$: two actions on $\mathbb{P}^2$ [Sak19, Proposition 3.16],
 - $\mathrm{ASL}_2(\mathbb{F}_3)$: two actions on $\mathbb{P}^2$ [Sak19, Proposition 3.16],
- rigid:
 - $\mathfrak{A}_5$: two actions on $\mathbb{P}^2$, conjugated in Cr_2 and $\mathrm{Bir}^G(\mathbb{P}^2) \simeq \mathfrak{S}_5$ [Che14, Lemma B.13],
 - a subgroup $G = C_3^2 \rtimes C_4$ of the Hessian group: two actions on $\mathbb{P}^2$, conjugated in Cr_2 .

We explain the last entry: there are eight faithful 3-dimensional representations of the central extension $C_3.G$, but only two, up to multiplication with characters, thus there are two G -actions on $\mathbb{P}^2$. The

outer automorphism group $\text{Out}(G)$ is C_2^2 . In the automorphism group $\text{Aut}(G)$, there are three conjugacy classes of subgroups of order 72, isomorphic to

$$\mathfrak{F}_9, \quad \mathfrak{S}_3 \wr \mathfrak{S}_2, \quad \text{PSU}_3(\mathbb{F}_2).$$

Recall that we have inclusions $G \subset \text{PSU}_3(\mathbb{F}_2) \subset \text{PGL}_3$. In particular,

$$\text{Aut}^G(\mathbb{P}^2) = \text{PSU}_3(\mathbb{F}_2).$$

Then the conjugation action of $\text{PSU}_3(\mathbb{F}_2)$ on G induces a map

$$\text{PSU}_3(\mathbb{F}_2) \xrightarrow{f} C_2 \subset \text{Out}(G).$$

The conjugation in Cr_2 can be seen as follows: according to [Sak19, Proposition 3.16], there are exactly two orbits of size 6 on $\mathbb{P}^2$, for each of the actions. Blowing up any of these orbits, we obtain a cubic surface S , which must be the Fermat cubic surface [DI09, Table 4], yielding an embedding

$$G \hookrightarrow \text{Aut}(S) = C_3^3 \rtimes \mathfrak{S}_4.$$

There is a unique subgroup $G_1 \subset \text{Aut}(S)$ of order 72. We know that $G_1 \simeq \mathfrak{S}_3 \wr \mathfrak{S}_2$ and $G \subset G_1$. Choosing an element in $G_1 \setminus G$, we obtain an element $\tau_1 \in \text{Bir}^G(\mathbb{P}^2)$ such that $\langle \tau_1, G \rangle \simeq G_1$. Similarly, blowing up another orbit of length 6, we obtain an element $\tau_2 \in \text{Bir}^G(\mathbb{P}^2)$ such that $\langle \tau_2, G \rangle \simeq G_1$. Then

$$\text{Bir}^G(\mathbb{P}^2) = \langle \tau_1, \tau_2, \text{PSU}_3(\mathbb{F}_2) \rangle,$$

and $\langle \tau_1, \tau_2, G \rangle$ is an amalgamated product of two copies of G_1 with common subgroup G . In particular, $\text{Bir}^G(\mathbb{P}^2)$ is infinite.

The conjugation action of G_1 on G induces a map

$$\mathfrak{S}_3 \wr \mathfrak{S}_2 \xrightarrow{g} C_2 \subset \text{Out}(G),$$

and the images of f and g generate $\text{Out}(G)$, since $\mathfrak{S}_3 \wr \mathfrak{S}_2$ and $\text{PSU}_3(\mathbb{F}_2)$ are distinct index-2 subgroups of $\text{Aut}(G)$. It follows that the two linear actions of G are conjugated in Cr_2 .

5. TRANSITIVE NON-PRIMITIVE ACTIONS

In this section, we assume that $G \subset \text{PGL}_3$ is one of the subgroups listed in Proposition 3.1. Observe that the points

$$(5.1) \quad [1 : 0 : 0], \quad [0 : 1 : 0], \quad [0 : 0 : 1]$$

form a G -orbit of length 3, and that

$$\iota \in \text{Bir}^G(\mathbb{P}^2),$$

where ι is the standard Cremona involution (1.1).

If $G \simeq \mathfrak{A}_4$, then $\text{Aut}^G(\mathbb{P}^2) \simeq \mathfrak{S}_4$ and

$$\bar{\beta}(\text{Aut}^G(\mathbb{P}^2)) = \text{Out}(G) \simeq C_2.$$

Similarly, if $G \simeq \mathfrak{S}_4$, then $\text{Aut}^G(\mathbb{P}^2) = G$ and $\text{Out}(G)$ is trivial. Hence, in these two cases, there is a unique G -action on $\mathbb{P}^2$.

Lemma 5.1 (Pinardin). *Suppose that $G \simeq \mathfrak{S}_4$. Then*

$$\text{Bir}^G(\mathbb{P}^2) = \langle G, \iota \rangle \simeq \mathfrak{S}_4 \times C_2.$$

Proof. Observe that the points

$$(5.2) \quad [1 : 1 : 1], \quad [-1 : 1 : 1], \quad [1 : -1 : 1], \quad [1 : 1 : -1],$$

form a G -orbit of length 4. Then (5.1) and (5.2) are the only G -orbits in $\mathbb{P}^2$ of length < 9 that are in general position. Thus, every G -Sarkisov link that starts at $\mathbb{P}^2$ must blow up one of these two orbits.

Let $\sigma: \tilde{S} \rightarrow \mathbb{P}^2$ be the blow up of the G -orbit (5.2). Then $\tilde{S}$ is a del Pezzo surface of degree 4, and we have the following G -Sarkisov link:

$$\begin{array}{ccc} & \tilde{S} & \\ \sigma \swarrow & & \searrow \tilde{\pi} \\ \mathbb{P}^2 & & \mathbb{P}^1 \end{array}$$

where $\tilde{\pi}$ is a conic bundle. The kernel of the G -action on $\mathbb{P}^1$ is isomorphic to C_2^2 , which implies that there are no G -Sarkisov links that start at $\tilde{S}$ except for the inverse of the constructed G -Sarkisov link, which brings us back to $\mathbb{P}^2$. The required assertion follows from the classification of equivariant Sarkisov links starting at $\mathbb{P}^2$. $\square$

Lemma 5.2. *Suppose that $G \simeq \mathfrak{A}_4$. For $\lambda \in k \setminus \{0, 1\}$ with $\lambda^6 \neq -1$, let ϑ_λ be the birational selfmap of $\mathbb{P}^2$ given by*

$$\vartheta_\lambda : (x_1, x_2, x_3) \mapsto (f_1, f_2, f_3),$$

where

$$\begin{aligned} f_1 &= x_1^4 x_3 - \frac{\lambda^{12} + 1}{\lambda^2} x_1^2 x_2^2 x_3 + \lambda^8 x_2^4 x_3 - 2\lambda^2 x_1^2 x_3^3 - 2\lambda^6 x_2^2 x_3^3 + \lambda^4 x_3^5, \\ f_2 &= \lambda^8 x_1^4 x_2 - 2\lambda^6 x_1^2 x_2^3 + \lambda^4 x_2^5 - \frac{\lambda^{12} + 1}{\lambda^2} x_1^2 x_2 x_3^2 - 2\lambda^2 x_2^3 x_3^2 + x_2 x_3^4, \\ f_3 &= \lambda^4 x_1^5 - 2\lambda^2 x_1^3 x_2^2 + x_1 x_2^4 - 2\lambda^6 x_1^3 x_3^2 - \frac{\lambda^{12} + 1}{\lambda^2} x_1 x_2^2 x_3^2 + \lambda^8 x_1 x_3^4. \end{aligned}$$

Then ϑ_λ is an involution in $\text{Bir}^G(\mathbb{P}^2)$, so, in particular, $\text{Bir}^G(\mathbb{P}^2)$ is infinite. Moreover, birational involutions ϑ_λ , the standard Cremona involution ι , and $\text{Aut}^G(\mathbb{P}^2) \simeq \mathfrak{S}_4$ generate the group $\text{Bir}^G(\mathbb{P}^2)$.

Proof. Observe that $\mathbb{P}^2$ contains one G -orbit of length 3, three G -orbits of length 4, and infinitely many G -orbits of length 6 such that each of them is a G -orbit of $[0 : 1 : \lambda]$, for $\lambda \in k^\times$. Let $\sigma_\lambda : S_\lambda \rightarrow \mathbb{P}^2$ be the blow up of the G -orbit of the point $[0 : 1 : \lambda]$. Then S_λ is a smooth cubic surface if and only if $\lambda \neq 0, 1$ and $\lambda^6 \neq -1$. In this case, we have the following G -commutative diagram:

$$\begin{array}{ccc} S_\lambda & \xrightarrow{\theta_\lambda} & S_\lambda \\ \sigma_\lambda \downarrow & & \downarrow \sigma_\lambda \\ \mathbb{P}^2 & \xrightarrow{\vartheta_\lambda} & \mathbb{P}^2 \end{array}$$

where θ_λ is a biregular involution in $\text{Aut}(S_\lambda)$. Note also that

$$\langle G, \vartheta_\lambda \rangle \simeq \mathfrak{S}_4.$$

Now, arguing as in the proof of Lemma 5.1, we see that $\text{Bir}^G(\mathbb{P}^2)$ is generated by $\text{Aut}^G(\mathbb{P}^2)$, the involution ι , and involutions ϑ_λ . $\square$

Recall from [Sak19, Theorem 1.3] that $\mathbb{P}^2$ is G -birationally rigid if and only if $G \not\simeq \mathfrak{A}_4, \mathfrak{S}_4$. Moreover, if

$$G \not\simeq \mathfrak{A}_4, \quad \mathfrak{S}_4, \quad C_3^2, \quad C_3 \rtimes \mathfrak{S}_3, \quad C_7 \rtimes C_3,$$

then it follows from [Sak19, Lemma 3.14] that (5.1) is the only G -orbit in $\mathbb{P}^2$ of length 3. From the proof of [Sak19, Theorem 1.3], we know that $\text{Bir}^G(\mathbb{P}^2)$ is finite, and that

$$(5.3) \quad \text{Bir}^G(\mathbb{P}^2) = \langle \text{Aut}^G(\mathbb{P}^2), \iota \rangle.$$

If $G \simeq C_3 \rtimes \mathfrak{S}_3$ or $G \simeq C_3^2$, then it follows from [Sak19, Lemma 3.14] that $\mathbb{P}^2$ has 4 orbits of length 3 (the group $C_3 \rtimes \mathfrak{S}_3$ is erroneously omitted in [Sak19, Lemma 3.14]). In these two cases, we have

$$\text{Aut}^G(\mathbb{P}^2) \simeq \text{ASL}_2(\mathbb{F}_3),$$

and all G -orbits of length 3 form one orbit of $\text{Aut}^G(\mathbb{P}^2)$. It follows from [Sak19, Theorem 1.3] that (5.3) also holds.

Lemma 5.3. *Suppose $G \simeq C_3^2$ or $C_3 \rtimes \mathfrak{S}_3$. Then $\text{Bir}^G(\mathbb{P}^2)$ is infinite.*

Proof. Set $\hat{G} = \text{Bir}^G(\mathbb{P}^2)$. If $\hat{G}$ is finite, its birational action on $\mathbb{P}^2$ is regularized on a smooth projective rational surface S such that

- S is a smooth del Pezzo surface with $\text{rkPic}(S)^{\hat{G}} = 1$, or

- S admits a $\hat{G}$ -equivariant conic bundle

$$\pi : S \rightarrow \mathbb{P}^1,$$

$$\text{with } \text{rk Pic}(S)^{\hat{G}} = 2.$$

The latter case is impossible, since $\mathbb{P}^2$ is G -solid [Sak19], so that S is a smooth del Pezzo surface. Using the classification of finite subgroups of automorphism groups of smooth del Pezzo surfaces [DI09], we conclude that $S \simeq \mathbb{P}^2$. Since $\text{Aut}^G(\mathbb{P}^2) \simeq \text{ASL}_2(\mathbb{F}_3)$, it follows from Corollary 3.3 that $\hat{G}$ is a primitive subgroup in $\text{Aut}(S) \simeq \text{PGL}_3$, which is a contradiction since $\text{ASL}_2(\mathbb{F}_3)$ is a maximal subgroup in PGL_3 . $\square$

If $G \simeq C_7 \rtimes C_3$, then $\text{Aut}^G(\mathbb{P}^2) = G$, and (5.1) is the only G -orbit in $\mathbb{P}^2$ of length 3. In this case, it follows from the proof of [Sak19, Theorem 1.3] that

$$\text{Bir}^G(\mathbb{P}^2) = \langle G, \iota, \tau_1, \tau_2, \tau_3 \rangle,$$

where τ_1, τ_2, τ_3 are Geiser involutions, implicitly described in the proof of [Sak19, Lemma 3.14]. One can show that the group $\langle \iota, \tau_1, \tau_2, \tau_3 \rangle$ is infinite, which gives:

Corollary 5.4. *If $G \simeq C_7 \rtimes C_3$, then $\text{Bir}^G(\mathbb{P}^2)$ is infinite.*

If $G \simeq \mathfrak{A}_4$ or $G \simeq \mathfrak{S}_4$, then $\bar{\beta}(\iota)$ is the identity, so that conjugation by ι does not change the G -action, which is not very surprising, since $\mathfrak{A}_4$ and $\mathfrak{S}_4$ admit one faithful action on $\mathbb{P}^2$. In all other cases, it follows from Remark 2.3 that

$$\bar{\beta}(\text{Bir}^G(\mathbb{P}^2)) = \langle \bar{\beta}(\text{Aut}^G(\mathbb{P}^2)), \bar{\beta}(\iota) \rangle,$$

where ι is the standard Cremona involution. We proceed to show how to compute $\bar{\beta}(\text{Bir}^G(\mathbb{P}^2))$ in several representative examples.

Example 5.5. Suppose that $G \simeq C_3^2$. Then $\text{Out}(G) \simeq \text{GL}_2(\mathbb{F}_3)$, the group $\text{Aut}^G(\mathbb{P}^2)$ is the Hessian subgroup $\text{ASL}_2(\mathbb{F}_3) \subset \text{PGL}_3$ and

$$\bar{\beta}(\text{Aut}^G(\mathbb{P}^2)) \simeq \text{SL}_2(\mathbb{F}_3),$$

and we have two non-isomorphic G -actions on $\mathbb{P}^2$. These actions are conjugated in Cr_2 , because $\bar{\beta}(\iota) \notin \bar{\beta}(\text{Aut}^G(\mathbb{P}^2))$. Thus,

$$\bar{\beta}(\text{Bir}^G(\mathbb{P}^2)) = \text{Out}(G),$$

and there is a unique imprimitive G -action up to conjugation in Cr_2 .

Example 5.6. Suppose that $G \simeq C_3 \rtimes \mathfrak{S}_3$. Then $\text{Aut}^G(\mathbb{P}^2) \simeq \text{ASL}_2(\mathbb{F}_3)$, $\text{Out}(G) \simeq \mathfrak{S}_4$, and

$$\bar{\beta}(\text{Aut}^G(\mathbb{P}^2)) \simeq \mathfrak{A}_4.$$

Therefore, there are two non-isomorphic G -actions on $\mathbb{P}^2$. They are conjugated in Cr_2 , because $\bar{\beta}(\iota) \notin \bar{\beta}(\text{Aut}^G(\mathbb{P}^2))$. Then

$$\bar{\beta}(\text{Bir}^G(\mathbb{P}^2)) = \text{Out}(G).$$

Example 5.7. Suppose that $G \simeq C_7 \rtimes C_3$. Then $\text{Out}(G) \simeq C_2$ and

$$\text{Aut}^G(\mathbb{P}^2) = G.$$

Hence, there are two non-isomorphic G -actions on $\mathbb{P}^2$, conjugated in Cr_2 by ι , and

$$\bar{\beta}(\text{Bir}^G(\mathbb{P}^2)) = \text{Out}(G),$$

since $\beta(\iota)$ is not an inner automorphism of G .

Example 5.8. In the notation of Proposition 3.1, suppose further that $\nu(G) = \mathfrak{S}_3$ and $G \simeq C_n^2 \rtimes \mathfrak{S}_3$ with $n \geq 3$. Then

$$\text{Aut}^G(\mathbb{P}^2) = G,$$

and $\text{Out}(G) \simeq (\mathbb{Z}/n\mathbb{Z})^\times$, and we have $|(\mathbb{Z}/n\mathbb{Z})^\times|$ non-isomorphic G -actions on $\mathbb{P}^2$. Then

$$\bar{\beta}(\text{Bir}^G(\mathbb{P}^2)) \simeq C_2,$$

since $\beta(\iota)$ is not an inner automorphism.

In all these examples, the conjugation by ι changes the action of the group G on $\mathbb{P}^2$. This is not a coincidence:

Lemma 5.9. *Suppose that $G \not\simeq \mathfrak{A}_4, \mathfrak{S}_4$. Then $\bar{\beta}(\iota) \notin \bar{\beta}(\text{Aut}^G(\mathbb{P}^2))$.*

Proof. Recall that G is one of the subgroups in Proposition 3.1. Let us use notation introduced in this proposition. Keeping in mind examples above, we may assume further that $G \not\simeq C_3^2, C_3 \rtimes \mathfrak{S}_3$. Then the G -orbit (5.1) is the only G -orbit of length 3 in $\mathbb{P}^2$, so that (5.1) is also an orbit of the group $\text{Aut}^G(\mathbb{P}^2)$.

Denote by $T \subseteq G$ the subgroup consisting of transformations given by diagonal matrices. Then G is generated by T , σ_{123} and in some cases by σ_{12} , where all possibilities for T are given by Proposition 3.1. Conjugation by ι gives an automorphism $\phi \in \text{Aut}(G)$ that acts trivially on σ_{123} and σ_{12} (if $\sigma_{12} \in G$), but

$$\phi(t) = \frac{1}{t},$$

for every $t \in T$. Hence, ϕ is not trivial, since $G \not\simeq \mathfrak{A}_4, \mathfrak{S}_4$.

Let $G' = \text{Aut}^G(\mathbb{P}^2)$, and let $T' \subseteq G'$ be the subgroup containing all transformations given by diagonal matrices. Note that $\sigma_{123} \in G'$, since $G \subset G'$. Similarly, if $\sigma_{12} \in G$, then $\sigma_{12} \in G'$, so that

$$G' = \langle T', \sigma_{123}, \sigma_{12} \rangle.$$

If $\sigma_{12} \notin G$, then it follows from the proof of Proposition 3.1 that

$$G' = \langle T', \sigma_{123} \rangle,$$

unless $\nu(G) = C_3$ and $G \simeq (C_n \times C_{n/3}) \rtimes C_3$ is generated by

$$\text{diag}(\zeta_n^3, 1, 1), \quad \text{diag}(\zeta_n^2, \zeta_n, 1), \quad \sigma_{123},$$

where $3 \mid n$. In this case, $\sigma_{12} \in G'$, and $G' = \langle T', \sigma_{123}, \sigma_{12} \rangle$.

Now, we suppose that ϕ is given by a conjugation by some $g \in G'$. Then $g = t'\sigma$ for some $\sigma \in \langle \sigma_{123}, \sigma_{12} \rangle$. Note that σ is not an identity, since conjugation by an element in T' acts trivially on T . If $g = t'\sigma_{123}$, then

$$\frac{1}{t} = \phi(t) = t'\sigma_{123}t(t'\sigma_{123})^{-1} = \sigma_{123}t(\sigma_{123})^{-1},$$

for every $t \in T$, which is a contradiction, since $\sigma_{123}t(\sigma_{123})^{-1} = t$ if and only if t is an identity, and T is nontrivial. Similarly, if $g = t'\sigma_{12}$, then

$$\frac{1}{t} = \phi(t) = t'\sigma_{12}t(t'\sigma_{12})^{-1} = \sigma_{12}t(\sigma_{12})^{-1},$$

for every $t \in T$. This implies that every element in T is given by the diagonal matrix

$$\text{diag}(a, a^{-1}, 1),$$

for some $a \in k^\times$, which is only possible when $G \simeq C_3^2$. But this case is treated in Example 5.5 and excluded, by assumption. Similarly, we obtain a contradiction for other possibilities for $\sigma \in \langle \sigma_{123}, \sigma_{12} \rangle$. $\square$

Summarizing, we obtain:

Theorem 5.10. *Let $\phi_1, \phi_2 : G \hookrightarrow \text{PGL}_3$ be different imprimitive actions such that $\phi_1(G) = \phi_2(G)$ is one of the subgroups in Proposition 3.1. Then ϕ_1 and ϕ_2 are conjugated in Cr_2 if and only if they are conjugated by the standard Cremona involution ι . Moreover, if $G \not\simeq \mathfrak{A}_4, \mathfrak{S}_4$, then conjugation by ι changes the isomorphism class of the G -action on $\mathbb{P}^2$.*

6. INTRANSITIVE

In this section, we prove Theorem 1.1. We assume that $G \subset \mathrm{PGL}_3$ is one of the subgroups listed in Proposition 3.2. Recall that the G -action on $\mathbb{P}^2 = \mathbb{P}(V)$ arises from a faithful linear representation of G , which decomposes as $V = \mathbf{1} \oplus V_2$. Let C_t be the generic stabilizer of the G -action on the line $\mathfrak{l} = \mathbb{P}(V_2) \subset \mathbb{P}(V)$, and χ the character of C_t in the normal bundle of $\mathfrak{l}$.

If $t = 1$, for two such G -actions $\mathbb{P}(V)$ and $\mathbb{P}(V')$ arising from

$$V = \mathbf{1} \oplus V_2, \quad V' = \mathbf{1} \oplus V'_2,$$

we have G -equivariant birationalities

$$\mathbb{P}(V) \sim \mathbb{P}(\mathbf{1} \oplus \mathbf{1}) \times \mathbb{P}(V_2) \sim \mathbb{P}(V'_2) \times \mathbb{P}(V_2) \sim \mathbb{P}(\mathbf{1} \oplus \mathbf{1}) \times \mathbb{P}(V'_2) \sim \mathbb{P}(V'),$$

by the no-name lemma. In fact, if G is nonabelian, we know that $G = \mathfrak{D}_n$, with odd n , when $t = 1$.

If $t \geq 2$, then the class of the action

$$(6.1) \quad [\mathbb{P}(V) \hookrightarrow G] \in \mathrm{Burn}_2^{\mathrm{inc}}(G)$$

equals

$$(6.2) \quad (C_t, \bar{G} \curvearrowright k(\mathfrak{l}), (\chi)) + (C_t, \bar{G} \curvearrowright k(\mathfrak{l}), (-\chi)),$$

where $\bar{G} = G/C_t$ acts generically freely on $\mathfrak{l} = \mathbb{P}(V_2)$. These symbols are incompressible, provided $\bar{G}$ is nonabelian. The main result in this section is

Theorem 6.1 (cf. Theorem 1.1). *Intransitive G -actions with $t \geq 2$ and G nonabelian are birational $\iff$ the corresponding $\bar{G}$ -actions on the invariant line $\mathfrak{l}$ are isomorphic and the corresponding characters χ are equal, up to ± 1 .*

Proof of $(\Leftarrow)$. Assume that the $\bar{G}$ -action and χ do not satisfy the assumption. Then the incompressible symbols (6.2) in the classes (6.1) of the two G -actions are different in $\mathrm{Burn}_2^{\mathrm{inc}}(G)$. The Burnside formalism implies that the two actions are not birational, see also [TYZ24, Section 7]. $\square$

The rest of this section is devoted to a proof of the $(\Rightarrow)$ direction of Theorem 6.1. The proof of Theorem 6.1 implies Theorem 1.1, namely, that the Burnside formalism is decisive for nonabelian intransitive actions. It suffices to identify actions with the same $C_t, \pm\chi$, and the same $\bar{G}$ -action on $\mathbb{P}^1$. We do this via a case-by-case analysis of $\bar{G}$, using the list of all possibilities for G in Proposition 3.2.

Remark 6.2. When $\bar{G} = \mathfrak{A}_4$ or $\mathfrak{S}_4$, there is a unique $\bar{G}$ -action on $\mathbb{P}^1$. Theorem 6.1 then implies that G -actions are not birational if and only if the corresponding characters χ do not differ by ± 1 .

Remark 6.3. For intransitive $G \subset \mathrm{PGL}_3$, the group $\mathrm{Bir}^G(\mathbb{P}^2)$ is always infinite – it contains a 1-dimensional torus generated by

$$\{\mathrm{diag}(1, a, a) : a \in k^\times\}.$$

6.1. When $\bar{G} = C_n$. In this case, G is abelian. Linear G -actions are birational if and only if the corresponding invariants in [RY02] coincide. In fact, birational classification of linear actions of *abelian* groups has been settled, in all dimensions, in [RY02, Theorem 7.1].

6.2. When $\bar{G} = \mathfrak{D}_n$ with odd n . Recall that all G -actions are conjugated in Cr_2 if the generic stabilizer C_t of the G -action on $\mathbb{P}(V_2)$ is trivial. Hence, we assume that $t \geq 2$. In each case, the standard Cremona involution ι is contained in $\mathrm{Bir}^G(\mathbb{P}^2)$. It would be interesting to describe generators of $\mathrm{Bir}^G(\mathbb{P}^2)$.

Proof of Theorem 6.1 in this case. By Proposition 3.2, the action of G depends on the choice of primitive weights t_1, t_2, t_3 in

$$\mathrm{diag}(1, \zeta_r^{t_1}, \zeta_r^{t_1}), \quad \mathrm{diag}(1, \zeta_{2n}^{t_2}, \zeta_{2n}^{-t_2}), \quad (x_1, \zeta_{2^m}^{t_3} x_3, \zeta_{2^m}^{t_3} x_2).$$

The action of the generic stabilizer of $\mathfrak{l} = \{x = 0\} \subset \mathbb{P}^2$ is generated by

$$\mathrm{diag}(1, \zeta_r^{t_1}, \zeta_r^{t_1}), \quad \mathrm{diag}(1, \zeta_{2^{m-1}}^{t_3}, \zeta_{2^{m-1}}^{t_3}).$$

The weights can be changed birationally as follows:

- The standard Cremona involution ι composed with transposition of x_2, x_3 preserves t_2 and inverts

$$t_1 \mapsto -t_1, \quad t_3 \mapsto -t_3,$$

- the transposition of x_2, x_3 preserves t_1, t_3 and inverts $t_2 \mapsto -t_2$,
- the coordinate change $x_3 \mapsto -x_3$ preserves t_1, t_2 and maps $t_3 \mapsto t_3 + 2^{m-1}$.

For two choices of weights (t_1, t_2, t_3) , if the corresponding characters χ differ by ± 1 and the induced $\mathfrak{D}_n$ -actions on $\mathfrak{l}$ are isomorphic, then the weights can be interchanged by a composition of the three maps above, and thus the actions are birational. $\square$

$$\begin{array}{ccc}
& S & \\
\pi \swarrow & & \searrow \phi \\
\mathbb{F}_1 & & \mathbb{F}_1 \\
\eta \downarrow & & \downarrow \eta \\
\mathbb{P}^2 & \dashrightarrow & \mathbb{P}^2
\end{array}$$

Observe that $\mathrm{Bir}^G(\mathbb{P}^2)$ also contains the birational selfmap of $\mathbb{P}^2$ given by

$$(6.3) \quad \gamma : (x_1, x_2, x_3) \mapsto (x_1 x_2^{n/2} x_3^{n/2}, x_2(x_2^n - x_3^n), x_3(x_2^n - x_3^n)).$$

$$\begin{array}{ccccccc}
& S_1 & & S_2 & & & S_r \\
& \swarrow \pi_1 & \searrow \phi_1 & \swarrow \pi_2 & \searrow \phi_2 & & \swarrow \pi_r \searrow \phi_r \\
\mathbb{F}_1 & & \mathbb{F}_{n-1} & & \mathbb{F}_{n-2} \cdots \mathbb{F}_3 & & \mathbb{F}_1 \\
\eta \downarrow & & & & & & \downarrow \eta \\
\mathbb{P}^2 & \text{---} & & & & \text{---} & \mathbb{P}^2 \\
& & & & \gamma & &
\end{array}$$

Proof of Theorem 6.1 in this case. In each of the families, the G -action depends on primitive weights t_1, t_2, t_3, t_4 as in

- $$\begin{aligned} (1) & \text{diag}(1, \zeta_r^{t_1}, \zeta_r^{t_1}), \quad \text{diag}(1, \zeta_{2n}^{t_2}, \zeta_{2n}^{-t_2}), \quad (x_1, \zeta_4 x_3, \zeta_4 x_2). \\ (2) & \text{diag}(1, \zeta_r^{t_1}, \zeta_r^{t_1}), \quad \text{diag}(1, \zeta_{m+1}^{t_2}, \zeta_{m+1}^{t_3} \zeta_{2n}^{-t_2}), \quad (x_1, \zeta_4 x_3, \zeta_4 x_2). \end{aligned}$$

$$(3) \text{ diag}(1, \zeta_r^{t_1}, \zeta_r^{t_1}), \quad \text{diag}(1, \zeta_{2n}^{t_2}, \zeta_{2n}^{-t_2}), \quad (x_1, \zeta_{2^{m+1}}^{t_4} \zeta_4 x_3, \zeta_{2^{m+1}}^{t_4} \zeta_4 x_2).$$

The generic stabilizer of $\mathfrak{l} = \{x_1 = 0\} \subset \mathbb{P}^2$ is generated by

$$\begin{aligned} (1) & \text{ diag}(1, \zeta_r^{t_1}, \zeta_r^{t_1}), \quad \text{diag}(1, -1, -1); \\ (2) & \text{ diag}(1, \zeta_r^{t_1}, \zeta_r^{t_1}), \quad \text{diag}(1, \zeta_{2^m}^{t_3}, \zeta_{2^m}^{t_3}); \\ (3) & \text{ diag}(1, \zeta_r^{t_1}, \zeta_r^{t_1}), \quad \text{diag}(1, \zeta_{2^m}^{t_4}, \zeta_{2^m}^{t_4}). \end{aligned}$$

The weights can be altered by the following maps, in all cases

- the birational map with image $(\frac{1}{x_1}, \frac{1}{x_3}, -\frac{1}{x_2})$ maps

$$t_1 \mapsto -t_1, \quad t_2 \mapsto t_2, \quad t_3 \mapsto -t_3, \quad t_4 \mapsto -t_4,$$

- the coordinate change $\text{diag}(1, 1, -1)$ maps

$$t_1 \mapsto t_1, \quad t_2 \mapsto t_2, \quad t_3 \mapsto t_3, \quad t_4 \mapsto t_4 + 2^m.$$

- the birational map γ maps

$$t_1 \mapsto t_1, \quad t_2 \mapsto t_2, \quad t_3 \mapsto t_3 + 2^m, \quad t_4 \mapsto t_4,$$

or equivalently,

$$t_1 \mapsto t_1, \quad t_2 \mapsto t_2 + n, \quad t_3 \mapsto t_3, \quad t_4 \mapsto t_4.$$

Similarly, for two actions, if the characters χ of the generic stabilizer of $\mathfrak{l}$ differ by ± 1 and the induced $\mathfrak{D}_n$ -actions on $\mathfrak{l}$ are isomorphic, we know that the corresponding weights can be interchanged by a composition of the three maps above; thus, the actions are birational. $\square$

To obtain an alternative proof of Theorem 6.1 in this case, one can explicitly describe generators of $\text{Bir}^G(\mathbb{P}^2)$: Recall that $\iota, \gamma \in \text{Bir}^G(\mathbb{P}^2)$. For every $\lambda \in k^\times$, with $\lambda^n \neq 1$, let $\tau_\lambda \in \text{Bir}^G(\mathbb{P}^2)$ be the birational map given by

$$(6.4) \quad \tau_\lambda : (x_1, x_2, x_3) \mapsto (x_1(x_2^{2n} - x_3^{2n}), x_2 f_\lambda, x_3 f_\lambda),$$

where

$$f_\lambda = \prod_{r=0}^{n-1} ((x_1 - \lambda \zeta_n^r x_2) \cdot (x_2 - \lambda \zeta_n^r x_1)).$$

Then τ_λ fits in the following G -commutative diagram:

$$\begin{array}{ccccccc} & & S' & & S'' & & S''' \\ & \swarrow \pi' & & \searrow \phi' & \swarrow \pi'' & & \searrow \phi'' \\ \mathbb{F}_1 & & & & \mathbb{F}_{2n-1} & & \mathbb{F}_{n-1} \\ & \swarrow \pi''' & & \searrow \phi''' & & & \\ & & & & & & \mathbb{F}_1 \\ \eta \downarrow & & & & & & \downarrow \eta \\ \mathbb{P}^2 & \xrightarrow{\quad \tau_\lambda \quad} & & & & & \mathbb{P}^2 \end{array}$$

where π' is the blowup of the orbit of length $2n$ on L that is mapped by χ to the G -orbit of $[0 : \lambda : 1]$, π'' and π''' are blowups of the orbits of length n on the strict transform of E , and ϕ' , ϕ'' and ϕ''' are contractions of the strict transforms of the fibers of the corresponding $\mathbb{P}^1$ -bundles passing through the blown up orbits.

Lemma 6.4. *Suppose $n \geq 10$. Then $\text{Bir}^G(\mathbb{P}^2)$ is generated by $\text{Aut}^G(\mathbb{P}^2)$, the standard Cremona transformation ι , the map γ given by (6.3), and τ_λ given by (6.4) for $\lambda \in k^\times$ with $\lambda^n \neq 1$.*

Proof. Take any $\varphi \in \text{Bir}^G(\mathbb{P}^2)$ such that $\varphi \notin \text{Aut}^G(\mathbb{P}^2)$. Then φ can be decomposed into a sequence of G -Sarkisov links. Since $n \geq 10$, $\mathbb{P}^2$ contains exactly two G -orbits of length < 9 : the G -fixed point, and the unique G -orbit of length 2, which is contained in the G -invariant line. Hence, the only G -Sarkisov links that start at $\mathbb{P}^2$ are the link

$$\begin{array}{ccc} & \mathbb{F}_1 & \\ \eta \swarrow & & \searrow \\ \mathbb{P}^2 & & \mathbb{P}^1 \end{array}$$

where $\mathbb{F}_1 \rightarrow \mathbb{P}^1$ is the natural projection, and the link

$$\begin{array}{ccc} & Y & \\ \swarrow & & \searrow \\ \mathbb{P}^2 & & \mathbb{P}^1 \times \mathbb{P}^1 \end{array}$$

where the left map is the blow up of the unique G -orbit of length 2 and the right map is the contraction of the strict transform of the G -invariant line $\mathfrak{l} \subset \mathbb{P}^2$. Moreover, the G -fixed point in $\mathbb{P}^1 \times \mathbb{P}^1$ is the only G -orbit of length < 8 in $\mathbb{P}^1 \times \mathbb{P}^1$, so the only G -Sarkisov link that starts at $\mathbb{P}^1 \times \mathbb{P}^1$ is the inverse of the link described above.

Moreover, since $C_t \neq 1$, any G -orbit of a point in $\mathbb{F}_1$ away from $E \cup L$ intersects some fiber of the natural projection $\mathbb{F}_1 \rightarrow \mathbb{P}^1$ at more than 1 point. Now, using the classification of two-dimensional G -Sarkisov

links [Isk96], we see that we have the following G -commutative diagram

$$\begin{array}{ccccc}
 & \mathbb{F}_1 & \overset{\rho}{\dashrightarrow} & \mathbb{F}_1 & \\
 & \downarrow & & \downarrow & \\
 \eta \swarrow & \mathbb{P}^1 & \xrightarrow{v} & \mathbb{P}^1 & \searrow \eta \\
 & \downarrow & & \downarrow & \\
 \mathbb{P}^2 & \dashrightarrow & \varphi & \dashrightarrow & \mathbb{P}^2
 \end{array}$$

where v is an isomorphism, and ρ is a G -birational map such that its indeterminacy points are G -orbits in $E \cup L$, and every fiber of the projection $\mathbb{F}_1 \rightarrow \mathbb{P}^1$ contains at most one of these indeterminacy points. Composing φ with a composition of appropriate maps τ_λ or $\iota \circ \tau_\lambda \circ \iota$, we may assume that the indeterminacy points of the map ρ are G -orbits of length 2 or n . Composing φ with γ or $\iota \circ \gamma \circ \iota$, we may further assume that the indeterminacy points of ρ are G -orbits of length 2. Hence, either $\rho \in \text{Aut}^G(\mathbb{P}^2)$ or $\rho \circ \iota \in \text{Aut}^G(\mathbb{P}^2)$, which implies the required assertion. $\square$

It seems possible to weaken (or remove) the condition $n \geq 10$ in Lemma 6.4; this would entail taking into account Geiser and Bertini involutions that may appear in $\text{Bir}^G(\mathbb{P}^2)$.

6.4. When $\bar{G} = \mathfrak{A}_4$. Recall that there is a unique $\mathfrak{A}_4$ -action on $\mathbb{P}^1$. We find birational maps between two G -actions when the corresponding characters χ differ by ± 1 .

Proof of Theorem 6.1 in this case. The action depends on choices of primitive (with the exception that t_2 can be 0) weights t_1, t_2, t_3 in

- (1) $\text{diag}(1, \zeta_r^{t_1}, \zeta_r^{t_1}), (x_1, \zeta_3^{t_2} \zeta_3^2 x_2, \zeta_3^{t_2} (\zeta_3 x_3 - x_2));$
- (2) $\text{diag}(1, \zeta_r^{t_1}, \zeta_r^{t_1}), (x_1, \zeta_{3^{m+1}}^{t_3} \zeta_3^{t_2} \zeta_3^2 x_2, \zeta_{3^{m+1}}^{t_3} \zeta_3^{t_2} (\zeta_3 x_3 - x_2))$

with $m \geq 1$. The respective action of the generic stabilizer of $\mathfrak{l}$ is generated by

- (1) $\text{diag}(1, \zeta_r^{t_1}, \zeta_r^{t_1}), \quad \text{diag}(1, -1, -1);$
- (2) $\text{diag}(1, \zeta_r^{t_1}, \zeta_r^{t_1}), \quad \text{diag}(1, \zeta_{3^m}^{t_3}, \zeta_{3^m}^{t_3}), \quad \text{diag}(1, -1, -1).$

Let

$$\begin{aligned}
 h_4 &= x_2^4 - \frac{8\zeta_3 + 4}{3} x_2^3 x_3 - 2x_2^2 x_3^2 + \frac{8\zeta_3 + 4}{3} x_2 x_3^3 + x_3^4, \\
 f_4 &= x_2^3 x_3 + (-2\zeta_3 - 1)x_2^2 x_3^2 - x_2 x_3^3, \\
 f_6 &= x_2^6 - (4\zeta_3 + 2)x_2^5 x_3 - 5x_2^4 x_3^2 - 5x_2^2 x_3^4 + (4\zeta_3 + 2)x_2 x_3^5 + x_3^6.
 \end{aligned}$$

We note that f_4, h_4, g_6 are semi-invariant forms under the G -action. They correspond to orbits of length 4 and 6 of the $\mathfrak{A}_4$ -action on $\mathbb{P}^1$. Then

- The birational map $\sigma : (x_1, x_2, x_3) \mapsto (x_1 h_4, x_2 f_4, x_3 f_4)$ maps

$$t_1 \mapsto t_1, \quad t_2 \mapsto t_2 + 1, \quad t_3 \mapsto t_3,$$

or equivalently

$$t_1 \mapsto t_1, \quad t_2 \mapsto t_2, \quad t_3 \mapsto t_3 + 3^{m-1}.$$

- The birational map $\gamma : (x_1, x_2, x_3) \mapsto (f_6 h_4, x_1 x_2 f_4^2, x_1 x_3 f_4^2)$ maps

$$t_1 \mapsto -t_1, \quad t_2 \mapsto -t_2, \quad t_3 \mapsto -t_3.$$

For two G -actions, as long as the characters of the stabilizer of $\mathfrak{l}$ differ by ± 1 , we can interchange the corresponding weights t_1, t_2, t_3 of the two actions using maps above, and thus they are birational. $\square$

6.5. **When $\bar{G} = \mathfrak{S}_4$.** Similarly, there is a unique $\mathfrak{S}_4$ -action on $\mathbb{P}^1$. We find birational maps between two actions when their corresponding characters χ differ by ± 1 .

Proof of Theorem 6.1 in this case. The action depends on choices of primitive weights t_1, t_2, t_3 in

$$\text{diag}(1, \zeta_r^{t_1}, \zeta_r^{t_1}), \quad (x_1, \zeta_{2^{m+1}}^{t_2} \zeta_4^{t_3} (-2x_2 + s_1 x_3), \zeta_{2^{m+1}}^{t_2} \zeta_4^{t_3} (s_2 x_2 + 2x_3))$$

with $m \geq 0$. The corresponding action of the generic stabilizer of $\mathfrak{l}$ is generated by

- $\text{diag}(1, \zeta_r^{t_1}, \zeta_r^{t_1}), \quad \text{diag}(1, -1, -1), \quad \text{when } m = 1,$
- $\text{diag}(1, \zeta_r^{t_1}, \zeta_r^{t_1}), \quad \text{diag}(1, -\zeta_{2^m}^{t_2}, -\zeta_{2^m}^{t_2}), \quad \text{when } m \neq 1.$

Let f_6, f_8, f_{12}, h_{12} be semi-invariant binary forms of degree 6, 8, 12 and 12, under the $\mathfrak{S}_4$ -action on $\mathbb{P}_{x_2, x_3}^1$ generated by

$$(-2x_2 + s_1 x_3, s_2 x_2 + 2x_3), \quad (-s_2 x_2 - x_3, s_1 x_2 + (s_1 + 1)x_3)$$

such that f_{12} and h_{12} are linearly independent. Such forms are unique up to scalars. Then

- The birational map $\sigma : (x_1, x_2, x_3) \mapsto (x_1 h_{12}, x_2 f_{12}, x_3 f_{12})$ maps

$$t_1 \mapsto t_1, \quad t_2 \mapsto t_2 + 2^m, \quad t_3 \mapsto t_3,$$

or equivalently

$$t_1 \mapsto t_1, \quad t_2 \mapsto t_2, \quad t_3 \mapsto t_3 + 2.$$

- The birational map $\gamma : (x_1, x_2, x_3) \mapsto (f_8, x_1 x_2 f_6, x_1 x_3 f_6)$ maps

$$t_1 \mapsto -t_1, \quad t_2 \mapsto -t_2, \quad t_3 \mapsto -t_3.$$

When the characters of the stabilizer of $\mathfrak{l}$ in two actions differ by ± 1 , we can interchange the corresponding weights t_1, t_2, t_3 using the maps above, and thus the actions are birational. $\square$

6.6. **When $\bar{G} = \mathfrak{A}_5$.** Recall that the group G is generated by

$$\text{diag}(1, \zeta_r, \zeta_r)$$

and

(6.5)

$$(x_1, (\zeta_5^3 + \zeta_5^4)x_2 - (\zeta_5^4 + 1)x_3, \zeta_5^3x_2 + (\zeta_5^2 + \zeta_5)x_3), (x_1, \zeta_5^2x_3, -\zeta_5^3x_2).$$

In particular, there is a unique minimal lift of $\mathfrak{A}_5 \subset \text{PGL}_2$ to GL_2 given by (6.5). This minimal lift is isomorphic to $\text{SL}_2(\mathbb{F}_5)$. It has two faithful 2-dimensional representations, giving rise to 2 non-isomorphic $\mathfrak{A}_5$ -actions on $\mathbb{P}^1$.

Proof of Theorem 6.1 in this case. Fixing the $\mathfrak{A}_5$ -action on $\mathbb{P}^1$, we see that the isomorphism class of the G -action only depends on the weight of the generic stabilizer of $\mathfrak{l}$. The action of the stabilizer is generated by

- $\text{diag}(1, \zeta_r^{t_1}, \zeta_r^{t_1})$, if r is even,
- $\text{diag}(1, \zeta_r^{t_1}, \zeta_r^{t_1})$, $\text{diag}(1, -1, -1)$, if r is odd,

where t_1 indicates the weight. Let f_{12} , f_{20} and f_{30} be the $\text{SL}_2(\mathbb{F}_5)$ -invariant binary forms of degrees 12, 20 and 30 in variables x_2 and x_3 , which correspond to the unique orbits of lengths 12, 20 and 30 of the $\mathfrak{A}_5$ -action on $\mathbb{P}^1$. Then the birational map

$$\sigma : (x_1, x_2, x_3) \mapsto (f_{12}f_{20}, x_1x_2f_{30}, x_1x_3f_{30})$$

changes the weight $t_1 \mapsto -t_1$. We conclude that two actions are birational if and only if the characters of the stabilizer of $\mathfrak{l}$ differ by ± 1 . $\square$

This completes the proof of Theorem 6.1, and Theorem 1.1 follows as a corollary.

REFERENCES

- [BB00] L. Bayle and A. Beauville. Birational involutions of $\mathbb{P}^2$. *Asian J. Math.*, 4(1):11–17, 2000.
- [BB04] A. Beauville and J. Blanc. On Cremona transformations of prime order. *C. R., Math., Acad. Sci. Paris*, 339(4):257–259, 2004.
- [Bea10] A. Beauville. Finite subgroups of $\text{PGL}_2(K)$. In *Vector bundles and complex geometry. Conference on vector bundles in honor of S. Ramanan on the occasion of his 70th birthday, Madrid, Spain, June 16–20, 2008.*

- pages 23–29. Providence, RI: American Mathematical Society (AMS), 2010.
- [Bla07] J. Blanc. Finite abelian subgroups of the Cremona group of the plane. *C. R., Math., Acad. Sci. Paris*, 344(1):21–26, 2007.
 - [Bli18] H. F. Blichfeldt. Blichfeldt’s finite collineation groups. *Bull. Amer. Math. Soc.*, 24(10):484–487, 1918.
 - [Che14] I. A. Cheltsov. Two local inequalities. *Izv. Math.*, 78(2):375–426, 2014.
 - [CLdC13] S. Cantat, St. Lamy, and Y. de Cornulier. Normal subgroups in the Cremona group. *Acta Math.*, 210(1):31–94, 2013.
 - [CS16] I. Cheltsov and C. Shramov. *Cremona groups and the icosahedron*. Monogr. Res. Notes Math. Boca Raton, FL: CRC Press, 2016.
 - [dF04] T. de Fernex. On planar Cremona maps of prime order. *Nagoya Math. J.*, 174:1–28, 2004.
 - [DI09] I. V. Dolgachev and V. A. Iskovskikh. Finite subgroups of the plane Cremona group. In *Algebra, arithmetic, and geometry: in honor of Yu. I. Manin. Vol. I*, volume 269 of *Progr. Math.*, pages 443–548. Birkhäuser Boston, Boston, MA, 2009.
 - [HKT21] B. Hassett, A. Kresch, and Yu. Tschinkel. Symbols and equivariant birational geometry in small dimensions. In *Rationality of varieties*, volume 342 of *Progr. Math.*, pages 201–236. Birkhäuser/Springer, Cham, [2021] ©2021.
 - [Isk96] V. A. Iskovskikh. Factorization of birational maps of rational surfaces from the viewpoint of Mori theory. *Russ. Math. Surv.*, 51(4):585–652, 1996.
 - [KT22a] A. Kresch and Yu. Tschinkel. Equivariant birational types and Burnside volume. *Ann. Sc. Norm. Super. Pisa Cl. Sci. (5)*, 23(2):1013–1052, 2022.
 - [KT22b] A. Kresch and Yu. Tschinkel. Equivariant Burnside groups and representation theory. *Selecta Math. (N.S.)*, 28(4):Paper No. 81, 39, 2022.
 - [Lon16] A. Lonjou. Non-simplicity of the Cremona group over any field. *Ann. Inst. Fourier*, 66(5):2021–2046, 2016.
 - [LS24] St. Lamy and J. Schneider. Generating the plane Cremona groups by involutions. *Algebr. Geom.*, 11(1):111–162, 2024.
 - [NvdPT08] K. A. Nguyen, M. van der Put, and J. Top. Algebraic subgroups of $GL_2(\mathbb{C})$. *Indag. Math. (N.S.)*, 19(2):287–297, 2008.
 - [RY02] Z. Reichstein and B. Youssin. A birational invariant for algebraic group actions. *Pacific J. Math.*, 204(1):223–246, 2002.
 - [Sak19] D. Sakovics. G -birational rigidity of the projective plane. *Eur. J. Math.*, 5(3):1090–1105, 2019.
 - [TYZ24] Yu. Tschinkel, K. Yang, and Zh. Zhang. Equivariant birational geometry of linear actions. *EMS Surv. Math. Sci.*, 11(2):235–276, 2024.
 - [YY93] St. S.-T. Yau and Yu. Yu. Gorenstein quotient singularities in dimension three. *Mem. Amer. Math. Soc.*, 105(505):viii+88, 1993.
 - [Zim18] S. Zimmermann. The abelianization of the real Cremona group. *Duke Math. J.*, 167(2):211–267, 2018.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF EDINBURGH, UK

Email address: `I.Cheltsov@ed.ac.uk`

COURANT INSTITUTE, 251 MERCER STREET, NEW YORK, NY 10012, USA

Email address: `tschinkel@cims.nyu.edu`

SIMONS FOUNDATION, 160 FIFTH AVENUE, NEW YORK, NY 10010, USA

COURANT INSTITUTE, 251 MERCER STREET, NEW YORK, NY 10012, USA

Email address: `zhijia.zhang@cims.nyu.edu`