

Introduction

- **Topic**: Probability and random processes
- Lectures: AT3 on Wednesday and Friday at 12:10pm
- Office hours: by appointment (email) at JCMB 6321

mi4a (Probability) Lecture 1

• **Email**: j.m.figueroa@ed.ac.uk

The future was uncertain



The future is still uncertain



José Figueroa-O'Farrill

2/23

The universe is fundamentally uncertain!

There is no god of Algebra, but...



The mathematical study of Probability

Some notable names

- Gerolamo Cardano (1501-1576)
- Pierre de Fermat (1601-1665)
- Blaise Pascal (1623-1662)
- Christiaan Huygens (1629-1695)
- Jakob Bernoulli (1654-1705)
- Abraham de Moivre (1667-1754)
- Thomas Bayes (1702-1761)
- Pierre-Simon Laplace (1749-1827)
- Adrien-Marie Legendre (1752-1833)
- Andrei Markov (1866-1922)
- Andrei Kolmogorov (1903-1987)
- Claude Shannon (1916-2001)



What it is all about

Mathematical probability aims to formalise everyday sentences of the type:

"The chance of A is p"

where A is some "event" and p is some "measure" of the likelihood of occurrence of that event.

Example

"There is a 20% chance of snow."

"There is 5% chance that the West Antarctic Ice Sheet will collapse in the next 200 years."

"There is a low probability of Northern Rock having a liquidity problem."

Trials and outcomes

This requires introducing some language.

Definition/Notation

By a trial (or an experiment) we mean any process which has a well-defined set Ω of outcomes. Ω is called the sample space.

Example

Tossing a coin: $\Omega = \{H, T\}$. Tossing two coins: $\Omega = \{(H, H), (H, T), (T, H), (T, T)\}$.

Example (Rolling a (6-sided) die)

 $\Omega = \{ \bullet, \bullet, \bullet, \bullet, \bullet, \bullet, \bullet, \bullet \}.$

José Figueroa-O'Farrill

mi4a (Probability) Lecture 1

9/23

The language of sets

Let us consider subsets of a set Ω .

Definition

The **complement** of $A \subset \Omega$ is denoted $A^c \subset \Omega$:

 $\omega \in A^c \iff \omega \not\in A$

Clearly $(A^c)^c = A$.



"Events are what we assign a probability to"

Definition

An event A is a subset of Ω . We say that an event A has (not) occurred if the outcome of the trial is (not) contained in A.

Example

Tossing a coin and getting a head: $A = \{H\}$. Tossing two coins and getting at least one head: $A = \{(H, H), (H, T), (T, H)\}$. Rolling a die and getting an even number: $A = \{\bullet, \bullet, \bullet, \bullet\}$ Rolling two dice and getting a total of 5: $A = \{(\bullet, \bullet), (\bullet, \bullet), (\bullet, \bullet), (\bullet, \bullet)\}$

Warning (for infinite O)

Not all subsets of Ω need be events!

José Figueroa-O'Farrill

José Figueroa-O'Farrill



mi4a (Probability) Lecture 1

mi4a (Probability) Lecture 1

10/23

Definition

The **intersection** of A and B is denoted $A \cap B$:

 $\omega \in A \cap B \iff \omega \in A \text{ and } \omega \in B$

If $A \cap B = \emptyset$ we say A and B are **disjoint**.



Distributivity identities

Union and intersection obey distributive properties.

Theorem

Let $(A_i)_{i\in I}$ be a family of subsets of Ω indexed by some index set I and let $B\subset \Omega$. Then

$$\bigcup_{i\in I} (B\cap A_i) = B\cap \bigcup_{i\in I} A_i$$

and

$$\bigcap_{i \in I} (B \cup A_i) = B \cup \bigcap_{i \in I} A_i$$

José Figueroa-O'Farrill mi4a (Probability) Lecture 1

14 / 23

De Morgan's Theorem

Union and intersection are "dual" under complementation.

Theorem (De Morgan's)

Let $(A_i)_{i\in I}$ be a family of subsets of Ω indexed by some index set I. Then

$$\left(\bigcup_{i\in I}A_i\right)^c=\bigcap_{i\in I}A_i^c \qquad \textit{and} \qquad \left(\bigcap_{i\in I}A_i\right)^c=\bigcup_{i\in I}A_i^c$$

Remark

This shows that complementation together with either union or interesection is enough, since, e.g.,

$$A \cup B = (A^c \cap B^c)^c$$

Proof.

$$\begin{split} \omega \in \bigcap_{i \in I} \left(B \cup A_i \right) \iff \forall i \in I, \quad \omega \in B \cup A_i \\ \iff \forall i \in I, \quad \omega \in B \quad \text{or} \quad \omega \in A_i \\ \iff \omega \in B \quad \text{or} \quad \omega \in A_i \quad \forall i \in I \\ \iff \omega \in B \quad \text{or} \quad \omega \in \bigcap_{i \in I} A_i \\ \iff \omega \in B \cup \bigcap_{i \in I} A_i . \end{split}$$

The other equality is proved similarly.



Definition

The difference $A \setminus B = A \cap B^c$ and the symmetric difference $A \triangle B = (A \setminus B) \cup (B \setminus A)$.



Probability/Set theory dictionary

Notation	Set-theoretic language	Probabilistic language
Ω	Universe	Sample space
$\omega\in\Omega$	member of Ω	outcome
$A\subset \Omega$	subset of Ω	some outcome in A occurs
Ac	complement of A	no outcome in A occurs
$A \cap B$	intersection	Both A and B
$A \cup B$	union	Either A or B (or both)
$A \setminus B$	difference	A, but not B
$A \bigtriangleup B$	symmetric difference	Either A or B, but not both
Ø	empty set	impossible event
Ω	whole universe	certain event

Which subsets can be events?

- For finite Ω , any subset can be an event.
- For infinite Ω, it is not always sensible to allow all subsets to be events. (Trust me!)
- If A is an event, it seems reasonable that A^c is also an event.
- Similarly, if A and B are events, it seems reasonable that $A \cup B$ and $A \cap B$ should also be events.

In summary, the collection of events must be closed under complementation and pairwise union and intersection. By induction, it must also be closed under finite union and intersection: if A_1, \ldots, A_N are events, so should be $A_1 \cap A_2 \cap \cdots \cap A_N$ and $A_1 \cup A_2 \cup \cdots \cup A_N$.

The following example, shows that this is not enough.

Example

Alice and Bob play a game in which they toss a coin in turn. The winner is the first person to obtain H. Intuition says that the the person who plays first has an advantage. We would like to quantify this intuition. Suppose Alice goes first. She wins if and only if the first H turns out after an odd number of tosses. Let ω_i be the outcome $TT \cdots TH$. Then the event that Alice wins is

 $A = \{\omega_1, \omega_3, \omega_5, ...\}$, which is a disjoint union of a countably infinite number of events. In order to compute the likelihood of Alice winning, it had better be the case that A is an event, so one demands that the family of events be closed under **countably infinite** unions; that is, if A_i , for i = 1, 2, ..., are events, then so is $\bigcup_{i=1}^{\infty} A_i$.

José Figueroa-O'Farrill

mi4a (Probability) Lecture 1

21/23

23 / 23

Summary

- With any experiment or trial we associate a pair (Ω, 𝔅), where
 - Ω, the **sample space**, is the set of all possible outcomes of the experiment; and
 - F, the family of events, is a σ-field of subsets of Ω: a family of subset of Ω containing the empty set and closed under complementation and countable unions.
- In the next lecture we will see how to enhance the pair (Ω, 𝔅) to a "probability space" by assigning a measure of likelihood (i.e., a "probability") to the events in 𝔅.
- There is a dictionary between the languages of set theory and of probability. In particular, we will use the set-theoretic language freely.

σ-fields

Definition

A family \mathfrak{F} of subsets of Ω is a σ -field if

- **1** $\Omega \in \mathcal{F}$
- **2** if $A_1, A_2, \ldots \in \mathcal{F}$, then $\bigcup_{i=1}^{\infty} A_i \in \mathcal{F}$
- **③** if $A \in \mathcal{F}$, then $A^c \in \mathcal{F}$

Remark

It follows from De Morgan's theorem that for a σ -field \mathcal{F} , if $A_1, A_2, \dots \in \mathcal{F}$ then $\bigcap_{i=1}^{\infty} A_i \in \mathcal{F}$. Also $\Omega \in \mathcal{F}$, since $\Omega = \varnothing^c$. Finally, a σ -field is closed under (symmetric) difference.

Example

The smallest σ -field is $\mathcal{F} = \{\emptyset, \Omega\}$. The largest is the **power set** of Ω (i.e., the collection of all subsets of Ω).

José Figueroa-O'Farrill mi

mi4a (Probability) Lecture 1

22 / 23