

MORSE THEORY FOR MANIFOLDS WITH BOUNDARY

MACIEJ BORODZIK, ANDRÁS NÉMETHI, AND ANDREW RANICKI

ABSTRACT. We develop Morse theory for manifolds with boundary. Beside standard and expected facts like the handle cancellation theorem and the Morse lemma for manifolds with boundary, we prove that under suitable connectedness assumptions a critical point in the interior of a Morse function can be moved to the boundary, where it splits into a pair of boundary critical points. As an application, we prove that every cobordism of connected manifolds with boundary splits as a union of left product cobordisms and right product cobordisms.

1. INTRODUCTION

For some time now, Morse theory has been a very fruitful tool in the topology of manifolds. One of the milestones was the h -cobordism theorem of Smale [Sm2], and its Morse-theoretic exposition by Milnor [Mi1, Mi2]. Recently, Morse theory has become even more popular, for two reasons. In the first instance, on account of its connections with Floer homology, see e.g. [Sa, Wi, Ni, KM]. Secondly, the stratified Morse theory developed by Goresky and MacPherson [GM]. In the last 20 years Morse theory has also had an enormous impact on the singularity theory of complex algebraic and analytic varieties.

Despite much previous interest in Morse theory, there still remain uncharted territories. Morse theory for manifolds with boundary is a particular example. The theory was initiated by Kronheimer and Mrowka in [KM], and there is also a recent paper of Laudenbach [La] devoted to the subject. Our paper is a further contribution.

In this paper we prove some new results in the Morse theory for manifolds with boundary. Beside some standard and expected results, like the boundary handle cancellation theorem (Theorem 5.1) and the topological description of passing critical points on the boundary (using the notions of right and left half-handles introduced in Section 2) we discover a new phenomenon. An interior critical point can be moved to the boundary and there split into two boundary critical points. In particular, if we have a cobordism of manifolds with boundary, then under a natural topological assumption we can find a Morse function which has only boundary critical points. We use this result to prove a structure theorem for connected cobordisms of connected manifolds with connected non-empty boundary: such a cobordism splits as a union of left and right product cobordisms. This is a topological counterpart to the algebraic splitting of cobordisms obtained in [BNR]: an algebraic splitting of the chain complex cobordism of a geometric cobordism can be realized topologically by a geometric splitting.

The structure of the paper is the following. After preliminaries in Section 1.1 we study in Section 2 the changes in the topology of the level sets when crossing a boundary critical point. Theorem 2.24 is the main result: passing a boundary stable (unstable) critical point produces a left (right) half-handle attachment. In Section 3 we prove Theorem 3.1,

Date: November 28, 2012.

2010 Mathematics Subject Classification. primary: 57R19, secondary: 58E05, 58A05.

Key words and phrases. Morse theory, manifold with boundary, cobordism, bifurcation of singular points.

The first author is supported by Polish MNiSzW Grant No N N201 397937. The second author is partially supported by OTKA Grant K100796.

which moves interior critical points to the boundary. Then we pass to some more standard results, namely rearrangements of critical points in Section 4. We finish the section with our most important — up to now — application, Theorem 4.18, about the splitting of a cobordism into left product and right product cobordisms. Finally, in Section 5 we discuss the possibility of cancelling a pair of critical points.

Acknowledgements. The first author wishes to thank Rényi Institute for hospitality, the first two authors are grateful for Edinburgh Mathematical Society for a travel grant to Edinburgh and Glasgow in March 2012. The authors thank András Juhász and András Stipsicz for fruitful discussions and to Rob Kirby for drawing their attention to [KM].

1.1. Notes on gradient vector fields. To fix the notation, let us recall what a cobordism of manifolds with boundary is.

Definition 1.1. Let Σ_0 and Σ_1 be compact oriented, n -dimensional manifolds with non-empty boundary M_0 and M_1 . We shall say that (Ω, Y) is a *cobordism* between (Σ_0, M_0) and (Σ_1, M_1) , if Ω is a compact, oriented $(n+1)$ -dimensional manifold with boundary $\partial\Omega = Y \cup \Sigma_0 \cup \Sigma_1$, where Y is nonempty, $\Sigma_0 \cap \Sigma_1 = \emptyset$, and $Y \cap \Sigma_0 = M_0$, $Y \cap \Sigma_1 = M_1$.

Remark 1.2. Strictly speaking, Ω is a manifold with corners, so around a point $x \in M_0 \cup M_1$ it is locally modelled by $\mathbb{R}^{n-1} \times \mathbb{R}_{\geq 0}^2$. Accordingly, sometimes we write that Σ_0 , Σ_1 and Y , as manifolds with boundary, have tubular neighbourhoods in Ω of the form $\Sigma_0 \times [0, 1)$, $\Sigma_1 \times [0, 1)$, or $Y \times [0, 1)$, respectively. Nevertheless, in most cases it is safe (and more convenient) to assume that Ω is a manifold with boundary, i.e. that the corners are smoothed along M_0 and M_1 . Whenever possible we make this simplification in order to avoid unnecessary technicalities.

Example 1.3. Given a manifold with boundary (Σ, M) , we call $(\Sigma, M) \times [0, 1]$ a trivial cobordism, with $\Omega = \Sigma \times [0, 1]$, $Y = M \times [0, 1]$, $\Sigma_i = \Sigma \times \{i\}$, $M_i = M \times \{i\}$ for $i = 0, 1$.

We recall the notion of a Morse function. For this it is convenient to fix a Riemannian metric g on Ω .

Definition 1.4. A function $F: \Omega \rightarrow [0, 1]$ is called a *Morse function* on the cobordism (Ω, Y) if $F(\Sigma_0) = 0$, $F(\Sigma_1) = 1$, F has only Morse critical points, the critical points are not situated on $\Sigma_0 \cup \Sigma_1$, and ∇F is everywhere tangent to Y .

There are two ways of doing Morse theory on manifolds. One can either consider the gradient flow of ∇F associated with F and the Riemannian metric (in the Floer theory, one often uses $-\nabla F$), or, the so-called gradient-like vector field.

Definition 1.5. (See [Mi2, Definition 3.1].) Let F be a Morse function on a cobordism (Ω, Y) . Let ξ be a vector field on Ω . We shall say that ξ is *gradient-like with respect to F* , if the following conditions are satisfied:

- (a) $\xi \cdot F > 0$ away from the set of critical points of F ;
- (b) if p is a critical point of F of index k , then there exist local coordinates x_1, \dots, x_{n+1} in a neighbourhood of p , such that

$$F(x_1, \dots, x_{n+1}) = F(p) - (x_1^2 + \dots + x_k^2) + (x_{k+1}^2 + \dots + x_{n+1}^2)$$

and

$$\xi = (-x_1, \dots, -x_k, x_{k+1}, \dots, x_{n+1}) \text{ in } U;$$

- (b') furthermore, if p is a boundary critical point, then the above coordinate system can be chosen so that $Y = \{x_j = 0\}$ and $U = \{x_j \geq 0\}$ for some $j \in \{1, \dots, n+1\}$.
- (c) ξ is everywhere tangent to Y ;

The conditions (a) and (b) are the same as in the classical case. Condition (b') is a specification of the condition (b) in the boundary case, compare Lemma 2.6.

Smale in [Sm1] noticed that for any gradient-like vector field ξ for a function F there exists a choice of the Riemannian metric such that $\xi = \nabla F$ in that metric. The situation is identical in the boundary case. This is stated explicitly in the following lemma, whose proof is straightforward and will be omitted.

Lemma 1.6. *Let U be a paracompact k -dimensional manifold and $F: U \rightarrow \mathbb{R}$ a Morse function without critical points. Assume that ξ is a gradient-like vector field on U . Then there exists a metric g on U such that $\xi = \nabla F$ in that metric.*

Similar statement holds if U has boundary and ξ is everywhere tangent to the boundary.

Hence the two approaches – gradients and gradient-like vector fields – are equivalent. However, we shall need both approaches. In Section 3 we use gradients of functions and a specific choice of a metric, because the argument becomes slightly simpler. In Section 5 we follow [Mi2] very closely; as he uses gradient-like vector fields, we use them as well.

The next result shows that the condition from Definition 1.4 that ∇F is everywhere tangent to Y can be relaxed. We shall use this result in Proposition 4.1.

Lemma 1.7. *Let Ω be a compact, Riemannian manifold of dimension $(n+1)$ and $Y \subset \partial\Omega$ be compact as well. Let g denote the metric. Suppose that there exists a function $F: \Omega \rightarrow \mathbb{R}$, and a relative open subset $U \subset Y$ such that ∇F is tangent to Y at each point $y \in U$. Suppose furthermore, that for any $y \in Y \setminus U$ we have*

$$\ker dF \not\subset T_y Y.$$

Then, for any open neighbourhood $W \subset \Omega$ of $Y \setminus U$, there exist a metric h on Ω , agreeing with g away from W , such that $\nabla_h F$ (the gradient in the new metric) is everywhere tangent to Y .

Proof. Let us fix a point $y \in Y \setminus U$ and consider a small open neighbourhood U_y of y in W , in which we choose local coordinates x_1, \dots, x_{n+1} such that $Y \cap U_y = \{x_{n+1} = 0\}$ and $U_y \subset \{x_{n+1} \geq 0\}$. In these coordinates we have $dF = \sum_{i=1}^{n+1} f_i(x) dx_i$ for some smooth functions f_1, \dots, f_{n+1} . By the assumption, for each $x \in U_y$, there exists $i \leq n$ such that $f_i(x) \neq 0$. Shrinking U_y if needed, we may assume that for each $x \in U_y$ we have $f_i(x) \neq 0$ for some i . We may suppose that $i = 1$, hence $\pm f_1(x) > 0$. Let us choose a symmetric positive definite matrix $A_y = \{a_{ij}(x)\}_{i,j=1}^{n+1}$ so that $a_{11} = \pm f_1(x)$ and for $i > 1$, $a_{1i} = a_{i1} = f_i(x)$. A_y defines a metric h_y on U_y such that $\nabla_{h_y} F = (\pm 1, 0, \dots, 0) \subset TY$ in that metric.

Now let us choose an open subset V of $\Omega \setminus (Y \setminus U)$ such that $V \cup \bigcup_{y \in Y \setminus U} U_y$ is a covering of Ω . Let $\{\phi_V\} \cup \{\phi_y\}_{y \in Y \setminus U}$ be a partition of unity subordinate to this covering. Define

$$h = \phi_V \cdot g + \sum_{y \in Y \setminus U} \phi_y h_y.$$

Then h is a metric, which agrees with g away from W . Moreover, as for each metric h_y , and $x \in U_y \cap Y$ we have $\nabla_{h_y} F(x) \in T_x Y$ by construction, the same holds for a convex linear combination of metrics. \square

2. BOUNDARY STABLE AND UNSTABLE CRITICAL POINTS

2.1. Morse function for manifolds with boundary. The whole discussion of Morse functions on manifolds with boundary would be pointless if we did not have the following.

Lemma 2.1. *Morse functions exist. In fact, for any Morse function $f: Y \rightarrow [0, 1]$ with $f(M_0) = 0$, $f(M_1) = 1$ there exists a Morse function $F: \Omega \rightarrow [0, 1]$ whose restriction to Y is f .*




FIGURE 1. Boundary stable (on the left) and unstable critical points.

Proof. Let $f: Y \rightarrow [0, 1]$ be a Morse function on the boundary, such that $f(M_0) = 0$ and $f(M_1) = 1$. We want to extend f to a Morse function on Ω .

First, let us choose a small tubular neighbourhood U of Y and a diffeomorphism $U \cong Y \times [0, \varepsilon]$ for some $\varepsilon > 0$. Let $\tilde{F}: U \rightarrow [0, 1]$ be given by the formula

$$(2.2) \quad U \cong Y \times [0, \varepsilon] \ni (x, t) \rightarrow \tilde{F}(x, t) = f(x) - f(x)(1 - f(x))t^2.$$

The factor $f(x)(1 - f(x))$ ensures that \tilde{F} attains values in the interval $[0, 1]$ and $\tilde{F}^{-1}(i) \subset \Sigma_i$ for $i \in \{0, 1\}$. It is obvious that there exists a smooth function $F: \Omega \rightarrow [0, 1]$, which agrees on $Y \times [0, \varepsilon/2)$ with \tilde{F} , and it satisfies the Morse condition on the whole Ω . The gradient ∇F is everywhere tangent to Y . \square

Remark 2.3. The above construction yields a function with the property that all its boundary critical points are boundary stable (see Definition 2.4 below). This is due to the choice of sign -1 in front of $f(x)(1 - f(x))t^2$ in (2.2). If we change the sign to $+1$, we obtain a function with all boundary critical points boundary unstable.

We fix a Morse function $F: \Omega \rightarrow [0, 1]$ and we start to analyze its critical points. Let z be such a point. If $z \in \Omega \setminus Y$, we shall call it an *interior* critical point. If $z \in Y$, it will be called a *boundary* critical point. There are two types of boundary critical points.

Definition 2.4. Let z be a boundary critical point. We shall call it *boundary stable*, if the tangent space to the unstable manifold of z lies entirely in $T_z Y$, otherwise it is called *boundary unstable*.

The index of the boundary critical point z is defined as the dimension of the stable manifold W_z^s . If z is boundary unstable, this is the same as the index of z regarded as a critical point of the restriction f of F on Y . If z is boundary stable, we have $\text{ind}_F z = \text{ind}_f z + 1$. In particular, there are no boundary stable critical point with index 0, nor boundary unstable critical points of index $n + 1$.

Remark 2.5. We point out that we use the flow of ∇F and not of $-\nabla F$ as Kronheimer and Mrowka [KM] do, hence our definitions and formulae are slightly different from theirs.

We finish this subsection with three standard results.

Lemma 2.6 (Boundary Morse Lemma). *Assume that F has a critical point $z \in Y$ such that the Hessian $D^2 F(z)$ at z is non-degenerate, and ∇F is everywhere tangent to Y .*

Then there are local coordinates (x_1, \dots, x_{n+1}) in an open neighbourhood $U \ni z$ such that $U = \{x_1^2 + \dots + x_{n+1}^2 \leq \varepsilon^2\} \cap \{x_1 \geq 0\}$ and $U \cap Y = \{x_1 = 0\}$ for some $\varepsilon > 0$, and F in these coordinates has the form $\pm x_1^2 \pm x_2^2 \pm \dots \pm x_{n+1}^2 + F(z)$.

Proof. We choose a coordinate system y_1, \dots, y_{n+1} in a neighbourhood $U \subset \Omega$ of z such that $z = (0, \dots, 0)$, $Y = \{y_1 = 0\}$, $U = \{y_1 \geq 0\}$, and the vector field $\frac{\partial}{\partial y_1}$ is orthogonal to Y . We might assume $F(z) = 0$. The tangency of ∇F to Y implies that at each point of Y

$$(2.7) \quad \frac{\partial F}{\partial y_1}(0, y_2, \dots, y_{n+1}) = 0.$$

By Hadamard's lemma one writes

$$(2.8) \quad F = y_1 K_1(y_1, \dots, y_{n+1}) + \sum_{j=2}^{n+1} y_j K_j(y_1, y_2, \dots, y_{n+1}),$$

for some functions K_1, \dots, K_{n+1} . We can assume that for $j > 1$, K_j does not depend on y_1 . Indeed, if it does depend, we might write $K_j(y_1, \dots, y_{n+1}) = K_j(0, y_2, \dots, y_{n+1}) + y_1 L_{1j}(y_1, \dots, y_{n+1})$, and then replace K_j by $K_j(0, y_2, \dots, y_{n+1})$ and K_1 by $K_1 + \sum y_j L_{1j}$.

The condition (2.7) implies now that $K_1(0, y_2, \dots, y_{n+1}) = 0$, hence $K_1 = y_1 H_{11}(y_1, \dots, y_{n+1})$. By Hadamard's lemma applied to K_2, \dots, K_{n+1} we get

$$(2.9) \quad F = y_1^2 H_{11}(y_1, \dots, y_{n+1}) + \sum_{j,k=2}^n y_j y_k H_{jk}(y_2, \dots, y_{n+1}).$$

The non-degeneracy of $D^2 F(z)$ means that $H_{11}(z) \neq 0$. Then, after replacing $y_1 \sqrt{\pm H_{11}}$ by x_1 , we can assume that $H_{11} = \pm 1$. Finally, the sum in (2.9) can be written as $\sum_{j \geq 2} \epsilon_j x_j^2$ ($\epsilon_j = \pm 1$) by the classical Morse lemma [Mi1, Lemma 2.2]. \square

The next result is completely standard by now.

Lemma 2.10. *Assume that F is a Morse function on a cobordism (Ω, Y) between (Σ_0, M_0) and (Σ_1, M_1) . If F has no critical points then $(\Omega, Y) \cong (\Sigma_0, M_0) \times [0, 1]$. Furthermore, we can choose the diffeomorphism to map the level set $F^{-1}(t)$ to the set $\Sigma_0 \times \{t\}$.*

Proof. The proof is identical to the classical case, see e.g. [Mi2, Theorem 3.4]. \square

2.2. Half-handles. For any k we consider the k -dimensional disc $D^k = \{x_1^2 + \dots + x_k^2 \leq 1\}$. In the classical theory, an n -dimensional handle of index k is the n -dimensional manifold $H = D^k \times D^{n-k}$ with boundary

$$\partial H = (\partial D^k \times D^{n-k}) \cup (D^k \times \partial D^{n-k}) = B_0 \cup B'_0.$$

Given an n -manifold with boundary $(\Sigma, \partial\Sigma)$ and a distinguished embedding $\phi: B_0 \rightarrow \partial\Sigma$, the effect of a classical handle attachment is the n -dimensional manifold with boundary

$$(\Sigma', \partial\Sigma') = (\Sigma \cup H, (\partial\Sigma \setminus B_0) \cup B'_0),$$

where we glue along $\phi(B_0)$ identified with B_0 . The boundary $\partial\Sigma'$ is the effect of surgery on $\phi(B_0) \subset \partial\Sigma$. We now extend this construction to relative cobordisms of manifolds with boundary, using 'half-handles'. Since our ambient space Ω is $(n+1)$ -dimensional, $(n+1)$ is the dimension of the handles, and they induce n -dimensional handle attachments on Y .

In order to do this, for any $k \geq 1$ we distinguish the following subsets of D^k : the 'half-disc' $D_+^k := D^k \cap \{x_1 \geq 0\}$, and its boundary subsets $S_+^{k-1} := \partial D^k \cap \{x_1 \geq 0\}$, $S_0^{k-2} := \partial D^k \cap \{x_1 = 0\}$ and $D_0^{k-1} := D^k \cap \{x_1 = 0\}$. Clearly, S_0^{k-2} is a boundary the two $(k-1)$ -discs S_+^{k-1} and D_0^{k-1} . We will call x_1 the *cutting coordinate*.




FIGURE 2. A right half-handle of index 1. The picture on the left is the handle, the two other pictures explain the notation. The two half-circles form B , C is the bottom rectangle.

Definition 2.11. Let $0 \leq k \leq n$. An $(n+1)$ -dimensional right half-handle of index k is the $(n+1)$ -dimensional manifold $H_{right} = D^k \times D_+^{n+1-k}$, with boundary subdivided into three pieces $\partial H_{right} = B \cup C \cup N$, where

$$B := \partial D^k \times D_+^{n+1-k}, \quad C := D^k \times D_0^{n-k}, \quad N := D^k \times S_+^{n-k}.$$

One has the following intersections too

$$B_0 := C \cap B = \partial D^k \times D_0^{n-k}, \quad N_0 := C \cap N = D^k \times S_0^{n-k-1}.$$

Hence the handle H is cut along C into two pieces, one of them is the half-handle H_{right} . Note that (C, B_0) is a n -dimensional handle of index k .

Symmetrically, we define the left half-handles by cutting the handle H along the left-component disc D^k .

Definition 2.12. Fix k with $1 \leq k \leq n+1$. An $(n+1)$ -dimensional left half-handle of index k is the $(n+1)$ -dimensional disk $H_{left} := D_+^k \times D^{n+1-k}$ with boundary subdivided into three pieces $\partial H_{left} = B \cup C \cup N$, where

$$B := S_+^{k-1} \times D^{n+1-k}, \quad C := D_0^{k-1} \times D^{n+1-k}, \quad N := D_+^k \times \partial D^{n+1-k}.$$

Furthermore, we specify $B_0 := C \cap B = S_0^{k-2} \times D^{n+1-k}$ and $N_0 := N \cap C = D_0^{k-1} \times \partial D^{n+1-k}$.

A *half-handle* will from now on refer to either a right half-handle or left half-handle. We pass to half-handle attachments. We will attach a half-handle along B . The definitions of the right half-handle attachment and the left half-handle attachment are formally very similar, but there are significant differences in the properties of the two operations.

Definition 2.13. Let $(\Omega, Y; \Sigma_0, M_0, \Sigma_1, M_1)$ be an $(n+1)$ -dimensional relative cobordism. Given an embedding $\Phi: (B, B_0) \hookrightarrow (\Sigma_1, M_1)$ define the relative cobordism $(\Omega', Y'; \Sigma_0, M_0, \Sigma'_1, M'_1)$ obtained from $(\Omega, Y; \Sigma_0, M_0, \Sigma_1, M_1)$ by *attaching a (right or left) half-handle of index k* by

$$\begin{aligned} \Omega' &= \Omega \cup_B H, & Y' &= Y \cup_{B_0} C, \\ \Sigma'_1 &= (\Sigma_1 \setminus B) \cup N, & M'_1 &= (M_1 \setminus B_0) \cup N_0. \end{aligned}$$

See Figure 4 and Figure 5 for right, respectively left half-handle attachments.

We point out that in the case of the right half-handle attachment, any embedding of B_0 into M_1 determines (up to an isotopy) an embedding of pairs $(B, B_0) \hookrightarrow (\Sigma_1, M_1)$. Indeed, as $(B, B_0) = (\partial D^k \times (D_+^{n+1-k}, D_0^{n-k}))$, a map $\phi: B_0 \hookrightarrow M_1$ extends to a map $\Phi: B \hookrightarrow \Sigma_1$ in a collar neighbourhood of M_1 in Σ_1 . (This is not the case in the left half-handle attachment.)

In particular, *in the case of right attachments, we specify only the embedding $B_0 \hookrightarrow M_1$.*




FIGURE 3. A left half-handle of dimension 3 and index $k = 2$. The two lines are B_0 , the bottom rectangle is C . B is the surface between the two half circles on the picture.

Example 2.14. (a) We exemplify the left half-handle attachment for $k = 1$. In this case B_0 is empty. If we are given an embedding of $B \cong \{1\} \times D^n$ into $\Sigma_1 \setminus M_1$, we glue $[0, 1] \times D^n$ to Ω along B . Then we set $Y' = Y \sqcup \{0\} \times D^n$, $\Sigma'_1 = (\Sigma_1 \setminus B) \cup [0, 1] \times \partial B$ and $M' = M \sqcup \{0\} \times \partial B$.

(b) The right half-handle attachment of index 0 is the disconnected sum $\Omega \sqcup D_+^{n+1}$ with boundary $\partial D_+^{n+1} = S_+^n \cup D_0^n$. We think of first disc S_+^n as a part of Σ'_1 , while the second disc as a part of Y' , and $M'_1 = M_1 \cup S_0^{n-1}$.

Remark 2.15. In the next subsection we shall see that crossing a boundary stable critical points corresponds to left half-handle attachment, while a boundary unstable critical point corresponds to a right half-handle attachment. Theorem 3.1 can be interpreted informally, as splitting a handle into a right half-handle and left half-handle. This also motivates the name ‘half-handle’.

2.3. Elementary properties of half-handle attachments. The following results are trivial consequences of the definitions.

Lemma 2.16. *Let Ω' be the result of a right half-handle attachment to Ω along $(B, B_0) \hookrightarrow (\Sigma_1, M_1)$. Let B' be B pushed slightly off M_1 into the interior of Σ_1 . Let $\tilde{\Omega}$ be a result of attaching a (standard) handle of index k to Ω along B' . Then Ω' and $\tilde{\Omega}$ are diffeomorphic.*

Proof. This is obvious, since when we forget about C and B_0 , the pair (H_{right}, B) is a standard $(n + 1)$ -dimensional handle of index k . \square

In particular, the effect of a right half-handle attachment on Ω is the same as the effect of a standard handle attachment of the same index.

The situation is completely different in the case of left half-handle attachments.

Lemma 2.17. *If Ω' is a result of a left half-handle attachment, then Ω' is diffeomorphic to Ω .*

Proof. By definition the pair (H_{left}, B) is diffeomorphic to the pair $(D^n \times [0, 1], D^n \times \{0\})$. Attaching H_{left} along B to Ω does not change the diffeomorphism type of Ω . \square

The effect on Y of a right/left half-handle attachments are ‘almost’ the same.

Lemma 2.18. *If $(\Omega', Y'; \Sigma_0, M_0, \Sigma'_1, M'_1)$ is the result of left (respectively, right) half-handle attachment to $(\Omega, Y; \Sigma_0, M_0, \Sigma_1, M_1)$ along $(B, B_0) \hookrightarrow (\Sigma_1, M_1)$, then Y' is a result of a classical handle attachment of index $k - 1$ (respectively k) along B_0 .*

Proof. This follows immediately from Definition 2.13. \square

The effects of half handle attachment on Σ are also easily described. The next lemma is a direct consequence of the definitions; its proof is omitted. We refer to Figures 4 and 5.




FIGURE 4. Right half-handle attachment. Here $k = 1$, $n = 2$. On the right, the two black points represent a sphere S^0 with a neighbourhood B_0 in M_1 and B in Σ_1 . On the picture on the right the dark green coloured part of the handle belongs to Σ_1 , the dashed lines belong to Σ_1 and are drawn only to make the picture look more 'three-dimensional'.

Lemma 2.19. (a) If $(\Omega', Y'; \Sigma_0, M_0, \Sigma'_1, M'_1)$ is the result of left half-handle attachment to $(\Omega, Y; \Sigma_0, M_0, \Sigma_1, M_1)$ along $(B, B_0) \hookrightarrow (\Sigma_1, M_1)$, then

$$\Sigma'_1 \cong \Sigma_1 \setminus B.$$

(b) If $(\Omega', Y'; \Sigma_0, M_0, \Sigma'_1, M'_1)$ is the result of index k right half-handle attachment to $(\Omega, Y; \Sigma_0, M_0, \Sigma_1, M_1)$ along $B_0 \hookrightarrow M_1$, then

$$\Sigma'_1 \cong \Sigma_1 \cup_{B_0} N,$$

where here N is an n -dimensional disk $D^k \times D^{n-k}$ and $B_0 = S^{k-1} \times D^{n-k}$.

These facts also emphasize that right half-handle attachments and left half-handle attachments are somehow dual operations on Σ . This can be seen also at the Morse function level: changing a Morse function F to $-F$ changes all right half-handles to left-half handles and conversely, see Section 2.4 and 2.5 below. But the above lemma shows another aspect as well: a right half handle attachment consists on gluing a disk, a left half-handle attachment consists of removing a disk. Indeed, in the case of right attachment, $(\Sigma'_1, M'_1) = (\Sigma_1 \cup D^k \times D^{n-k}, \partial \Sigma_1)$ associated with an embedding $\Phi: \partial D^k \times D^{n-k} \rightarrow M_1$. On the other hand, by definition, for an embedding $\Phi': (D^{k-1} \times D^{n+1-k}, \partial D^{k-1} \times D^{n+1-k}) \rightarrow (\Sigma_1, M_1)$ the pair

$$(\Sigma'_1, M'_1) = (\text{closure of } (\Sigma_1 \setminus D^{k-1} \times D^{n+1-k}), \partial \Sigma'_1)$$

is obtained from (Σ_1, M_1) by a *handle detachment* of index $k - 1$. Hence we obtain:

Corollary 2.20. The effect on (Σ_1, M_1) of a right half-handle attachment of index k (at the level of cobordism) is a handle attachment of index k at the level of (Σ_1, M_1) . Conversely, the effect on (Σ_1, M_1) of a left half-handle attachment of index k (at the level of cobordism) is a handle detachment of index $k - 1$ at the level of (Σ_1, M_1) . In particular, M'_1 is obtained from M_1 as the result of a k surgery in the first case, and $(k - 1)$ surgery in the second.

The duality can be seen also as follows: we can cancel any handle attachment by a suitably defined handle detachment, and conversely.

The following definition introduces a terminology which is rather self-explanatory. We include it for completeness of the exposition.

Definition 2.21. We shall say that a cobordism (Ω', Y') between (Σ, M) and (Σ', M') is a *right (respectively left) half-handle attachment* of index k , if $(\Omega', Y', \Sigma', M')$ is a result of right (respectively left) half-handle attachments of index k (in the sense of Definition 2.13) to $(\Sigma \times [0, 1], M \times [0, 1], \Sigma \times \{0\}, M \times \{0\}, \Sigma \times \{1\}, M \times \{1\})$.




FIGURE 5. Left handle attachment with $k = 2$ and $n = 2$. This time the sphere on the left (denoted by two points) bounds a disk in Σ_1 .

We conclude this section by studying homological properties of the handle attachment. These properties will be used in [BNR]. The proofs are standard and are left to the reader.

Let (H_+, C, B, N) be a half-handle of index k .

Lemma 2.22. *If $(H_{\text{right}}, C, B, N)$ is a right half-handle, then the pair (C, B_0) is a strong deformation retract of (H_+, B) , while $(D^k, \partial D^k)$ is a strong deformation retract of (C, B_0) . In particular, $H_j(H_+, B) \cong H_j(C, B_0) = \mathbb{Z}$ for $j = k$, and it is zero otherwise.*

The situation is completely different for left half-handles.

Lemma 2.23. *If $(H_{\text{left}}, C, B, N)$ is a left half-handle, then the pair (H_{left}, B) retracts onto the trivial pair (point, point). In particular, all the relative homologies $H_*(H_{\text{left}}, B)$ vanish. On the other hand, (D_0^{k-1}, S_0^{k-2}) is a strong deformation retract of (C, B_0) , hence $H_j(C, B_0) = \mathbb{Z}$ for $j = k - 1$, and it is zero otherwise. Therefore, the inclusion $(C, B_0) \hookrightarrow (H_{\text{left}}, B)$ induces a surjection on homologies.*

2.4. Boundary critical points and half-handles. Consider a Morse function F on a cobordism (Ω, Y) and assume that it has a single boundary critical point z of index k with critical value c .

Theorem 2.24. *If z is boundary stable (unstable), then the cobordism is a left (right) half-handle attachment of index k respectively.*

Proof. We can assume that $c = F(z) = 0$. Let ρ be a positive number small enough and let U be a ‘half’ ball around z of radius 2ρ such that in U we can choose Morse coordinates x_1, \dots, x_{n+1} (cf. Lemma 2.6) with

$$U = \{x_1^2 + \dots + x_{n+1}^2 \leq 4\rho^2\} \cap \{x_1 \geq 0\},$$

and $Y \cap U$ defined by $\{x_1 = 0\}$, and

$$F(x_1, \dots, x_{n+1}) = -a^2 + b^2,$$

where if z is boundary stable we set

$$(2.25) \quad a^2 = x_1^2 + x_2^2 + \dots + x_k^2, \quad b^2 = x_{k+1}^2 + \dots + x_{n+1}^2 \quad (k \geq 1),$$

and if z is boundary unstable

$$(2.26) \quad a^2 = x_2^2 + \dots + x_{k+1}^2, \quad b^2 = x_1^2 + x_{k+2}^2 + \dots + x_{n+1}^2 \quad (k \geq 0).$$

We also assume that x_1, \dots, x_{n+1} is an Euclidean orthonormal coordinate system.

Next, we consider $\varepsilon > 0$ such that $\varepsilon \ll \rho$, and we define the space \tilde{H} bounded by the following conditions (see Figure 6)

$$\tilde{H} := \{-a^2 + b^2 \in [-\varepsilon^2, \varepsilon^2], \quad a^2 b^2 \leq (\rho^4 - \varepsilon^4)/4, \quad x_1 \geq 0\}.$$




FIGURE 6. A schematic presentation of \tilde{H} , \tilde{B} , \tilde{P} , \tilde{K} from the proof of Theorem 2.24. To each point (a^2, b^2) in \tilde{H} on the picture, correspond all those points (x_1, \dots, x_{n+1}) for which (2.25) or (2.26) holds and $x_1 \geq 0$.

Observe that

$$\tilde{H} \subset U.$$

Let us now define the following parts of the boundary of \tilde{H}

$$(2.27) \quad \begin{aligned} \tilde{B} &= \partial\tilde{H} \cap \{-a^2 + b^2 = -\varepsilon^2\} \subset F^{-1}(-\varepsilon^2), \\ \tilde{P} &= \partial\tilde{H} \cap \{-a^2 + b^2 = \varepsilon^2\} \subset F^{-1}(\varepsilon^2), \\ \tilde{K} &= \partial\tilde{H} \cap \{a^2 b^2 = (\rho^4 - \varepsilon^4)/4\}, \\ \tilde{C} &= \partial\tilde{H} \cap \{x_1 = 0\} \subset Y. \end{aligned}$$

We have $\tilde{B} \cup \tilde{P} \cup \tilde{K} \cup \tilde{C} = \partial\tilde{H}$ (on Figure 6 we do not see \tilde{C} , because this would require one more dimension). If z is boundary unstable and $k = 0$ in (2.26) then the term a^2 is missing and $\tilde{B} = \emptyset$. Otherwise $\tilde{B} \neq \emptyset$.

Lemma 2.28. *The flow of ∇F is tangent to \tilde{K} .*

Proof. Assume the critical point is boundary stable. Then the time t flow acts by

$$(x_1, \dots, x_{n+1}) \rightarrow (e^{-2t}x_1, \dots, e^{-2t}x_k, e^{2t}x_{k+1}, \dots, e^{2t}x_{n+1}),$$

hence $a^2 \rightarrow e^{-4t}a^2$ and $b^2 \rightarrow e^{4t}b^2$, and the hypersurface $a^2 b^2 = \text{const}$ is preserved. \square

Lemma 2.29. *The inclusion of pair of spaces*

$$\left(F^{-1}(-\varepsilon^2) \cup_{\tilde{B}} \tilde{H}, Y \cap \left(F^{-1}(-\varepsilon^2) \cup_{\tilde{B}} \tilde{H} \right) \right) \subset (\Omega, Y)$$

admits a strong deformation retract (inducing an isomorphism of these pairs).

Proof. By Lemma 2.10 we can assume that (Ω, Y) is $(F^{-1}([-\varepsilon^2, \varepsilon^2]), Y \cap F^{-1}([-\varepsilon^2, \varepsilon^2]))$.

First we assume that \tilde{B} is not empty, and it is given by the equation (2.27) in U . Set $\Sigma_- = \text{closure of } (F^{-1}(-\varepsilon^2) \setminus \tilde{B})$ and let T_- be the part of the boundary of Σ_- given by

$$T_- = \text{closure of } (\partial\Sigma_- \setminus \partial F^{-1}(-\varepsilon^2)).$$

Obviously $T_- \subset \tilde{B}$, see Figure 7. Let us choose a collar of T_- in Σ_- , that is a subspace $U_- \subset \Sigma_-$ diffeomorphic to $T_- \times [0, 1]$, T_- identified with $T_- \times \{0\}$ and $\partial T_- \times [0, 1] \subset \partial\Sigma_- \cap \partial F^{-1}(-\varepsilon^2)$. Let T'_- be the space identified with $T_- \times \{1\}$ by this diffeomorphism.

Similarly, let $\Sigma_+ = \text{closure of } (F^{-1}(\varepsilon^2) \setminus \tilde{P})$, and $T_+ = \text{closure of } (\partial\Sigma_+ \setminus \partial F^{-1}(\varepsilon^2))$. We also define Ω_0 as the closure of $\Omega \setminus \tilde{H}$. Clearly F has no critical points in Ω_0 and ∇F is




FIGURE 7. Notation used in Lemma 2.29. Please note that the left picture is drawn on Σ_- , while the right one is on Ω .




FIGURE 8. Contractions Π_H and Π_V from the proof of Lemma 2.29. The set V is now drawn as a rectangle.

everywhere tangent to $\partial\Omega_0 \setminus (\Sigma_- \cup \Sigma_+) = (Y \cap \Omega_0) \cup \tilde{K}$ by Lemma 2.28. In particular, by Lemma 2.10, the flow of ∇F on Ω_0 yields a diffeomorphism between Σ_- and Σ_+ , mapping T_- to T_+ . We define $V \subset \Omega$ as the closure of the set of points v such that a trajectory going through v hits U_- . Lemma 2.10 implies that there is a diffeomorphism $V \cong T_- \times [0, 1] \times [-\varepsilon^2, \varepsilon^2]$ such that for $(x, t, s) \in V$ we have $F(x, t, s) = s$. Finally, we also define $V^* := \{(x, t, s) \in V : s \leq \varepsilon^2(1 - 2t)\}$.

We define the contraction in two steps: vertical and horizontal. The vertical contraction is defined as follows. For $v \in \tilde{H} \cup V^*$ we define $\Pi_V(v) = v$. For a point $v \in \Omega_0 \setminus V$ we take for $\Pi_V(v)$ the unique point $s \in \Sigma_-$ such that a trajectory of ∇F goes from s to v . Finally if $v = (x, t, s) \in V \setminus V^*$ we define $\Pi_V(v) = (x, t, \varepsilon^2(1 - 2t))$.

By construction, the image of Π_V is $\tilde{H} \cup V^* \cup F^{-1}(-\varepsilon^2)$. Next, we define Π_H .

It is an identity on $\tilde{H} \cup F^{-1}(-\varepsilon^2)$, and maps $(x, t, s) \in V^*$ to $(x, t - (\varepsilon^2 + s)/(2\varepsilon^2), -\varepsilon^2)$ if $s \leq \varepsilon^2(2t - 1)$, and to $(x, 0, s - 2\varepsilon^2 t)$ otherwise. Note that the expressions agree for any (x, t, s) with $s = \varepsilon^2(2t - 1)$ and these points are sent to $(x, 0, -\varepsilon^2)$. Both Π_H and Π_V are continuous retractions, by smoothing corners we can modify them into smooth retractions; also they can be extended in a natural way to strong deformation retracts. By construction, the retracts preserve Y too. See also Figure 8.

If \tilde{B} is empty, then \tilde{H} is necessarily a unstable (right) half-handle of index 0, $F^{-1}([-\varepsilon^2, \varepsilon^2])$ is a disconnected sum of \tilde{H} and the manifold $F^{-1}(-\varepsilon^2) \times [-\varepsilon^2, \varepsilon^2]$. \square

Continuation of the proof of Theorem 2.24. We want to show that \tilde{H} is a half-handle.

By subsection 2.2 we have the following description in local coordinates of the left half-handle (2.30) and right half-handle (2.31) with cutting coordinate x_1 :

$$(2.30) \quad H_{left} = \{x_1^2 + \dots + x_k^2 \leq 1\} \cap \{x_{k+1}^2 + \dots + x_{n+1}^2 \leq 1\} \cap \{x_1 \geq 0\}$$

$$(2.31) \quad H_{right} = \{x_2^2 + \cdots + x_{k+1}^2 \leq 1\} \cap \{x_1^2 + x_{k+2}^2 + \cdots + x_{n+1}^2 \leq 1\} \cap \{x_1 \geq 0\}.$$

We consider the subsets R and S of \mathbb{R}^2 and a diffeomorphism $\psi: R \rightarrow S$ continuous on the boundary, where

$$R = \{(u, v) \in \mathbb{R}^2: u \geq 0, v \geq 0, uv \leq (\rho^4 - \varepsilon^4)/4, -u + v \in [-\varepsilon^2, \varepsilon^2]\},$$

$$S = \{(u, v) \in \mathbb{R}^2: u \in [0, \varepsilon], v \in [0, \varepsilon]\}.$$

Assume that ψ maps the edge of R given by $\{-u + v = -\varepsilon^2\}$ to the edge $\{u = \varepsilon\}$ of S and the images of coordinate axes are the corresponding coordinate axes. (Note that R can be seen on Figure 6 if we replace a^2 by u and b^2 by v .)

We lift ψ to a diffeomorphism Ψ between \tilde{H} and H_{right} (respectively H_{left}) as follows. First let us write $\psi(u, v) = (\psi_1(u, v), \psi_2(u, v))$. As ψ maps axes to axes, we have $\psi_1(0, v) = 0$ and $\psi_2(u, 0) = 0$. Furthermore $\psi_1, \psi_2 \geq 0$. By Hadamard's lemma there exists smooth functions ξ and η such that

$$\psi(u, v) = (u\xi(u, v)^2, v\eta(u, v)^2).$$

We define now

$$\Psi(x_1, \dots, x_{n+1}) = (\xi(a, b)x_1, \dots, \xi(a, b)x_k, \eta(a, b)x_{k+1}, \dots, \eta(a, b)x_{n+1})$$

if z is boundary stable, and

$$\Psi(x_1, \dots, x_{n+1}) = (\eta(a, b)x_1, \xi(a, b)x_2, \dots, \xi(a, b)x_k, \eta(a, b)x_{k+1}, \dots, \eta(a, b)x_{n+1})$$

if z is boundary unstable. Here a and b are given by (2.25) or (2.26). By the very construction, Ψ maps $(\tilde{H}, \tilde{B}, \tilde{C})$ diffeomorphically to the triple (H, B, C) , where

$$H = \{a^2 \in [0, \varepsilon^2], b^2 \in [0, \varepsilon^2], x_1 \geq 0\}$$

$$B = \{a^2 = -\varepsilon^2, b^2 \in [0, \varepsilon^2], x_1 \geq 0\}$$

$$C = \{a^2 \in [0, \varepsilon^2], b^2 \in [0, \varepsilon^2], x_1 = 0\}.$$

After substituting for a and b the values from (2.25) or (2.26) (depending on whether z is boundary stable or unstable), we recover the model (2.31) of a right half-handle if z is boundary unstable; or the model (2.30) of a left half-handle (both of index k). \square

The fact that each half-handle can be presented in a left or right model will be now used to show the following converse to Theorem 2.24.

Proposition 2.32. *Let $(\Omega, Y) = (\Sigma_0 \times [0, 1], M_0 \times [0, 1])$ be a product cobordism between (Σ_0, M_0) and $(\Sigma_1, M_1) \cong (\Sigma_0, M_0)$. Let us be given a half-handle (H, C, B) of index k and an embedding of $B_0 = C \cap B$ into M_1 (respectively an embedding of (B, B_0) into (Σ_1, M_1)), and let (Ω', Y') be the result of a right half-handle attachment along B_0 (respectively, a left half-handle attachment along (B, B_0)) of index k . Then, there exists a Morse function $F: (\Omega', Y') \rightarrow \mathbb{R}$, which has a single boundary unstable critical point (respectively, a single boundary stable critical point) of index k on H and no other critical points. In particular, F is a Morse function on a cobordism (Ω', Y') .*

Proof. We shall prove the result for right half-handle attachment, the other case is completely analogous. In this case we have B_0 embedded into M_1 and we extend this embedding to an embedding of B into Σ_1 (see Definition 2.13). We shall be using the notation from the proof of Theorem 2.24, with $\varepsilon^2 = 1$ and $\rho^2 = 4$. We have $\Omega' = \Omega \cup_B H$, where we identify (H, B, C) with $(\tilde{H}, \tilde{B}, \tilde{C})$ using the diffeomorphism Ψ . Let now $\Sigma_- = \text{closure of } (\Sigma_1 \setminus B)$ and $T_- = \text{closure of } (\partial\Sigma_- \setminus \partial\Sigma_1)$.

By Lemma 2.29 used ‘back to front’ we deduce that Ω'' is diffeomorphic to Ω' , where

$$(2.33) \quad \Omega'' = \Omega' \cup \Sigma_- \times [-1, 1].$$

The gluing in (2.33) is as follows: we glue $\Sigma_- \times \{-1\}$ to Σ_1 and $T_- \times [-1, 1]$ to \tilde{K} (in the proof of Lemma 2.29 we proved that \tilde{K} is $T_- \times [-1, 1]$). Now let us define a function

$$F(x) = \begin{cases} t & \text{if } x = (v, t) \in \Sigma_0 \times \{t\} \subset \Sigma_0 \times [0, 1] = \Omega \\ 2 + t & \text{if } x = (v, t) \in \Sigma_- \times \{t\} \subset \Sigma_- \times [-1, 1] \\ 2 - \sum_{j=1}^k x_j^2 + \sum_{j=k+1}^{n+1} x_j^2 & \text{if } x = (x_1, \dots, x_{n+1}) \in H. \end{cases}$$

Checking continuity of F is straightforward. Moreover F is piecewise smooth and smooth away from B . Let us choose a vector field \vec{e} on B , normal to B and pointing towards H . Then $\langle \nabla F, \vec{e} \rangle$ is positive on both sides of B . By a standard analytic argument F can be approximated by a smooth function equal to F away from some neighbourhood of B , which has no new critical point. \square

2.5. Left and right product cobordisms and traces of handle attachments. In this subsection we relate half-handle attachments to handle attachments and detachments in the sense subsection 2.3. This also creates a dictionary between surgery theoretical notions (traces of handle attachments and detachments) and Morse theoretical (additions of half-handles). To begin with, let (Ω, Y) be a cobordism between (Σ_0, M_0) and (Σ_1, M_1) .

Definition 2.34. We shall say that Ω is a *left product cobordism* if $\Omega \cong \Sigma_0 \times [0, 1]$. Similarly, if $\Omega \cong \Sigma_1 \times [0, 1]$, then we shall say that Ω is a *right product cobordism*.

Proposition 2.35. (a) *If (Ω, Y) is a cobordism between (Σ_0, M_0) and (Σ_1, M_1) consisting only of left half-handle attachments, then it is a left-product cobordism. Conversely, if it consists only on right half-handle attachments, then it is a right product cobordism.*

(b) *Let $F: \Omega \rightarrow [0, 1]$ be a Morse function in the sense of Definition 1.4. Assume that F has no critical points in the interior of Ω . If all critical points on the boundary are boundary stable, then F is a left-product cobordism. If all critical points are boundary unstable, then F is a right product cobordism.*

Proof. The two statements (a) and (b) are equivalent via Theorem 2.24 and Proposition 2.32. The stable-unstable (right-left) statements are also equivalent by replacing the Morse function F by $-F$. The stable case follows from Lemma 2.17. \square

The next results of this subsection will be not used in this paper, but we insert them because they bridge surgery techniques and applications, e.g. with [Ra] or [BNR].

In order to clarify what we wish, let us recall that by Theorem 2.24 if a Morse function F defined on a cobordism (Ω, Y) has only one critical point of boundary type then (Ω, Y) is a half-handle attachment. Proposition 2.32 is the converse of this, the (total) space of a half-handle attachment can be thought as a cobordism with a Morse function on it with only one critical point.

We wish to establish the analogues of these statements ‘at the level of Σ ’. In Subsection 2.3 we proved that the output of a right/left half-handle attachment at the level of Σ induces a handle attachment/detachment. The next lemma is the converse of this statement. (In fact, the output cobordism provided by it can be identified with the cobordism constructed in Proposition 2.32.)

Lemma 2.36. *Assume that (Σ_1, M_1) is a result of a handle attachment (respectively detachment) to (Σ_0, M_0) . Then, there exists a cobordism $(\Omega, Y; \Sigma_0, M_0, \Sigma_1, M_1)$ such that $\Omega \cong \Sigma_1 \times [0, 1]$ (respectively $\Omega \cong \Sigma_0 \times [0, 1]$).*




FIGURE 9. Lemma 2.36. On the left a 1-handle is attached to Σ_0 . On the right there is a cobordism between Σ_0 and Σ_1 , which is a right product cobordism.

Proof. Assume that (Σ_1, M_1) arises from a handle attachment to (Σ_0, M_0) , i.e. $\Sigma_1 = \Sigma_0 \cup D^k \times D^{n-k}$. Let us define $\Omega = \Sigma_1 \times [0, 1]$. The boundary $\partial\Omega$ can be split as

$$\begin{aligned} \partial\Omega &= \left(\Sigma_0 \cup D^k \times D^{n-k} \right) \times \{0\} \cup (M_1 \times [0, 1]) \cup (\Sigma_1 \times \{1\}) \\ &= \Sigma_0 \times \{0\} \cup Y \cup \Sigma_1 \times \{1\}, \end{aligned}$$

where $Y = D^k \times D^{n-k} \cup (M_1 \times [0, 1])$. Its $D^k \times D^{n-k}$ part can be ‘pushed inside’ Ω transforming (diffeomorphically) Ω into a cobordism, see Figure 9.

Analogous construction can be used in the case of a handle detachment. If (Σ'_1, M'_1) is a result of a handle detachment from (Σ_0, M_0) , then the trace of the handle detachment is the cobordism between (Σ_0, M_0) and (Σ'_1, M'_1) such that

$$(\Omega', Y') = (\Sigma_0 \times [0, 1], M_0 \times [0, 1] \cup D^k \times D^{n-k}).$$

□

Definition 2.37. The cobordism $(\Omega, Y; \Sigma_0, M_0, \Sigma_1, M_1)$ determined by the Lemma 2.36 is called the *trace* of a handle attachment of (Σ_0, M_0) (respectively the *trace* of a handle detachment).

3. SPLITTING INTERIOR HANDLES

We prove here the theorem about moving critical points to the boundary.

Theorem 3.1. *Assume that on a cobordism (Ω, Y) between (Σ_0, M_0) and (Σ_1, M_1) we have a Morse function F with a single critical point z of index $k \in \{1, \dots, n\}$ in the interior of Ω situated on the level set $\Sigma_{1/2} = F^{-1}(F(z))$. If*

(3.2) the connected component of $\Sigma_{1/2}$ containing z has non-empty intersection with Y ,

then there exists a function $G: \Omega \rightarrow [0, 1]$, such that

- G agrees with F in a neighbourhood of $\Sigma_0 \cup \Sigma_1$;
- ∇G is everywhere tangent to Y ;
- G has exactly two critical points z^s and z^u , which are both on the boundary and of index k . The point z^s is boundary stable and z^u is boundary unstable.
- There exists a Riemannian metric such that there is a single trajectory of ∇G from z^s to z^u inside Y .

Remark 3.3. A careful reading of the proof shows that we can in fact construct a smooth homotopy G_t such that $F = G_0$, $G = G_1$ and there exists $t_0 \in (0, 1)$ such that G_t has




FIGURE 10. The trajectories of the gradient vector field of D for values of $a > 0$, $a = 0$ and $a < 0$.

a single interior critical point for $t < t_0$, two boundary critical points for $t > t_0$ and a degenerate critical point on the boundary for $t = t_0$. See Remark 3.15.

The proof of Theorem 3.1 occupies Sections 3.2 to 3.4. We make a detailed discussion of Condition (3.2) in Section 3.5.

3.1. About the proof. The argument is based on the following two-dimensional picture. Consider the set $Z = \{(x, y) \in \mathbb{R}^2 : x \geq 0\}$ and the function $D : Z \rightarrow \mathbb{R}$ given by

$$D(x, y) = y^3 - yx^2 + ay,$$

where $a \in \mathbb{R}$ is a parameter. Observe that the boundary of Z given by $\{x = 0\}$ is invariant under the gradient flow of D (see Figure 10). The proof of the following lemma is completely straightforward and will be omitted.

Lemma 3.4. *For $a > 0$, D has a single Morse critical point in the interior of Z . For $a < 0$, D has two critical points on the boundary of Z .*


For $a = 0$, D acquires a D_4^- singularity at the origin (see e.g. [AGV]).

The proof of Theorem 3.1 starts with introducing ‘local/global’ coordinates $(x, y, u_1, \dots, u_{n-1})$ at z , in which F has the form $D(x, y) \pm u_1^2 \pm \dots \pm u_{n-1}^2$, hence it also parametrizes a neighbourhood of a path connecting z with a point of Y . Then we change the parameter a (which we originally assume to be equal to 1) to $-\delta$, where δ is very small positive number (which corresponds to the move of the critical point to the boundary along the chosen path).

3.2. Proof of Theorem 3.1 under additional assumption. We first give the proof assuming the existence of such coordinate system as in 3.1, described explicitly in the next proposition (which is proved in Section 3.4).

Proposition 3.5. *There exists $\eta > 0$, $\eta \ll 1$ and an open ‘half-disc’ $U \subset \Omega$, intersecting Y along a disk, and coordinates $x, y, u_1, \dots, u_{n-1}$ such that in these coordinates U is given by*

$$0 \leq x < 3 + \eta, \quad |y| < \eta, \quad \sum_{j=1}^{n-1} u_j^2 < \eta^2,$$

FIGURE 11. Sets U_1, U_{21}, S_1 and S_2 in two dimensions (coordinates x and y).

$U \cap Y$ is given by $\{x = 0\}$, and in these coordinates F is given by

$$y^3 - yx^2 + y + \frac{1}{2} + \sum_{j=1}^{n-1} \epsilon_j u_j^2,$$

where $\epsilon_1, \dots, \epsilon_{n-1} \in \{\pm 1\}$ are choices of signs. In particular $\#\{j: \epsilon_j = -1\} = k - 1$, where $k = \text{ind}_z F$.

Assuming the proposition, we prove Theorem 3.1. Let us introduce some abbreviations.

$$(3.6) \quad \vec{u} = (u_1, \dots, u_{n-1}), \quad \vec{u}^2 = \sum_{j=1}^{n-1} \epsilon_j u_j^2, \quad \|\vec{u}\|^2 = \sum_{j=1}^{n-1} u_j^2.$$

We fix a small real number $\varepsilon > 0$ such that $\varepsilon \ll \eta$ and two subsets $U_1 \subset U_2$ of U by

$$\begin{aligned} U_1 &= \{|y| \leq \varepsilon, x \leq 3\} \cup \{(x-3)^2 + y^2 \leq \varepsilon^2\} \\ U_2 &= \{|y| \leq 2\varepsilon, x \leq 3\} \cup \{(x-3)^2 + y^2 \leq 4\varepsilon^2\}. \end{aligned}$$

The difference $U_{21} := \overline{U_2} \setminus U_1$ splits into two subsets $S_1 \cup S_2$ (see Figure 11), where

$$S_1 = U_{21} \cap \{x \leq 3\}, \quad S_2 = U_{21} \cap \{x \geq 3\}.$$

For a point $v = (x, y, u_1, \dots, u_{n-1}) \in U$, let us define:

$$\tilde{s}(v) = \begin{cases} 1 & \text{if } v \in U_1, \\ 0 & \text{if } v \in \overline{U \setminus U_2}, \\ 2 - \frac{|y|}{\varepsilon} & \text{if } v \in S_1, \\ 2 - \frac{\sqrt{(x-3)^2 + y^2}}{\varepsilon} & \text{if } v \in S_2. \end{cases}$$

The above formula defines a continuous function $\tilde{s}: U_2 \rightarrow [0, 1]$. It is smooth away of $\partial S_1 \cup \partial S_2$. We can perturb it to a C^∞ function s , with the following properties:

- (S1) $s^{-1}(1) = U_1$, $s^{-1}(0) = \{|y| \geq 2\varepsilon - \varepsilon^2\} \cup \{(x-3)^2 + y^2 \geq 4\varepsilon^2 - \varepsilon^3\}$;
- (S2) $\frac{\partial s}{\partial u_j} = 0$ for any $j = 1, \dots, n-1$;
- (S3) $\frac{\partial s}{\partial x} = 0$, and $\left| \frac{\partial s}{\partial y} \right| < \frac{2}{\varepsilon}$ at all points of S_1 . Furthermore $y \frac{\partial s}{\partial y} < 0$ at all points of S_1 ;
- (S4) if $v \in S_2$ and we choose radial coordinates $x = 3 + r \cos \phi$, $y = r \sin \phi$ (where $r \in [\varepsilon, 2\varepsilon]$ and $\phi \in [-\frac{\pi}{2}, \frac{\pi}{2}]$), then $\left| \frac{\partial s}{\partial r} \right| < \frac{2}{\varepsilon}$ and $\left| \frac{\partial s}{\partial \phi} \right| < \varepsilon$

Observe that \tilde{s} satisfies (S1)–(S4) at every point, where it is smooth, the only issue is that on $S_1 \cap S_2$, \tilde{s} fails to be C^2 .

Now let us choose smooth decreasing function $\phi: [0, \eta^2] \rightarrow [0, 1]$, which is equal to 0 on $[\frac{3}{4}\eta^2, \eta^2]$ and $\phi(0) = 1$. We define now a new function $b: U_2 \rightarrow [0, 1]$ by the formula

$$(3.7) \quad b(x, y, \vec{u}) = s(x, y, \vec{u}) \cdot \phi(\|\vec{u}\|^2).$$

Let us finally define the function $G: \Omega \rightarrow [0, 1]$ by

$$(3.8) \quad G(w) = \begin{cases} F(w) & \text{if } w \notin U_2 \\ y^3 - yx^2 + y - (\delta + 1)b(x, y, \vec{u})y + \frac{1}{2} + \vec{u}^2 & \text{if } w = (x, y, \vec{u}) \in U_2, \end{cases}$$

where $\delta > 0$ is a very small number. Later we shall show that it is enough to take $\delta < \varepsilon^2/2$. In the following lemmas we shall prove that G satisfies the conditions of Theorem 3.1.

Lemma 3.9. *The function G is smooth.*

Proof. It is a routine checking and we leave it for the reader. \square

In the next two lemmas we show that G has no critical points in U_{21} .

Lemma 3.10. *G has no critical points on $U_{21} \cap \{y = 0\}$.*

Proof. If $(x, 0, u_1, \dots, u_{n-1}) \in U_{21}$ then $x > 3$. Consider the derivative over y of G :

$$(3.11) \quad \frac{\partial G}{\partial y} = 3y^2 - x^2 + 1 - (\delta + 1)b - (\delta + 1)\phi(u_1^2 + \dots + u_{n-1}^2) \frac{\partial s}{\partial y} y.$$

Taking $y = 0$ we get $-x^2 + 1 - (\delta + 1)b$. Since $b \in [0, 1]$ and $x > 3$, one gets $\frac{\partial G}{\partial y} < 0$. \square

Lemma 3.12. *If $\delta < 3\varepsilon^2$, then G has no critical points on $U_{21} \cap \{y \neq 0\}$.*

Proof. Assume that $\frac{\partial G}{\partial x} = 0$ for some (x, y, \vec{u}) . Then

$$y \left(-2x - (\delta + 1) \frac{\partial s}{\partial y} \phi \right) = 0.$$

As $y \neq 0$, the expression in parentheses should be zero. If $0 < x \leq 3$, then by (S3) we have $\frac{\partial s}{\partial x} = 0$. Hence the above equality can not hold. Assume that $x = 0$. In the derivative over y (see equation (3.11)), the expression $-(\delta + 1)\frac{\partial s}{\partial y} y \cdot \phi$ is non-negative by (S3). Furthermore $b < 1$, hence

$$\frac{\partial G}{\partial y} \geq 3y^2 - \delta.$$

Now if $\delta < 3\varepsilon^2$ then there is no critical points with $x = 0$. It remains to deal with the case $(x, y, u_1, \dots, u_{n-1}) \in S_2$. Consider the derivative $\frac{\partial G}{\partial y}$. By (S4) we have

$$\left| y \frac{\partial s}{\partial y} \right| = \left| \frac{y^2}{r} \cdot \frac{\partial s}{\partial r} \right| + \left| \frac{(x-3)y}{r^2} \cdot \frac{\partial s}{\partial \phi} \right| < r \frac{2}{\varepsilon} + \varepsilon < 5.$$

Furthermore $|1 - (\delta + 1)b| \leq 1$, and $|3y^2| < 1$ because ε is small. As $x \geq 3$, we have $\frac{\partial G}{\partial y} < 0$ on S_2 . \square

On U_1 the function G is given by

$$(3.13) \quad G(x, y, \vec{u}) = y^3 - yx^2 - \delta y + \vec{u}^2 + \frac{1}{2}.$$

As in Section 3.1 we study the critical points in U_1 .

Lemma 3.14. *G has two critical points on U_1 at*

$$\begin{aligned} z^s &:= (0, \sqrt{\delta/3}, 0, \dots, 0) \\ z^u &:= (0, -\sqrt{\delta/3}, 0, \dots, 0). \end{aligned}$$

Both critical points are boundary, both of Morse indices k , z^s is stable, while z^u is unstable.

Proof. The derivative of G vanishes only at z^s and z^u . Indices are immediately computed from (3.13). The point z^s is boundary stable, because for z^s the expression $-yx^2$ is negative and the boundary is given by $x = 0$, hence it is attracting in the normal direction. Similarly we prove for z^u . See also Figure 10 for the two-dimensional picture. \square

Remark 3.15. If we define $G_t = y^3 - yx^2 + y - t(\delta + 1)b \cdot y + \frac{1}{2} + \vec{u}^2$ for $t \in [0, 1]$, then the same argument as in Lemmas 3.10 and 3.12 shows that G_t has no critical points in $U_2 \setminus U_1$. As for critical points in U_1 , observe that on U_1 we have

$$G_t = y^3 - yx^2 + (1 - t(1 + \delta))y + \frac{1}{2} + \vec{u}^2.$$

Let $t_0 = \frac{1}{1+\delta}$. If $t > t_0$, the function G_t has two critical points on the boundary Y , while for $t < t_0$, G_t has a single critical point in the interior $U_1 \setminus Y$. If $t = t_0$, G_t has a single degenerate critical point on Y . In this way we construct an ‘isotopy’ between F and G .

Let us now choose a Riemannian metric g' on

$$U'_1 := U_1 \cap \{||\vec{u}|| < \varepsilon\}$$

by the condition that $(x, y, u_1, \dots, u_{n-1})$ be orthonormal coordinates. Clearly, any metric g on Ω can be changed near U_1 so as to agree with g' on U'_1 . In this metric the gradient of G is

$$(-2xy, 3y^2 - x^2 - \delta, 2\epsilon_1 u_1, \dots, 2\epsilon_{n-1} u_{n-1}).$$

We want to show that there is a single trajectory starting from z^s and terminating at z^u . Clearly, there is one trajectory from z^s to z^u which stays in U'_1 (having $y = 0$ and $\vec{u} = 0$). In order to eliminate the others, we need the following lemma.

Lemma 3.16. *Let γ be a trajectory of ∇G starting from z^s . Let w be the point, where γ hits $\partial U'_1$ for the first time. If δ is sufficiently small, then $G(w) > G(z^u)$.*

Proof. Assume that $\gamma(t)$ is such trajectory. Assume that among numbers ϵ_i , we have $\epsilon_i = -1$ for $i \leq k-1$ and $\epsilon_i = 1$ otherwise. As z^s is a critical point of the vector field ∇G with a non-degenerate linear part, we conclude that the limit

$$\lim_{t \rightarrow -\infty} \frac{\gamma'(t)}{||\gamma'(t)||} =: v = (x_0, y_0, u_{01}, \dots, u_{0,n-1})$$

exists. The vector v is the tangent vector to the curve γ at the point z^s , and it lies in the unstable space. Hence $x_0 = 0$ as $(1, 0, \dots, 0)$ is a stable direction; similarly $u_{01} = \dots = u_{0,k-1} = 0$. Therefore, until γ hits the boundary of U'_1 for the first time, we have

$$x = u_1 = \dots = u_{k-1} = 0.$$

Set also $g(y) = y^3 - \delta y$. One has the following cases, depending the position of w , where γ hits $\partial U'_1$ for the first time: (a) $y = -\varepsilon$, (b) $y = \varepsilon$, or (c) $||\vec{u}||^2 = \eta^2$. The case (a) cannot happen since G is increasing along the trajectory, hence $G(w) > G(z^s)$, a fact which contradicts $g(-\varepsilon) < g(\sqrt{\delta/3})$ valid for $2\delta < \varepsilon^2$. In case (b), $G(w) > G(z^u)$ follows from $g(\varepsilon) > g(-\sqrt{\delta/3})$. Finally, assume the case (c). Then, as $u_{01} = \dots = u_{0,k-1} = 0$, we obtain $\vec{u}^2 = ||\vec{u}||^2 = \eta^2$. Then $G(w) - G(z^s) \geq \eta^2$, because the contribution to G from $y^3 - \delta y$ increases along γ . Hence $G(w) > G(z^u)$ follows again since $\varepsilon \ll \eta$. \square




FIGURE 12. Sets Z_1 and Z_2 from Section 3.3. There are drawn also the stable and unstable manifolds of ∇A .

Given the above lemma it is clear that if a trajectory γ leaves U'_1 , then G becomes bigger than $G(z^u)$. As G increases along any trajectory, it is impossible that such trajectory limits in z^u . The proof of Theorem 3.1, up to Proposition 3.5, is accomplished.

3.3. An auxiliary construction. We provide now a construction, a crucial ingredient in the proof of Proposition 3.5, see next Section. Set

$$Z = \{(x, y) \in \mathbb{R}^2 : x \geq 0\},$$

and define the two functions

$$(3.17) \quad A(x, y) = \frac{x^3}{3\sqrt{3}} - \sqrt{3}xy^2 - \frac{x}{\sqrt{3}} + \frac{2}{3\sqrt{3}}, \quad B(x, y) = y^3 - yx^2 + y.$$

They are the real/imaginary parts of a complex function; indeed (with $i = \sqrt{-1}$)

$$A + iB = \left(\frac{x}{\sqrt{3}} - iy \right)^3 - \left(\frac{x}{\sqrt{3}} - iy \right) + \frac{2}{3\sqrt{3}}.$$

It follows from the Cauchy–Riemann equations (or straightforward computations), that the level sets of A and B are orthogonal, away from the point $(1, 0)$. If we choose a Riemannian metric such that (x, y) are orthonormal coordinates, then A is constant on trajectories of ∇B (i.e. A is a first integral of ∇B). Let us fix $\delta > 0$ smaller than $\frac{2}{3\sqrt{3}}$. Consider two sets

$$(3.18) \quad Z_1 = \{(x, y) \in Z, x < 1, A(x, y) \geq \delta\}, \quad Z_2 = \{(x, y) \in Z, x > 1, A(x, y) \geq \delta\}.$$

We have the following result.

Lemma 3.19. *The map $\psi(x, y) = (A(x, y), B(x, y))$ maps Z_1 and Z_2 diffeomorphically onto E_1 and E_2 respectively, where*

$$E_1 = \left\{ (a, b) \in \mathbb{R}^2 : a \in \left[\delta, \frac{2}{3\sqrt{3}} \right] \right\}, \quad E_2 = \left\{ (a, b) \in \mathbb{R}^2 : a \geq \delta \right\}.$$

Moreover, ψ maps trajectories of ∇B onto vertical lines.

Proof. This follows from the covering theory of one-variable complex functions. Alternatively, one readily checks that $\psi: Z_1 \rightarrow V_1$ and $\psi: Z_2 \rightarrow V_2$ are bijections. As $D\psi$ is non-degenerate on $Z_1 \cup Z_2$, ψ is a diffeomorphism between the two pairs of sets. Now A is a first integral of the flow ∇B , so each trajectory of ∇B lies in the set $A^{-1}(c)$. \square




FIGURE 13. Notation used in Section 3.4. The top line is the picture on Ω , the middle line is in coordinates such that F is equal to $y^3 - yx^2 + y + \frac{1}{2} + \vec{u}^2$. The bottom line is in coordinates such that $F = b + \frac{1}{2}$. There is no mistake, the line $a = \delta$ appears twice on the picture, in coordinates on C_1 and on C_2 .

3.4. Proof of Proposition 3.5. First, as z is a critical point of index $k \in \{1, \dots, n\}$, by Morse lemma we can find a neighbourhood \tilde{V} of z and a chart $h_1: \tilde{V} \rightarrow \mathbb{R}^{n+1}$, with coordinates (x', y, \vec{u}) such that

$$F \circ h_1^{-1}(x', y, \vec{u}) = x'y + \vec{u}^2 + \frac{1}{2}.$$

Let us define a map $h_2(x, y, \vec{u}) = (x', y, \vec{u})$, where $x' = y^2 + 1 - x^2$. By the inverse function theorem, h_2 is a local diffeomorphism near $(1, 0, \dots, 0)$. Shrinking \tilde{V} if needed, and considering $h_3 = h_2^{-1} \circ h_1$, we obtain $h_3(z) = (1, 0, \dots, 0)$ and

$$F \circ h_3^{-1}(x, y, \vec{u}) = y^3 - yx^2 + y + \vec{u}^2 + \frac{1}{2} = B(x, y) + \vec{u}^2 + \frac{1}{2}.$$

Let us pick now $\xi > 0$ such that the cylinder

$$V = \{|x - 1| < \xi, |y| < \xi, \|\vec{u}\| < \xi\} \subset \mathbb{R}^{n+1}$$

lies entirely in $h_3(\tilde{V})$. By shrinking \tilde{V} we may in fact assume that $h_3(\tilde{V}) = V$. If $0 < \delta \ll 1$ is sufficiently small then $A(x, 0) < \delta$ implies $|x - 1| < \xi$. Choose such a δ , and set

$$V_1 := V \cap \{x < 1, A(x, y) \geq \delta\}$$

(compare (3.18)). By Lemma 3.19 the map

$$(3.20) \quad \Psi_1(x, y, \vec{u}) = (A(x, y), B(x, y) + \vec{u}^2, \vec{u}),$$

is a diffeomorphism (being the composition of ψ and a ‘triangular’ map). Set $C_1 := \Psi_1(V_1)$ and $\tilde{V}_1 := h_3^{-1}(V_1)$. Finally, let

$$h = \Psi_1 \circ h_3.$$

Using Lemma 3.19 again we obtain that

$$F \circ h^{-1}(a, b, \vec{u}) = b + \frac{1}{2}.$$

Let $\theta > \delta$ be sufficiently closed to δ satisfying the inclusion

$$D_1 := [\delta, \theta] \times (-\theta, \theta) \times (-\theta, \theta)^{n-1} \subset C_1.$$

Let $\tilde{D}_1 = h^{-1}(D_1) \subset \tilde{V}_1$, see Figure 13.

Lemma 3.21. *If θ and δ are small enough, there is an open/closed ball \tilde{W} in Ω , containing \tilde{D}_1 , such that h extends to a diffeomorphism between \tilde{W} and $[\delta, \frac{2}{3\sqrt{3}}] \times (-\theta, \theta) \times (-\theta, \theta)^{n-1}$ with $F \circ h^{-1}(a, b, \vec{u}) = b + \frac{1}{2}$, sending points with $a = \frac{2}{3\sqrt{3}}$ to Y .*

In the proof we shall use the following result.

Lemma 3.22. *There exists a smooth curve $\gamma: [\delta, \frac{2}{3\sqrt{3}}] \rightarrow \Omega$, such that $\gamma(\frac{2}{3\sqrt{3}}) \in Y$, $\gamma(t) \in \Sigma_{1/2}$, $\gamma(t) \in \tilde{D}_1$ if and only if $t \in [\delta, \theta]$ and $h(\gamma(t)) = (t, 0, \dots, 0)$ and γ omits $\tilde{V} \setminus \tilde{V}_1$. Furthermore, γ is transverse to Y .*

Proof of Lemma 3.22. Let $p = h^{-1}(\theta, 0, \dots, 0) \in \Sigma_{1/2}$. Let $B \subset \Sigma_{1/2}$ be an open ball with centre z and $p \in \partial B$. Let Σ' be the connected component of $\Sigma_{1/2}$ containing p . We consider two cases.

If $\Sigma' \setminus B$ is connected, it is also path connected. By (3.2), there exists a path $\tilde{\gamma} \subset \Sigma' \setminus B$ joining p with a point on the boundary. We can assume that $\tilde{\gamma}$ is transverse to Y . We choose $\gamma = h^{-1}([\delta, \theta] \times \{0, \dots, 0\}) \cup \tilde{\gamma}$ (and we smooth a possible corner at p). It is clear that γ omits $\tilde{V} \setminus \tilde{V}_1$ and that we can find a parametrization of γ by the interval $[\delta, \frac{2}{3\sqrt{3}}]$.

If $\Sigma' \setminus B$ is not connected, then as Σ' is connected, by a homological argument we have $n = 1$ and $k = 1$. Since Σ' is connected and has boundary, then Σ' is an interval and B is an interval too. Then $\Sigma' \setminus B$ consists of two intervals, each intersecting Y . One of these intervals contains p . So p is connected with Y by an interval, which omits B . We conclude the proof by the same argument as in the above case, when $\Sigma' \setminus B$ was connected. \square

Proof of Lemma 3.21. Given Lemma 3.22, let us choose a tubular neighbourhood X of γ in $F^{-1}(1/2) \setminus (\tilde{V} \setminus \tilde{V}_1)$. Shrinking X if needed we can assume that it is a disk and $X_1 := X \cap \tilde{V} = \tilde{D}_1 \cap F^{-1}(1/2)$. Now let τ be a vector field on \tilde{D}_1 given by $(Dh)^{-1}(1, 0, \dots, 0)$, where Dh denotes the derivative of h . This vector field is everywhere tangent to X_1 and

$$(3.23) \quad \tau|_{\gamma \cap \tilde{D}_1} = \frac{d}{dt} \gamma(t)$$

by the very definition of γ . We extend τ to a smooth vector field on the whole X , such that (3.23) holds on the whole γ . For any point $z \in \gamma$, the trajectory of τ (which is γ) hits eventually Y and, on the other end, it hits the ‘right wall’

$$\tilde{R} = h^{-1}(\{\delta\} \times \{0\} \times (-\theta, \theta)^{n-1}).$$

(compare Figure 14; note that the horizontal coordinate there increases from right to left for consistency with Figure 13). Since γ is transverse to \tilde{R} and to Y , by implicit function




FIGURE 14. Proof of Lemma 3.21. Construction of the vector field τ . Picture on $F^{-1}(1/2)$. The parallel vector field from the region on the right is extended to the whole X so that it is tangent to γ .

theorem trajectories close to τ also start at \tilde{R} and end up at Y . Shrinking X if necessary we may assume that each point of X lies on the trajectory of τ which connects a point of \tilde{R} to some point of Y , and all the trajectories are transverse to both Y and \tilde{R} .

We can now rescale τ (that is multiply by a suitable smooth function constant on trajectories) so that all the trajectories go from \tilde{R} to Y in time $\frac{2}{3\sqrt{3}} - \delta$, i.e. the same time as γ does. The rescaled vector field allows us to introduce coordinates on X in the following way. For any point $z \in X$, let γ_z be the trajectory of τ , going through z . We can assume that $\gamma_z(\delta) \in \tilde{R}$. Let $t_z = \gamma_z^{-1}(z)$, i.e. the moment when γ_z passes through z . Since we normalized γ_z , we know that $t_z \in [\delta, \frac{2}{3\sqrt{3}}]$ and $t_z = \frac{2}{3\sqrt{3}}$ if and only if $z \in Y \cap X$.

Let \vec{u}_z be such that $h(\gamma_z(\delta)) = (\delta, 0, \vec{u}_z)$. The vector \vec{u}_z might be thought off as a coordinate on \tilde{R} . We define now

$$h(z) = (t_z, 0, \vec{u}_z).$$

This maps clearly extends h to the whole X .

Now let \tilde{W} be a tubular neighbourhood of X in $\Omega \setminus (\tilde{V} \setminus \tilde{V}_1)$. We use the flow of ∇F to extend coordinates from X to \tilde{W} . More precisely, shrinking \tilde{W} if needed we may assume that for each $w \in \tilde{W}$ the trajectory of ∇F intersects X . This intersection is necessarily transverse and it is in one point, which we denote by $z_w \in X$. We define now

$$h(w) = (t_{z_w}, F(w) - F(z_w), \vec{u}_{z_w}).$$

As h is a local diffeomorphism on X (because ∇F is transverse to X), it is also a local diffeomorphism near X . We put $W = h(\tilde{W})$. Clearly both definitions of h on \tilde{V} and \tilde{W} agree. We may now decrease θ and shrink W so that

$$W = [\delta, \frac{2}{3\sqrt{3}}] \times (-\theta, \theta) \times (-\theta, \theta)^{n-1}.$$

We have $F \circ h^{-1}(a, b, \vec{u}) = b + \frac{1}{2}$. We now extend h_3 over \tilde{W} by the formula $h_3 = \Psi^{-1} \circ h$. \square

Consider now

$$V_2 := V \cap \{x > 1, A(x, y) \geq \delta\}.$$

Let $\Psi_2: V \rightarrow \mathbb{R}^{n+1}$ be given by $\Psi_2(x, y, \vec{u}) = (a, b, \vec{u}) = (A(x, y), B(x, y) + \vec{u}^2, \vec{u})$, provided by the same formula as Ψ_1 in (3.20) but the image now satisfies $a \geq \delta$, cf. Lemma 3.19.

Let $C_2 = \Psi_2(V_2)$, and let us choose θ' sufficiently small such that

$$D_2 := [\delta, \theta'] \times (-\theta', \theta') \times (-\theta', \theta')^{n-1} \subset C_2.$$

We shall denote $h = \Psi_2 \circ h_3$ and $\tilde{D}_2 = h^{-1}(D_2)$.

Let us now fix $M > 0$ large enough and consider a map $h_4: \mathbb{R}^{n+1} \rightarrow \mathbb{R}^{n+1}$ of the form

$$h_4(a, b, \vec{u}) = (\phi(a), b, \vec{u}),$$

where $\phi: [\delta, \theta'] \cong [\delta, M]$ is a strictly increasing smooth function, which is an identity near δ . Consider the map $h'_3: \Psi_2^{-1} \circ h_4 \circ h: \tilde{D}_2 \rightarrow \mathbb{R}^{n+1}$. Since h is an identity for a close to δ , this map agrees with h_3 for a close to δ . Furthermore $F \circ h^{-1}(a, b, \vec{u}) = F \circ h^{-1} \circ h_4^{-1}(a, b, \vec{u}) = b + \frac{1}{2}$ by a straightforward computation. On the other hand, the point $h^{-1}(\theta', 0, \dots, 0) \in \tilde{D}_2$ is mapped by h'_3 to $(M, 0, \dots, 0) \in \mathbb{R}^{n+1}$, where M can be arbitrary large, e.g. $M > 3$.

Having gathered all the necessary maps, we now conclude the proof. Let

$$\tilde{U} = \tilde{W} \cup (\tilde{V} \setminus h_3^{-1}(V_1 \cup V_2)) \cup \tilde{D}_2.$$

The map $h_3: \tilde{U} \rightarrow [0, \infty) \times \mathbb{R}^n$ is given by h_3 on \tilde{W} and on $\tilde{V} \setminus h_3^{-1}(V_2)$, and by h'_3 on \tilde{D}_2 . This map is clearly a diffeomorphism onto its image, so it is a chart near z . By construction $F \circ h_3^{-1}$ is equal to $y^3 - yx^2 + y + \vec{u}^2 + 1/2$ and $h_3(\tilde{W})$ contains the segment with endpoints $(0, 0, \dots, 0)$ and $(3, 0, \dots, 0)$. Since it is an open subset, it contains $[0, 3 + \eta) \times (-\eta, \eta) \times (-\eta, \eta)^{n-1}$ for $\eta > 0$ small enough. The inverse image of this cube gives the required chart.

This ends the proof of Theorem 3.1 which moves a single interior critical point to the boundary. Section 4 generalizes this fact for more critical points; one of the needed tools will be the rearrangements of the critical values/points.

3.5. Condition (3.2) revisited. We will provide two characterizations of Condition (3.2). One is valid for arbitrary $n \geq 1$, the other one holds only in the case $n = 1$. We shall keep the notation from previous subsections, in particular (Ω, Y) is a cobordism between (Σ_0, M_0) and (Σ_1, M_1) , $F: \Omega \rightarrow [0, 1]$ is a Morse function with a single critical point z in the interior of Ω , and $F(z) = 1/2$. Let $\Sigma_{1/2} = F^{-1}(1/2)$ and Σ' be the connected component of $\Sigma_{1/2}$ such that $z \in \Sigma'$.



Proposition 3.24. *If Σ_0 , Σ_1 and Ω have no closed connected components, then $\Sigma' \cap Y \neq \emptyset$. In particular, in Theorem 3.1 we can assume that Σ_0, Σ_1 and Ω have no closed connected components instead of (3.2).*

Proof. Let $p = h^{-1}(\theta, 0, \dots, 0) \in \tilde{D}_1 \subset \Omega$ and let B be an open ball in Σ' near z , such that $p \in \partial B$. It is enough to show that p can be connected to Y by a path in $\Sigma_{1/2}$, which omits B (compare Lemma 3.22).

Let us choose a Riemannian metric on Ω . Let W_z^s be the stable manifold of z and let T be intersection of W_z^s and Σ_0 . This is a $(k-1)$ -dimensional sphere. The flow of ∇F induces a diffeomorphism $\Phi: \Sigma_{1/2} \setminus B \cong \Sigma_0 \setminus B_0$, where B_0 is a tubular neighbourhood of T in Σ_0 (here we tacitly use the fact that δ and θ are small enough), see Figure 15. Let $p_0 = \Phi(p)$. Let Σ'_0 be the connected component of Σ_0 , which contains B_0 .

Now we will analyze several cases. Recall that $k = \text{ind}_z F \in \{1, \dots, n\}$. First we assume that $k < n$. Then $\Sigma'_0 \setminus T$ is connected, so p_0 can be connected to the boundary of Σ'_0 — which is non-empty by assumptions of the proposition — by a path γ_0 . Now the inverse image $\Phi^{-1}(\gamma_0)$ is the required path.

If $k = n > 1$ then we reverse the cobordism and look at $-F$, hence this case is covered by the previous one (since $k = n$ will be replaced by $k = 1 < n$).

FIGURE 15. Notation on Σ_0 .FIGURE 16. Proof of Proposition 3.24. Case $k = 1$ and $n = 1$ and T lies in two components of Σ . Σ_0 is the horizontal segment. The points p'_0 and p''_0 are the two possible positions of the point p_0 .FIGURE 17. Proof of Proposition 3.24. Case $k = 1$ and $n = 1$ and T lies in two components of Σ_0 . The points p'_0 and p''_0 are the two possible positions of the point p_0 . Both can be connected to the boundary M_0 .

Finally, it remains to deal with the situation $k = n = 1$. Then $\dim \Sigma_0 = 1$. T consists of two points. Assume first that they lie in a single connected component Σ'_0 of Σ_0 . We shall show that this is impossible. As Σ'_0 is connected with non-trivial boundary, it is an interval. The situation is like on Figure 16. Now as F has precisely one Morse critical point of index 1, Σ_1 is the result of a surgery on Σ_0 . This surgery consists of removing two inner segments from Σ_0 and gluing back two other segments, which on Figure 16 are drawn as dashed arc. But then Σ_1 has a closed connected component, which contradicts assumptions of Theorem 3.1.

Therefore, T lies in two connected components of Σ_0 . The situation is drawn of Figure 17, and it is straightforward to see that p_0 can be connected to M_0 by a segment omitting B_0 . \square

The proof of Proposition 3.24 suggests that the case $n = 1$ is different than case $n > 1$. We shall provide now a full characterization of a failure to (3.2).

Proposition 3.25. *Assume that $k = n = 1$ and Ω is connected. If (3.2) does not hold, then Ω is a pair of pants, Σ_0 is a circle and Σ_1 is a disjoint union of two circles; or converse: $\Sigma_1 = S^1$ and Σ_0 is a disjoint union of two circles. In particular, $Y = \emptyset$.*

Proof. A one-handle attached to a surface changes the number of boundary components by ± 1 . Let us assume that Σ_1 has less components than Σ_0 , if not we can reverse the cobordism. As Ω is connected, Σ_0 has two components and Σ_1 only one. Let $A_0 \subset \Sigma_0$ be the attaching region, i.e. the union of two closed intervals to which the one-handle is attached. With the notation of Section 3.5 we have $(\Sigma', z) \cong (\Sigma_0/A_0, A_0/A_0)$, where the quotient denotes collapsing a space to a point. In particular z can not be joined to Y by a path in Σ' if and only if Σ_0 is disjoint from Y . Hence Σ_0 is closed, that is, it is a union of two circles. \square

4. REARRANGEMENTS OF BOUNDARY HANDLES

4.1. Preliminaries. Let (Ω, Y) be a cobordism between two n -dimensional manifolds with boundary (Σ_0, M_0) and (Σ_1, M_1) . Let F be a Morse function, with critical points $w_1, \dots, w_k \in \text{Int } \Omega$ and $y_1, \dots, y_l \in Y$. In the classical theory (that is, when $Y = \emptyset$), the Thom–Milnor–Smale theorem (see [Mi2, Section 4]) says that we can alter F without introducing new critical points such that if $\text{ind } w_i < \text{ind } w_j$, then $F(w_i) < F(w_j)$ as well. We want to prove similar results in our more general case.

In this section we rely very strongly on [Mi2, Section 4].

4.2. Elementary rearrangement theorems. We shall begin with the case $k + l = 2$, i.e. F has two critical points. For a critical point p we shall denote by K_p the union $W_p^s \cup \{p\} \cup W_p^u$, i.e. the set of all points $x \in \Omega$, such that the trajectory $\phi_t(x)$ ($t \in \mathbb{R}$), of the gradient vector field ∇F has p as its limit set. Elementary rearrangement theorems deal with the case when the two sets K_{p_1} and K_{p_2} for the two critical points are disjoint.

Proposition 4.1 (Rearrangement of critical points). *Let p_1 and p_2 be two critical points, and assume that $K_1 := K_{p_1}$ and $K_2 := K_{p_2}$ are disjoint. Let us choose $a_1, a_2 \in (0, 1)$. Then, there exist a Morse function $G: \Omega \rightarrow [0, 1]$, with critical points exactly at p_1 and p_2 , such that $G(p_i) = a_i$, $i = 1, 2$; furthermore, near p_1 and p_2 , the difference $F - G$ is a constant function.*

Remark 4.2. If both critical points are on the boundary, in order to guarantee the above existence, we need even to change the Riemannian metric away from K_1 and K_2 .

Proof. Similarly as in [Mi2, Section 4] we first prove an auxiliary result.

Lemma 4.3. *There exists a smooth function $\mu: \Omega \rightarrow [0, 1]$ with the following properties:*

- (M1) $\mu \equiv 0$ in a neighbourhood of K_1 ;
- (M2) $\mu \equiv 1$ in a neighbourhood of K_2 ;
- (M3) μ is constant on trajectories of ∇F .

Furthermore, if at least one of the critical points is interior, we have

- (M4) μ is constant on Y .

Given the lemma, we choose also a smooth function $\Psi: [0, 1] \times [0, 1] \rightarrow [0, 1]$ with

- (PS1) $\frac{\partial \Psi}{\partial x}(x, y) > 0$ for all $(x, y) \in [0, 1] \times [0, 1]$;
- (PS2) there exists $\delta > 0$, such that $\Psi(x, y) = x$ for all $x \in [0, \delta] \cup [1 - \delta, 1]$ and $y \in [0, 1]$;
- (PS3) for any $s \in (-\delta, \delta)$ we have $\Psi(F(p_1) + s, 0) = a_1 + s$ and $\Psi(F(p_2) + s, 1) = a_2 + s$.

For any $\eta \in \Omega$ we define $G(\eta) = \Psi(F(\eta), \mu(\eta))$. From the properties (PS3), (M1) and (M2) we see that near p_i , G differs from F by a constant. The property (PS2) ensures that G agrees with F in a neighbourhood of Σ_0 and Σ_1 . Let us show that ∇G does not vanish away from p_i . By a chain rule we have

$$(4.4) \quad \nabla G = \frac{\partial \Psi}{\partial x} \nabla F + \frac{\partial \Psi}{\partial y} \nabla \mu.$$

Since μ is constant on all trajectories of ∇F , the scalar product $\langle \nabla F, \nabla \mu \rangle = 0$. Then the property (PS1) guarantees that $\langle \nabla G, \nabla F \rangle > 0$ away from p_1 and p_2 .

We need to show that ∇G is everywhere tangent to Y . If one of the points is interior, by (M4) $\nabla \mu$ vanishes on Y , hence ∇G is parallel on Y to ∇F and we are done. Next assume that both critical points are on the boundary. Let us choose an open subset U of Y such that $\nabla \mu|_U = 0$ and $K_1 \cup K_2 \subset U$. This is possible, because of the properties (M1) and (M2). Then let us choose a neighbourhood W in Ω of $Y \setminus U$, disjoint from K_1 and K_2 . Observe that $dG(\nabla F) = \langle \nabla G, \nabla F \rangle > 0$. As $\nabla F \in TY$ one has $\ker dG \not\subset TY$, so by Lemma 1.7 we can change the metric in W so that ∇G is everywhere tangent to Y . \square

Proof of Lemma 4.3. Let us define $T_1 = K_1 \cap \Sigma_0$ and $T_2 = K_2 \cap \Sigma_0$. Assume that T_1 and T_2 are not empty. For each $\eta \in \Omega \setminus K_1 \cup K_2$, let $\pi(\eta)$ be the intersection of the trajectory of η under ∇F with Σ_0 . This gives a map $\pi: \Omega \setminus (K_1 \cup K_2) \rightarrow \Sigma_0 \setminus (T_1 \cup T_2)$.

Let us define μ first on Σ_0 by the following conditions: $\mu \equiv 1$ in a neighbourhood of T_2 , $\mu \equiv 0$ in a neighbourhood of T_1 . Furthermore, if either T_1 or T_2 is disjoint from the boundary M_0 we extend μ to a constant function on M_0 . Finally, we extend μ to the whole Ω by picking $\mu(\eta) = \mu(\pi(\eta))$ if $\eta \notin K_1 \cup K_2$, and $\mu|_{K_i}(\eta) = i - 1$, $i = 1, 2$.

If $T_1 = \emptyset$, then $\text{ind}_F p_1 = 0$ and the proof of the rearrangement theorem is completely straightforward. \square

4.3. Morse–Smale condition on manifolds with boundary. In the classical theory, the Morse–Smale condition imposed on a Morse function $F: M \rightarrow \mathbb{R}$ means that for each pair of two critical points p_1, p_2 of M the intersection of stable manifold $W_{p_1}^s$ with the unstable manifold of $W_{p_2}^u$ is transverse. (Note that this Morse–Smale condition also depends on the choice of Riemannian metric on M .) Following [KM, Definition 2.4.2], we reformulate the Morse–Smale condition in the following way

Definition 4.5. The function F is called *Morse–Smale* if for any two critical points p_1 and p_2 , the intersection of $\text{Int } \Omega \cap W_{p_1}^s$ with $\text{Int } \Omega \cap W_{p_2}^u$ is transverse (as the intersection in the $(n+1)$ -dimensional manifold Ω) and the intersection of $Y \cap W_{p_1}^s$ with $Y \cap W_{p_2}^u$ is transverse (as an intersection in the n -dimensional manifold Y).

It is clear, that regular functions form an open-dense subset of all C^2 functions, which satisfy the Morse condition from Definition 1.4.

Assume now that F is Morse–Smale. Given two critical points of F , p_1 and p_2 , we want to check whether $W_{p_1}^s \cap W_{p_2}^u = \emptyset$. This depends not only on the index, but also on whether any of the two points is boundary stable. We show this in a tabulated form in Table 1, where $\text{ind } p_1 = k$ and $\text{ind } p_2 = l$. In studying the intersection, we remark that $W_{p_1}^s \cap W_{p_2}^u$ is formed from trajectories, so if for dimensional reasons we have $\dim W_{p_1}^s \cap W_{p_2}^u < 1$, it immediately follows that this intersection is empty.

4.4. Global rearrangement theorem. Let us combine the rearrangement theorems from Section 4.2 with the computations in Table 1.

Proposition 4.6. *Let F be a Morse function on a cobordism (Ω, Y) between (Σ_0, M_0) and (Σ_1, M_1) . Let w_1, \dots, w_m be the interior critical points of F and y_1, \dots, y_{k+l} be the boundary*

type of p_1	type of p_2	$\dim_\Omega W_{p_1}^s$	$\dim_Y W_{p_1}^s$	$\dim_\Omega W_{p_2}^u$	$\dim_Y W_{p_2}^u$	empty if
interior	interior	k	\emptyset	$n+1-l$	\emptyset	$k \leq l$
interior	b. stable	k	\emptyset	\emptyset	$n+1-l$	always
interior	b. unstable	k	\emptyset	$n+1-l$	$n-l$	$k \leq l$
b. stable	interior	k	$k-1$	$n+1-l$	\emptyset	$k \leq l$
b. stable	b. stable	k	$k-1$	\emptyset	$n-l$	$k \leq l$
b. stable	b. unstable	k	$k-1$	$n+1-l$	$n-l$	$k \leq l$
b. unstable	interior	\emptyset	k	$n+1-l$	\emptyset	always
b. unstable	b. stable	\emptyset	k	\emptyset	$n+1-l$	$k < l$
b. unstable	b. unstable	\emptyset	k	$n+1-l$	$n-l$	$k \leq l$

TABLE 1. Under the Morse–Smale condition, the last column shows, whether there might exist trajectories from z_2 to z_1 . We write $\dim_\Omega W^s = \dim(W^s \cap \text{Int } \Omega)$ and $\dim_Y W^s = \dim(W^s \cap Y)$. \emptyset means that the manifold in question is empty.

critical points, where y_1, \dots, y_k are boundary stable and y_{k+1}, \dots, y_{k+l} boundary unstable. Let us choose real numbers $0 < c_0 < c_1 < \dots < c_{n+1} < 1$, $0 < c_1^s < \dots < c_{n+1}^s < 1$, $0 < c_0^u < \dots < c_n^u < 1$ satisfying

$$(4.7) \quad \begin{aligned} c_{i-1}^s &< c_i < c_{i+1}^s \\ c_{i-1}^u &< c_i < c_{i+1}^u \\ c_{i-1}^u &< c_i^s < c_i^u \end{aligned}$$

for all $i \in \{0, \dots, n+1\}$ (we can assume that $c_{-1} = c_0^s = c_{-1}^u = 0$, $c_{n+2} = c_{n+2}^s = c_{n+1}^u = 1$ so that (4.7) makes sense for all i).

Then, there exists another Morse function G on the cobordism (Ω, Y) with critical points w_1, \dots, w_m in the interior, y_1, \dots, y_{k+l} on the boundary, such that if $\text{ind } w_j = l$, then $G(w_j) = c_l$, if $\text{ind } y_i = l$ and y_i is boundary stable then $G(y_i) = c_l^s$, and if $\text{ind } y_i = l$ and y_i is boundary unstable, then $G(y_i) = c_l^u$.

Proof. Given the elementary rearrangement result (Proposition 4.1), the proof is completely standard (see the proof of Theorem 4.8 in [Mi2]). Note only that we need to have $c_i^s < c_i^u$ in the statement, because there might be a trajectory from a boundary stable critical point to a boundary unstable of the same index. However, we are free to choose $c_i < c_i^s$ or $c_i \in (c_i^s, c_i^u)$ or $c_i > c_i^u$. \square

4.5. Moving more handles to the boundary at once. Before we formulate Theorem 4.10, let us introduce the following technical notion.

Definition 4.8. The Morse function F on the cobordism (Ω, Y) is called *technically good* if it has the following properties.

- (TG1) If p_1, p_2 are (interior or boundary) critical points of F then $\text{ind } p_1 < \text{ind } p_2$ implies $F(p_1) < F(p_2)$;
- (TG2) There exist regular values of F , say $c, d \in [0, 1]$, with $c < d$ such that $F^{-1}[0, c]$ contains those and only those critical points, which have index 0 or which are boundary stable critical points of index 1, and $F^{-1}[d, 1]$ contains those and only those critical points, which have index $n+1$ and boundary unstable critical points of index n .
- (TG3) There are no pairs of 0 and 1 (interior) handles of F that can be cancelled (in the sense of Section 5);
- (TG4) There are no pairs of n and $n+1$ (interior) handles of F that can be cancelled.

Lemma 4.9. *Each function F can be made technically good without introducing new critical points.*

Proof. By Proposition 4.6 we can rearrange the critical points of F , proving (TG1) and (TG2). The properties (TG3) and (TG4) can be guaranteed, using the handle cancellation theorem (e.g. [Mi2, Theorem 5.4]). We refer to the beginning of Section 5 for an explanation, that one can use the handle cancellation theorem if the manifold in question has boundary. \square

Theorem 4.10. *Let (Ω, Y) be a cobordism between (Σ_0, M_0) and (Σ_1, M_1) . Let F be a technically good Morse function on that cobordism, which has critical points y_1, \dots, y_k on the boundary Y , z_1, \dots, z_{l+m} in the interior $\text{Int } \Omega$, of which z_{m+1}, \dots, z_{l+m} have index 0 or $n+1$ and the indices of z_1, \dots, z_m are in $\{1, \dots, n\}$. Suppose furthermore the following properties are satisfied:*

- (I1) Σ_0 and Σ_1 have no closed connected components;
- (I2) Ω has no closed connected component.

Then, there exist a Morse function $G: \Omega \rightarrow [0, 1]$, on the cobordism (Ω, Y) , with critical points $y_1, \dots, y_k \in Y$, z_{m+1}, \dots, z_{l+m} and $z_1^s, z_1^u, \dots, z_m^s, z_m^u$ such that:

- $\text{ind}_G y_i = \text{ind}_F y_i$ for $i = 1, \dots, k$ and for $j = m+1, \dots, m+l$ we have $\text{ind}_G z_j = \text{ind}_F z_j$;
- for $j = 1, \dots, m$, $\text{ind}_G z_j^s = \text{ind}_G z_j^u = \text{ind}_F z_j$;
- for $j = 1, \dots, m$, z_j^s and z_j^u are on the boundary Y , furthermore z_j^s is boundary stable, z_j^u is boundary unstable and $G(z_j^s) < G(z_j^u)$.

In other words, we can move all critical points to the boundary at once. To prove Theorem 4.10 we use Theorem 3.1 independently for each critical point z_1, \dots, z_m . We need to ensure that Condition 3.2 holds. This is done in Proposition 4.11 stated below. Given these two ingredients the proof is straightforward.

4.6. Topological ingredients needed in the proof of Theorem 4.10.

Proposition 4.11. *Let F be a technically good Morse function on the cobordism (Ω, Y) . Assume that Σ_0, Σ_1 and Ω have no closed connected components. Let c, d be as in Definition 4.8. Then*

- (a) *If $n > 1$, then for any $y \in [c, d]$, the inverse image $F^{-1}(y)$ has no closed connected component.*
- (b) *If $n = 1$, then after possibly rearranging the critical values of the interior critical points of index 1, for any interior critical point $z \in \Omega$ of F of index 1, z can be connected with Y by a curve lying entirely in $F^{-1}(F(z))$; and furthermore all the critical points are on different levels.*

Remark 4.12. The distinction between cases $n > 1$ and $n = 1$ is necessary. Point (a) of Proposition 4.11 is not necessarily valid if $n = 1$, see Figure 18 for a simple counterexample.

First let us prove several lemmas, which are simple consequences of the assumptions of Proposition 4.11. We use assumptions and notation of Proposition 4.11.

Lemma 4.13. *Let $x, y \in [0, 1]$ with $x < y$. If Ω' is a connected component of $F^{-1}[x, y]$ then either $\Omega' \cap Y = \emptyset$, or for any $u \in [x, y] \cap [c, d]$ we have $F^{-1}(u) \cap \Omega' \cap Y \neq \emptyset$.*

Proof. Assume that for some $u \in [x, y] \cap [c, d]$ the intersection $F^{-1}(u) \cap \Omega' \cap Y = \emptyset$ and $\Omega' \cap Y \neq \emptyset$. Then either $\Omega' \cap Y \cap F^{-1}[0, u]$ or $\Omega' \cap Y \cap F^{-1}[u, 1]$ is not empty. Assume the first possibility (the other one is symmetric) and let $Y' = \Omega' \cap Y \cap F^{-1}[0, u]$. Let $f = F|_{Y'}$.




FIGURE 18. The statement of Proposition 4.11(a) does not hold if $n = 1$. Here, F is the height function. The level set $F^{-1}(1/2)$, drawn on the picture, has two connected components, one of which is closed.

be the restriction. Then Y' is compact and f has a local maximum on Y' . This maximum corresponds to a critical point of f of index n , so either a boundary stable critical point of F of index $n + 1$, or a boundary unstable critical point of index n . But the corresponding critical value is smaller than u , so smaller than d , which contradicts property (TG2). \square

Lemma 4.14. *For any $x \in [c, 1]$ the set $F^{-1}[0, x]$ cannot have a connected component disjoint from Y .*

Proof. Assume the contrary, and let Ω' be a connected component of $F^{-1}[0, x]$ disjoint from Y . Let Ω_1 be the connected component of Ω containing Ω' . Suppose that $\Omega_1 \cap Y = \emptyset$. Then either $\Omega_1 \cap (\Sigma_0 \cup \Sigma_1) = \emptyset$ or $\Omega_1 \cap (\Sigma_0 \cup \Sigma_1) \neq \emptyset$. In the first case Ω_1 is a closed connected component of Ω , in the second either $\Omega_1 \cap \Sigma_0$, or $\Omega_1 \cap \Sigma_1$ is not empty, so either Σ_0 or Σ_1 has a closed connected component. The contradiction implies that $\Omega_1 \cap Y \neq \emptyset$.

By Lemma 4.13 we have then $F^{-1}(x) \cap \Omega_1 \cap Y \neq \emptyset$, hence $\Omega'' := (F^{-1}[0, x] \cap \Omega_1) \setminus \Omega'$ is not empty and is disjoint from Ω' . As Ω' and Ω'' both belong to Ω_1 which is connected, there must be a critical point $z \in \Omega_1$ of index 1, which joins Ω' to Ω'' . We have $F(z) > x$ and Ω', Ω'' belong to two different connected components of $F^{-1}[0, F(z))$ and to a single connected component of $F^{-1}[0, F(z)]$. The connected component of $F^{-1}[0, F(z))$ containing Ω' has empty intersection with Y (by Lemma 4.13) hence z must be an interior critical point of index 1. We also remind that all critical points of F on Ω' are interior critical points, because $\Omega' \cap Y = \emptyset$.

Let W^s be the stable manifold of z of the vector field ∇F . Then $W^s \cap \Omega_1$ must be a connected curve, with non-empty intersection with Ω' . One of its boundaries is either on $\Sigma_0 \cap \Omega'$ or it is a critical point of F in Ω' , necessarily interior and by the Morse–Smale condition, its index is 0. In the first case, $\Sigma_0 \cap \Omega'$ is not empty and since it is disjoint from Y , Σ_0 has a closed connected component. In the other case, we have in Ω_1 a single trajectory between a critical point of index 0 and a critical point of index 1. This contradicts (TG3). \square

Lemma 4.15. *Assume that for some $y \in [c, d]$ Σ_1 and Σ_2 are two disjoint connected components of $F^{-1}(y)$. If F has no interior or boundary unstable critical points of index n*

with critical value in $[c, y)$, then Σ_1 and Σ_2 belong to two different connected components of $F^{-1}[0, y]$.

Proof. For $x < y$ and close to y the sets Σ_1 and Σ_2 lie in two different connected components of $F^{-1}(x, y]$. Let $x_0 \geq 0$ be the smallest x with that property. Assume $x_0 > 0$. Then x_0 is a critical value of F . The number of connected components of $F^{-1}(x)$ increases as x crosses x_0 . Thus the corresponding critical point is either an interior critical point of index n , or a boundary unstable critical point of index n . But then $x_0 \geq c$ because of (TG2), so we have $x_0 \in [c, y]$ which contradicts the assumptions of the lemma.

It follows that $x_0 = 0$. As F has no critical points on Σ_0 , it follows that Σ_1 and Σ_2 belong to different components of $F^{-1}[0, y]$. \square

Lemma 4.16. *Let $y \in [c, d]$ be chosen so that there are no interior or boundary unstable critical points of index n with critical values in $[c, y)$. Then $F^{-1}(y)$ has no closed connected components.*

Proof. Assume that Σ' is a closed connected component of $F^{-1}(y)$. Let $\Sigma'' = F^{-1}(y) \setminus \Sigma'$, it is not empty by Lemma 4.13 (applied for $x = 0, y = 1$), for otherwise $\Omega \cap Y = \emptyset$. Let Ω' be the connected component of $F^{-1}[0, y]$ containing Σ' . By Lemma 4.15, Ω' and Σ'' are disjoint, in particular $\Omega' \cap F^{-1}(y) \cap Y = \Sigma' \cap Y = \emptyset$. By Lemma 4.13, $\Omega' \cap Y = \emptyset$. But this contradicts Lemma 4.14. \square

Remark 4.17. There exists a symmetric formulation of the last three lemmas, which can be obtained by considering the function $1 - F$ instead of F . For instance, in Lemma 4.16, the symmetric assumption is that there are no interior or boundary stable critical points of index 1 in $(y, d]$. The statement is the same.

Proof of Proposition 4.11. Case $n > 1$. Let $x \in [c, d]$ be a non-critical value such that all the critical points of F with index n have critical value greater than x , and all critical points with index smaller than n have critical values smaller than x . Such x exists because of (TG1). If $y \leq x$, then Lemma 4.16 guarantees that $F^{-1}(y)$ has no closed connected components. If $y > x$, then $F^{-1}[y, d]$ has no critical points of index 1 (as $n > 1$), so we apply the symmetric counterpart of Lemma 4.16. \square

Proof of Proposition 4.11. Case $n = 1$. The property (TG2) implies that the only critical points of $F|_{[c, d]}$ are the interior critical points of index 1. Let us call them z_1, \dots, z_m . Since they are all of the same index, by Proposition 4.6 we are able to rearrange the values $F(z_1), \dots, F(z_m)$ at will. Let us fix c_1, \dots, c_m with the property that $c < c_1 < \dots < c_m < d$. Let us first rearrange the points z_1, \dots, z_m so that $F(z_1) = \dots = F(z_m) = c_1$.

The singular level set $F^{-1}(c_1)$ is a manifold with m singular points z_1, \dots, z_m , which are double points. By Lemma 4.16, $F^{-1}(c_1)$ has no closed connected components. In particular each of the points z_1, \dots, z_m , can be connected to Y by a curve lying in $F^{-1}(c_1)$. At least one of those points can be connected to Y by a curve γ , which omits all the other critical points. We relabel the critical points so that this point is z_m . We rearrange the critical points so that $F(z_m) = c_m$ and the value $F(z_1) = \dots = F(z_{m-1}) = c_1$. By construction, z_m can be connected with Y by a curve lying in $F^{-1}(c_m)$.

The procedure now is repeated, i.e. assume that we have already moved z_{k+1}, \dots, z_m to levels c_{k+1}, \dots, c_m respectively. Then $F^{-1}(c_1)$ still has no closed connected components by Lemma 4.16. We assume that z_k can be connected to Y by a curve in $F^{-1}(c_1)$ omitting all the other critical points. Then we rearrange the critical values so that $F(z_k) = c_k$. The proof is accomplished by an inductive argument. \square

4.7. Splitting of cobordisms. We have now all the ingredients needed to prove our main theorem about splitting cobordisms. We slightly change the notation in this subsection, the cobordism will be between (Σ, M) and (Σ', M') .

Theorem 4.18. *Let (Ω, Y) be a cobordism between (Σ, M) and (Σ', M') . If the following conditions are satisfied*

- Σ and Σ' have no closed connected components;
- Ω has no closed connected component;

Then the relative cobordism can be expressed as a union

$$\Omega = \Omega_0 \cup \Omega_{1/2} \cup \Omega_1 \cup \Omega_{3/2} \cup \cdots \cup \Omega_{n+1/2} \cup \Omega_{n+1}$$

such that $\partial\Omega_s = \Sigma_s \cup \Sigma_{s+1/2} \cup Y_s$ with $\Sigma_0 = \Sigma$, $\Sigma_{n+3/2} = \Sigma'$, $Y = Y_0 \cup \cdots \cup Y_{n+1}$. In other words (Ω_s, Y_s) is a cobordism between (Σ_s, M_s) and $(\Sigma_{s+1/2}, M_{s+1/2})$, where $M_s = \partial\Sigma_s = \Sigma_s \cap Y_s$. Furthermore

- (Ω_0, Y_0) is a cobordism given by a sequence of index 0 handle attachments;
- for $k = 1, \dots, n+1$, $(\Omega_{k-1/2}, Y_{k-1/2})$ is a left product cobordism, given by a sequence of index k left half-handle attachments;
- for $k = 1, \dots, n$, (Ω_k, Y_k) is a right product cobordism, given by a sequence of index k right half-handle attachments;
- (Ω_{n+1}, Y_{n+1}) is a cobordism provided by a sequence of index $(n+1)$ handle attachments.

Proof. Let us begin with a Morse function F on the cobordism which has only boundary stable critical points (see Remark 2.3). Assume that w_1, \dots, w_m are the interior critical points and y_1, \dots, y_k are the boundary critical points. After a rearrangement of critical points and the cancellation of pairs of critical points as in Lemma 4.9 we can make F technically good. After applying Theorem 4.10 we get that F can have only 0 handles and $n+1$ handles as interior handles. Let us write $\theta = 1/(4n+6)$ and choose $c_0 = \theta$, $c_1^s = 3\theta$, $c_1^u = 5\theta$, \dots , $c_k^s = (4k-1)\theta$, $c_k^u = (4k+1)\theta$, \dots , $c_{n+1}^s = 1-3\theta$, $c_{n+1} = 1-\theta$. We rearrange the function F according to Proposition 4.6. Then we define for $k = 0, 1/2, 1, \dots, n+1$ the manifold $\Omega_k = F^{-1}[4k\theta, (4k+2)\theta]$, $Y_k = \Omega_k \cap Y$ and $\Sigma_k = F^{-1}(4k\theta)$.

By construction, each part (Ω_k, Y_k) contains critical points only of one type: for $k = 0$ and $n+1$ they are interior critical points, for $k = 1, \dots, n$ they are boundary unstable of index k and for $k = 1/2, \dots, n+1/2$, they are boundary stable of index $k+1/2$ and we conclude the proof by Proposition 2.35. \square

Remark 4.19. If the cobordism is a product on the boundary, i.e. $Y = M \times [0, 1]$, we can choose the initial Morse function to have no critical points on the boundary. Then all the critical points of F come in pairs, z_j^s and z_j^u with z_j^s boundary stable, z_j^u boundary unstable and there is a single trajectory of ∇F going from z_j^s to z_j^u .

The strength of Theorem 4.18 is that it is much easier to study the difference between the intersection forms on (Σ_k, M_k) and on $(\Sigma_{k\pm 1/2}, M_{k\pm 1/2})$. We refer to [BNR] for an application of this fact.

5. THE CANCELLATION OF BOUNDARY HANDLES

In this section we assume that F is a Morse function on the cobordism (Ω, Y) satisfying the Kronheimer–Mrowka–Morse–Smale regularity condition (Definition 4.5). We assume that F has precisely two critical points z and w , with $\text{ind } z = k$ and $\text{ind } w = k+1$ and that there exists a single trajectory γ of ∇F going from z to w . If z and w are both interior critical points, then [Mi2, Theorem 5.4] implies that (Ω, Y) is a product cobordism. In fact,

Milnor's proof modifies F only in a small neighbourhood of γ , which avoids the boundary Y . Hence, it does not matter, that in our case the cobordism has a boundary.

We want to extend this result to the case of boundary critical points. In some cases an analogue of the Milnor's theorem holds, in other cases we can show that it cannot hold.

5.1. Elementary cancellation theorems.

Theorem 5.1. *Let z and w be a boundary critical points of index k and $k+1$, respectively. Assume that γ is a single trajectory joining z and w . Furthermore, assume that both z and w are boundary stable, or both boundary unstable. Then (Ω, Y) is a product cobordism.*

As usual, it is enough to prove the result for boundary unstable critical points, the other case is covered if we change F to $1 - F$. Note also, cf. Section 5.2, the assumption that both critical points are boundary stable, or both boundary unstable is essential.

A careful reading of Milnor [Mi2, pages 46–66] shows that the proof there applies to this situation with only small modifications. Below we present only three steps of that proof, adjusted to our situation. We refer to [Mi2] for all the missing details.

Let ξ be the gradient vector field of F . The proof relies on the following proposition (see the Preliminary Hypothesis 5.5 in [Mi2], proved on pages 55–66).

Proposition 5.2. *There exist an open neighbourhood U of γ and a coordinate map $g: U \rightarrow \mathbb{R}_{\geq 0} \times \mathbb{R}^n$ and a gradient-like vector field ξ' agreeing with ξ away from U such that*

- $g(Y) \subset \{x_1 = 0\}$, and $g(U) \subset \{x_1 \geq 0\}$;
- $g(z) = (0, 0, \dots, 0)$;
- $g(w) = (0, 1, \dots, 0)$;
- $g_*\xi' = \eta = (x_1, v(x_2), -x_3, \dots, -x_k, x_{k+1}, \dots, x_{n+1})$, where v is a smooth function positive in $(0, 1)$, zero at 0 and 1 and negative elsewhere. Moreover $|\frac{dv}{dx_2}| = 1$ near 0 and 1.

Furthermore, U can be made arbitrary small (around γ).

Given the proposition, we argue in the same way, as in the classical case, cf. [Mi2, pages 50–55]: we improve the vector field ξ' in U so that it becomes a gradient like vector field of a function F' , which has no critical points at all. Then the cobordism is a product cobordism.

The proof of Proposition 5.2 is a natural modification of the Milnor's proof. After applying arguments as in [Mi2, pages 55–58] the proof boils down to the following result.

Proposition 5.3 (compare [Mi2, Theorem 5.6]). *Let $a + b = n$, $a \geq 1$ and $b \geq 0$ and write a point $x \in \mathbb{R}_{\geq 0} \times \mathbb{R}^{a-1} \times \mathbb{R}^b$ as (x_a, x_b) with $x_a \in \mathbb{R}_{\geq 0} \times \mathbb{R}^{a-1}$ and $x_b \in \mathbb{R}^b$. Assume that $h: (\mathbb{R}_{\geq 0} \times \mathbb{R}^{n-1}, \{0\} \times \mathbb{R}^{n-1}) \rightarrow (\mathbb{R}_{\geq 0} \times \mathbb{R}^{n-1}, \{0\} \times \mathbb{R}^{n-1})$ is an orientation preserving diffeomorphism such that $h(0) = 0$. Suppose that $h(\mathbb{R}_{\geq 0} \times \mathbb{R}^{a-1} \times \{0\})$ intersects $\{0\} \times \{0\} \times \mathbb{R}^b$ only at the origin and the intersection is transverse and the intersection index is $+1$. Then, given any neighbourhood N of $0 \in \mathbb{R}_{\geq 0} \times \mathbb{R}^{n-1}$, there exists a smooth isotopy h'_t for $t \in [0, 1]$ of diffeomorphisms from $(\mathbb{R}_{\geq 0} \times \mathbb{R}^{n-1}, \{0\} \times \mathbb{R}^{n-1})$ to itself with $h'_0 = h$ such that*

- (I) $h'_t(x) = h(x)$ away from N ;
- (II) $h'_1(x) = x$ in some small neighbourhood N_1 of 0 such that $\overline{N_1} \subset N$;
- (III) $h'_1(\mathbb{R}_{\geq 0} \times \mathbb{R}^{a-1} \times \{0\}) \cap \{0\} \times \{0\} \times \mathbb{R}^b = \{0\} \in \mathbb{R}_{\geq 0} \times \mathbb{R}^n$.

Remark 5.4. The transversality assumption from the assumption of Proposition 5.3 is equivalent to the flow of ξ being Morse–Smale.

The proof of Proposition 5.3 in Milnor's book is given on pages 59–66. We prove here only the analogue of [Mi2, Lemma 5.7]. For all other results we refer to Milnor's book.

Lemma 5.5. *Let h be as in hypothesis of Proposition 5.3. Then there exists a smooth isotopy $h_t: \mathbb{R}_{\geq 0} \times \mathbb{R}^{n-1} \rightarrow \mathbb{R}_{\geq 0} \times \mathbb{R}^{n-1}$, with h_0 the identity map and $h_t = h$, such that for each t we have $h_t(\mathbb{R}_{\geq 0} \times \mathbb{R}^{a-1}) \cap \mathbb{R}^b = 0$.*

Proof. We follow the proof of [Mi2, Lemma 5.7]. We shall construct the required isotopy in two steps. First we isotope h by $h_t(x) = \frac{1}{t}h(tx)$. Then $h_1 = h$ and h_0 is a linear map. If this is an identity, we are done. Otherwise h_0 is just a nondegenerate linear map and clearly it maps $\mathbb{R}_{\geq 0} \times \mathbb{R}^{a-1} \times \mathbb{R}^b$ diffeomorphically onto itself. It means that under the decomposition $\mathbb{R}^n = \mathbb{R} \oplus \mathbb{R}^{a-1} \oplus \mathbb{R}^b$, h_0 has the following block structure

$$h_0 = \begin{pmatrix} a_{11} & 0 & 0 \\ * & A & B \\ * & C & D \end{pmatrix},$$

where $a_{11} > 0$, and stars denote unimportant terms. As h_0 is orientation-preserving, $\det h_0 > 0$. We can apply a homotopy of linear maps which changes the first column of h_0 to $(a_{11}, 0, \dots, 0)$ and preserves all the other entries of h_0 . We do not change the determinant and the condition $h_0(\mathbb{R}_{\geq 0} \oplus \mathbb{R}^{a-1}) \cap \mathbb{R}^b = 0$ is preserved (it means that $a_{11} \det A > 0$). Let

$$h_{00} = \begin{pmatrix} A & B \\ C & D \end{pmatrix}.$$

Obviously $\det h_0 = a_{11} \det h_{00}$, so $\det h_{00} > 0$. We use the same reasoning as in the Milnor's proof to find a homotopy of h_{00} to the identity matrix, finishing the proof. \square

5.2. Non-cancellation results. The two results below have completely obvious proofs, we state them to contrast with Theorem 5.1.

Lemma 5.6. *Assume that a Morse function F on the cobordism (Ω, Y) between (Σ_0, M_0) and (Σ_1, M_1) has two critical points z and w . Suppose z is an interior critical point and w is a boundary critical point. Then (Ω, Y) is not a product cobordism.*

Proof. F restricted to Y has a single critical point, so the cobordism between M_0 and M_1 cannot be trivial. \square

Lemma 5.7. *Suppose that F has two critical points z and w . Assume that z is boundary stable and w is boundary unstable. Then (Ω, Y) is not a product.*

Proof. If it were a product, we would have $H_*(\Omega, \Sigma_0) = 0$. We shall show that this is not the case.

If $F(z) = F(w)$, then there are no trajectories between z and w , so by Proposition 4.1 we can ensure that $F(z) < F(w)$. So we can always assume that $F(z) \neq F(w)$. For simplicity assume that $F(z) < F(w)$. Let c be a regular value such that $F(z) < c < F(w)$.

By Lemma 2.17 $F^{-1}[0, c] \sim \Sigma_0 \times [0, c]$. Then $H_*(\Omega, \Sigma_0) \cong H_*(\Omega, F^{-1}[0, c])$. Now Ω arises from $F^{-1}[0, c]$ by a right half-handle addition, hence $H_*(\Omega, F^{-1}[0, c]) \cong H_*(H, B)$, where (H, B) is the corresponding right half-handle. But $H_*(H, B)$ is not trivial by Lemma 2.22 (or Lemma 2.16). \square

REFERENCES

- [AGV] V. I. Arnold, S. Gusein-Zade, A. Varchenko, *Singularities of differentiable maps. Vol. I. The classification of critical points, caustics and wave fronts* Monographs in Mathematics, **82**. Birkhäuser Boston, Inc., Boston, 1985.
- [BNR] M. Borodzik, A. Némethi, A. Ranicki, *Codimension 2 embeddings, algebraic surgery and Seifert forms*, preprint 2012, arxiv: 1211.5964.
- [Bo] R. Bott, *Morse theory indomitable*, Inst. Hautes Études Sci. Publ. Math. No. **68** (1988), 99–114.
- [GM] M. Goresky, R. McPherson, *Stratified Morse theory*, Ergebnisse der Mathematik und ihrer Grenzgebiete (3), **14**. Springer-Verlag, Berlin,

- [KM] P. Kronheimer, T. Mrowka, *Monopoles and three manifolds*, New Mathematical Monographs, **10**. Cambridge University Press, Cambridge, 2007.
- [La] F. Laudenbach, *A Morse complex on manifolds with boundary*, Geom. Dedicata **153** (2011), 47–57.
- [Mi1] J. Milnor, *Morse Theory*, Annals of Math. Studies **51**. Princeton Univ. Press, Princeton (1963)
- [Mi2] J. Milnor, *Lectures on the h-cobordism theorem*, Princeton University Press, Princeton, N.J. 1965.
- [Ni] L. Nicolaescu, *An invitation to Morse theory*, Universitext. Springer, New York, 2007.
- [Ra] A. Ranicki, *High-dimensional knot theory. Algebraic surgery in codimension 2*. Springer-Verlag, New York, 1998.
- [Sa] D. Salamon, *Morse theory, the Conley index and Floer homology*, Bull. London Math. Soc. **22** (1990), no. 2, 113–140.
- [Sm1] S. Smale, *On gradient dynamical systems*, Ann. of Math. (2) **74** (1961), 199–206.
- [Sm2] S. Smale, *On the structure of manifolds*, Amer. J. Math. **84** (1962), 387–399.
- [Wi] E. Witten, *Supersymmetry and Morse theory*, J. Diff. Geo. **17** (1982), 661–692.

INSTITUTE OF MATHEMATICS, UNIVERSITY OF WARSAW, UL. BANACHA 2, 02-097 WARSAW, POLAND
E-mail address: mcboro@mimuw.edu.pl

A. RÉNYI INSTITUTE OF MATHEMATICS, 1053 BUDAPEST, REÁLTANODA U. 13-15, HUNGARY.
E-mail address: nemethi@renyi.hu

SCHOOL OF MATHEMATICS, UNIVERSITY OF EDINBURGH, EDINBURGH EH9 3JZ, SCOTLAND UK.
E-mail address: a.ranicki@ed.ac.uk