Annals of Mathematics

Nondegenerate Critical Manifolds Author(s): Raoul Bott Source: *The Annals of Mathematics*, Second Series, Vol. 60, No. 2 (Sep., 1954), pp. 248-261 Published by: Annals of Mathematics Stable URL: <u>http://www.jstor.org/stable/1969631</u> Accessed: 08/08/2009 12:00

Your use of the JSTOR archive indicates your acceptance of JSTOR's Terms and Conditions of Use, available at http://www.jstor.org/page/info/about/policies/terms.jsp. JSTOR's Terms and Conditions of Use provides, in part, that unless you have obtained prior permission, you may not download an entire issue of a journal or multiple copies of articles, and you may use content in the JSTOR archive only for your personal, non-commercial use.

Please contact the publisher regarding any further use of this work. Publisher contact information may be obtained at http://www.jstor.org/action/showPublisher?publisherCode=annals.

Each copy of any part of a JSTOR transmission must contain the same copyright notice that appears on the screen or printed page of such transmission.

JSTOR is a not-for-profit organization founded in 1995 to build trusted digital archives for scholarship. We work with the scholarly community to preserve their work and the materials they rely upon, and to build a common research platform that promotes the discovery and use of these resources. For more information about JSTOR, please contact support@jstor.org.



Annals of Mathematics is collaborating with JSTOR to digitize, preserve and extend access to The Annals of Mathematics.

NONDEGENERATE CRITICAL MANIFOLDS

By RAOUL BOTT

(Received February 4, 1953)

Introduction

Let J be a smooth¹ function defined on an open set U of Euclidean n space E_n . A point $x \in U$ is called a critical point of J if the gradient of J vanishes at x.² The nullity and index of a critical point x of J are defined as the dimension of the null space and the number of negative characteristic roots of the Hessian³ of f evaluated at x, respectively.

It is a basic result of the Morse theory [1] that if x is a critical point of nullity 0, then for all sufficiently small neighborhoods V of x (in U) the following isomorphisms hold:

(I)
$$H_k (J_x \cap V, J_x \cap V) \approx H_{k-\lambda_x} (x)$$

In this formula $J_x[J_x^-]$ denotes the subset of U on which $J \leq J(x) [J < J(x)]$, while λ_x is the index of x. H_k ($J_x \cap V$, $J_x^- \cap V$) and $H_k(x)$ ($k = 0, 1, \cdots$) refer to the k^{th} singular homology group [9] of the pair ($J_x \cap V$, $J_x^- \cap V$) at the left and of the point x at the right. The coefficients are immaterial; however, we shall assume throughout the paper that all homology groups considered are computed mod 2.

The purpose of this paper is the proof and application of the following extension of (I).

DEFINITION. Let $M \subset U$ be a compact manifold smoothly imbedded in U such that (a) all points of M are critical points of a smooth function J, defined on U, and (b) the nullity of all $x \in M$ equals the dimension of M; then M will be called a nondegenerate critical manifold of J on U.

THEOREM I. If M is a nondegenerate critical manifold of J on U, then for all sufficiently small neighborhoods V of M

(I*) $H_k (J_M \cap V, J_M \cap V) \approx H_{k-\lambda_M}(M)$

where $J_M[J_M]$ refers to the subset of U on which $J \leq J(x) [J < J(x)] x \in M$, and λ_M denotes the index of some $x \in M$.⁴

As an application we compute the "sensed circular connectivities" of the n-sphere. Our result is the following one: (see §2 for exact definitions) if we de-

¹ The term "smooth" will mean "at least three times continously differentiable" throughout the paper.

² i.e. $\partial J/\partial x_i(x) = 0$ $(i = 1 \cdots n)$.

³ The Hessian of J is the $n \times n$ matrix of second partial derivatives of J with respect to x_i $(i = 1 \cdots n)$.

 $^{{}^{4}\}lambda_{M}$ and J_{M} , $[J_{M}]$ are independent of $x \in M$, as our conditions guarantee that J and the nullity are constants on M.

note by $\mathfrak{M}(S^n; z) = \sum_{k=0}^{\infty} p_k z^k$ the formal power series with p_k equal to the k^{th} sensed circular connectivity of S^n we show that

(II)
$$\mathfrak{M}(S^{n}; z) = P(G_{n-1}^{2}; z) z^{n-1} (1 - z^{2(n-1)})^{-1}$$

where $P(G_{n-1}^2; z)$ is the Poincaré polynomial of the mod 2 Betti numbers of the Grassmann manifold of oriented 2-planes in Euclidean n + 1 space. For example, $\mathfrak{M}(S^2; z) = (1 + z^2) (z + z^3 + z^5 \cdots).$

Our method also yields a formula for the circular connectivities \bar{p}_k of the *n*-sphere, namely:

(III)
$$\overline{\mathfrak{M}}(S^{n}; z) = P(\bar{G}_{n-1}^{2}; z) \, z^{n-1} (1 - z^{2(n-1)})^{-1}$$

where now $\overline{\mathfrak{M}}(S^n; z) = \sum_{0}^{\infty} \overline{p}_k z^k$, while \overline{G}_{n-1}^2 stands for the Grassmann manifold of unoriented 2-planes in E_{n+1} .

In [1] Morse characterised \bar{p}_k as the number of solutions of certain diophantine equations ([1]; p. 349). It appears therefore that in his classic derivation of $\overline{\mathfrak{M}}(S^n; z)$ Morse obtained the Betti numbers of \bar{G}_{n-1}^2 as a byproduct.

The factorization of \mathfrak{M} and $\overline{\mathfrak{M}}$ as given by II and III is not without a certain intuitive appeal. Recall that Morse introduced the numbers \bar{p}_k for the purpose of counting closed geodesics on manifolds. On the other hand the *n*-sphere in its usual representation in E_{n+1} admits a continuum of closed geodesics, which in a rather obvious topology is homeomorphic to G_{n-1}^2 if the geodesics are taken as oriented, and to \bar{G}_{n-1}^2 if they are taken as unoriented. Furthermore, a given geodesic can be traversed an integral number of times; this would then account for the factor $z^{n-1}(1 - z^{2(n-1)})^{-1}$.

The sensed circular connectivities p_k introduced here can be shown to have the same relation to the "nonreversible problem" that the \bar{p}_k have to the "reversible problem" considered by Morse. This relationship and some of its applications will be reported at another time.

1. Proof of (I^*)

We recall from [1] that the proof of (I) is constructed as follows: Consider the linear manifold E_x of vectors x + y passing through the critical point x, spanned by those vectors y which are linear combinations of the characteristic vectors associated with the negative characteristic roots of the Hessian of J at x. The dimensionality of this manifold is λ_x . The formula (I) is established by the construction of a homotopy retraction of the pair $(J_x, J_x) \cap V_{\varepsilon}$ onto the pair $(E_x, E_x - x) \cap V$ valid for all sufficiently small spherical neighborhoods V_{ε} of x (see [2] p. 34).

To prove (I*) we first surround M by an ε neighborhood V_{ε} so small that the mapping $p: V_{\varepsilon} \to M$ defined by the orthogonal projection of points of V_{ε} onto M is well defined and constitutes a fiber decomposition of V_{ε} , with base space M, and fiber the orthogonal cells F'_x of dimension $n - \dim M$. Thus \bar{V}_{ε} can be considered as the mapping cylinder of the normal sphere bundle of M. For each $x \in M$ the manifold E_x is well defined and will, as a consequence of condition b,

be completely orthogonal to M at x. Let $A_{\varepsilon} \subset V_{\varepsilon}$ be the subbundle of V_{ε} consisting of those $x \in V_{\varepsilon}$ for which $x \in E_{p(x)}$. A_{ε} is thus a bundle whose fibers F_x are λ_M -cells. M is imbedded in A_{ε} . We set A^- equal to $A_{\varepsilon} - M$. Now compare the pair $(A_{\varepsilon}, A_{\varepsilon}^-)$ with $(J_M, J_M^-) \cap V_{\varepsilon}$. Clearly $(A_{\varepsilon}, A_{\varepsilon}^-) \subset (J_M, J_M^-) \cap V_{\varepsilon}$ for sufficiently small ε . However, the following much stronger proposition holds:

PROPOSITION. For $0 < \varepsilon$ small enough $(A_{\varepsilon}, A_{\varepsilon})$ is a fiber preserving deformation retract of $(J_M, J_M) \cap V_{\varepsilon}$.

In particular, therefore

$$H_m\{(J_M, J_M^-) \cap V_{\varepsilon}\} \approx H_m(A_{\varepsilon}, A_{\varepsilon}^-)$$

for small enough ε .

PROOF. The proof proceeds in entire analogy to that of (I) given in [2. p. 34]. We set V equal to the "abstract" normal bundle [8] of $M \subset U. E_x$, $x \in M$ is a well defined subspace of the fiber of V at x. Let D_x be the complementary subspace to E_x and let E, D denote the subbundle consisting of $\bigcup_{x \in M} E_x$, $\bigcup_{x \in M} D_x$ respectively. We set H^+ and H^- equal to the projections defined on V which map the normal space at x into D_x and E_x respectively. H⁺ and H⁻ are continuous fiber mappings of V onto D and E respectively. Consider now the following deformation which establishes E as a fiber and norm preserving deformation retract of V - D. This deformation will be parametrized by $\pi/4 \leq \theta \leq \pi/2$, and assigns to each $y \in V$ the element $y_{\theta} = \frac{\cos \theta H^+ y + \sin \theta H^- y}{|\cos \theta H^+ y + \sin \theta H^- y|} |y|$ for $y \neq 0$, and $y_{\theta} = y$ for y = 0. (We use a Euclidean metric in the fiber and |y| denotes the length of a vector y.)

On each fiber this deformation reduces to the one used by Seifert and Threlfall in [2].

Now for small enough ε we can identify V and V_{ε} so as to take radial lines into radial lines. Then the restriction of J to a given fiber F will have a critical point of index 0 at the intersection of F and M. Hence the argument of [2] applies and for sufficiently small ε the proposition to be proved will be true on each individual fiber. By continuity and compactness it follows that the proposition as stated is correct.

Hence (I*) will be demonstrated if we prove that $H_k(A_{\varepsilon}, A_{\varepsilon}) \approx H_{k-\lambda_M}(M)$. As we are using mod 2 coefficients throughout, it is sufficient to prove the equality of the ranks of the groups in question.

Consider now the exact sequence of the pair $(A_{\varepsilon}, A_{\varepsilon}^{-})$:

(1.1)
$$\rightarrow H_n(A_{\varepsilon}) \rightarrow H_n(A_{\varepsilon}) \rightarrow H_n(A_{\varepsilon}, A_{\varepsilon}) \rightarrow H_{n-1}(A_{\varepsilon}) \rightarrow$$

 A_{ε}^{-} is a bundle with fiber deformable into a $\lambda_{M} - 1$ sphere. Accordingly we have the exact Gysin, Thom, Chern – Spanier [3] [4] [5] sequence of A_{ε}^{-} as follows:

(1.2)
$$\rightarrow H_n(A_{\varepsilon}) \rightarrow H_n(M) \rightarrow H_{n-\lambda_M}(M) \rightarrow H_{n-1}(A_{\varepsilon}) \rightarrow$$
.

250

Comparing the two sequences it is easily seen that $H_{n-\lambda_M}(M)$ and $H_n(A_{\varepsilon}, A_{\varepsilon}^-)$ are equivalent group extensions and therefore have the same rank. (We merely have to establish the commutativity of the square

$$\begin{array}{ccc} H_n(A_{\varepsilon}) & \to & H_n(A_{\varepsilon}) \\ \uparrow & & \uparrow \\ H_n(A_{\varepsilon}^-) & \to & H_n(M) \end{array}$$

which of course is well known [4]. We might add that in Thom's abstract [4] of the proof of (1.1) the isomorphism $H_{m-\lambda_M}(M) \stackrel{\phi_*}{\leftarrow} H_m(A_{\varepsilon}, A_{\varepsilon})$ is stated as the basic lemma and (1.2) deduced from it. The Thom result asserts the isomorphism in question to exist for a suitable local system of coefficients on M.)

REMARK. The local nature of the whole problem makes it clear that the theorem is equally true if we replace the imbedding space U by an open subset of an arbitrary Riemann manifold.

2. The circular connectivities

The m^{th} circular connectivity \bar{p}_m of a Riemann manifold R is defined by Morse [1] as the rank of the limit group $\overline{\mathfrak{M}}_m(R)$ of the $m^{\text{th}} \mod 2$ homology groups of a directed system of pairs [9] { $(\bar{\pi}_k, \bar{\Delta}_k); h$ }. $\bar{\pi}_k(R)$ (written $\bar{\pi}_k$ when there is no danger of confusion!) is defined as follows: In $R^k = R \times \cdots \times R$ (k factors) consider the subset of points $(x) = \{x_1 \cdots x_k\}$ such that the distance between consecutive points in the array $x_1, x_2, \cdots x_n, x_1$ on R is less than the elementary length ρ of geodesics on R. (See [1] p. 196.) We denote this subset in R^k by Π_k . A point (x) of Π_k uniquely determines a closed geodesic polygon on R, namely the polygon obtained by joining consecutive points of the above array by elementary geodesics. If in Π_k we identify those points which differeither by a cyclic permutation of the factors of R^k , or by a reversal of the order of the factors of R^k , we obtain the space $\bar{\pi}_k$ (i.e. $(x_1, \cdots, x_k) \sim (x_k, x_1 \cdots x_{k-1}) \sim (x_k, x_{k-1}, \cdots x_1)$ in $\bar{\pi}_k$). The points of $\bar{\pi}_k$ therefore correspond to closed unsensed k edged geodesic polygons on R, none of whose vertices are distinguished.

By introducing s equally spaced vertices on each of the sides of a polygon $(x) \epsilon \bar{\pi}_k$ a new polygon with sk vertices is obtained. This subdivision operation gives rise to maps h_s of $\bar{\pi}_k$ into $\bar{\pi}_{sk}$ which make the family $\bar{\pi}_k$ into the directed system $\{\bar{\pi}_k; h\}$. If we now set $\bar{\Delta}_k$ equal to the diagonal in $\bar{\pi}_k$, (that is, the set of points of the form $(x) = (x, x, \dots, x)$) we obtain the desired family of pairs $\{(\bar{\pi}_k, \bar{\Delta}_k); h\}$.

The length of a polygon is a continuous function J defined on all $\bar{\pi}_k$. We clearly have $J(x) = J(h_s(x)), x \in \bar{\pi}_k$ (all s); hence if $\bar{\pi}_k^c[\bar{\pi}_k^{c-}]$ denotes the subspace of $\bar{\pi}_k(R)$ on which $J \leq c [J < c]; c \geq 0, h$ restricted to $\bar{\pi}_k^c[\bar{\pi}_k^{c-}]$ makes the family $\{\bar{\pi}_k^c, \bar{\Delta}_k; h\}$ [$\{\bar{\pi}_k^{c-}, \bar{\Delta}_k; h\}$] a directed system. The group $\overline{\mathfrak{M}}(J_c; R)$ [$\overline{\mathfrak{M}}(J_{c-}; R)$] is defined as the limit homology group of $\{(\bar{\pi}_k^c, \bar{\Delta}_k); h\}$ [$\{(\bar{\pi}_k^{c-}, \bar{\Delta}_k); h\}$].

By the symbol $\overline{\mathfrak{M}}(J_c, J_c, ; R) c > c'$ we will mean the limit relative homology group of the family of pairs $\{\overline{\pi}_k^c, \overline{\pi}_k^{c'}; h\}$.

RAOUL BOTT

REMARK. The definition of $\overline{\mathfrak{M}}(R)$ seems to be dependent on Riemann structure of R. However, Morse showed that $\overline{\mathfrak{M}}(R)$ is a topological invariant of R. [See [1] p. 297.] His proof can easily be modified to yield also the topological invariance of the group $\mathfrak{M}(R)$ defined in the next section:

3. The sensed circular connectivities

In the previous construction $\bar{\pi}_k(R)$ was obtained from $\Pi_k(R)$ be identifying on $\Pi_k(R)$ points which were images under the group of cyclic permutations, or reversal of order. Let $\pi_k(R)$ be the space obtained from $\Pi_k(R)$ by only identifying those points which are cyclically equivalent. Then if we replace $\bar{\pi}_k$ by π_k in the construction of §2, we obtain the "sensed" counterparts of $\overline{\mathfrak{M}}(R)$ etc. which we denote by $\mathfrak{M}(R)$, etc. We will adhere to the notation that the barred letters stand for the equivalent in the "unoriented" theory of the unbarred letters in the "oriented" theory.

4. On the computation of $\mathfrak{M}(S^n; z)$

Let S^n be represented by the subset $\sum_{i=1}^{n+1} x_i^2 = 1/(2\pi)^2$ in Euclidean n + 1 space. The metric induced on S^n by the Euclidean metric makes S^n into an analytic Riemann manifold. It is well known that then the closed geodesics on S^n "are" the intersection of S^n with two-planes $E^2 \subset E^{n+1}$. The length of these geodesics is t, t being the positive integral value of the number of times the point set $E^2 \subset S^{n+1}$ is traversed. For ρ we take the number $\frac{1}{3}$.

The computation of $\mathfrak{M}(S^n; z)$ is broken up into two steps. First we compute the group $E_1 = \sum_{i=1}^{\infty} \mathfrak{M}(J_t, J_{t-1})$ (direct sum). E_1 will turn out to be finitely generated in each dimension.

Next let $E_{\infty} = \sum_{1}^{\infty} D_t / D_{t-1}$ (direct sum) where D_t is the image of $\mathfrak{M}(J_t)$ in $\mathfrak{M}(R)$ under the inclusion map induced by the inclusions $(\pi_k^t, \Delta) \to (\pi_k, \Delta)$. As $\pi_k^t = \Delta$ and $\pi_k^t = \pi_k$ for sufficiently large t, we have

$$0 = D_0 \subset D_1 \subset \cdots \subset \mathfrak{M}(R) = \bigcup_0^\infty D_t.$$

The second step of the computation consists of showing that E_1 and E_{∞} are isomorphic in a dimension preserving fashion. It follows then that $\mathfrak{M}_n(R)$ is also finitely generated in each dimension and as we are working over a field of coefficients E_{∞} is (unnaturally) isomorphic to $\mathfrak{M}(R)$ in a dimension preserving fashion. The procedure of going from E_1 to E_{∞} is of course the basic method of the classical work of Morse as well as of the modern spectral theory of Leray. E_1 is obtained by local considerations while the step from $E_1 \to E_{\infty}$ usually involves arguments "in the large". In the case under consideration however, it turns out very fortuitously that $E_1 \approx E_{\infty}$ for purely "dimensional" reasons.

5. Computation of E_1

Let $\tilde{\mu}_k [\mu_k]$ denote the subset of $\Pi_k [\pi_k]$, each point of which represents a polygon whose k edges are equal in length. A point $z \in \tilde{\mu}_k [\mu_k]$ is called critical if it represents a closed geodesic. Clearly in our case J takes on only integral values on critical points of $\tilde{\mu}_k [\mu_k]$. We set $\tilde{\sigma}_{t,k} [\sigma_{t,k}]$ equal to the totality of critical points of length t in $\tilde{\mu}_k [\mu_k]$. As $\rho = \frac{1}{3}$, $\tilde{\sigma}_{t,k} [\sigma_{t,k}]$ will be nonvacuous for t < 3k, and we assume in the following that this condition is always satisfied.

It is shown by Morse ([1], Theorem 5.1, p. 261) that for sufficiently small neighborhoods $\tilde{V}_{t,k}$ of $\tilde{\sigma}_{t,k}$ in $\tilde{\mu}_k$, $\tilde{V}_{t,k}$ is an analytic manifold (in the subset topology induced by the imbedding of $\tilde{\mu}_k \subset \Pi_k$). As points of $\tilde{V}_{t,k}$ are well away from the diagonal in Π_k , the projection $\tilde{\pi}_k \to \pi_k$ induces a manifold structure on the image $V_{t,k}$ of $\tilde{V}_{t,k}$.

The following basic proposition due to Morse establishes the local character of E_1 . (We have phrased it to fit into our context.)

THEOREM A. (Morse) Let $V_{t,k}$ be any family of sufficiently small neighborhoods of $\sigma_{t,k} \subset \mu_k$ (t < 3k), such that $h_s: V_{t,k} \subset V_{t,sk}$ and let $V_{t,k}^*[V_{t,k}]$ be the subset of $V_{t,k}$ on which $J \leq t$, [J < t]. Then

$$\mathfrak{M}_m(J_t, J_{t-1}) \approx \lim_{(k)} \{H_m(V_{t,k}^*, V_{t,k}); h\}.$$

REMARK. Theorem A can be deduced roughly in the following fashion: One first shows that $\mathfrak{M}_m(J_t, J_{t-1}) \approx \mathfrak{M}_m(J_t, J_{t-1})$ using the fact that there are no critical points of J between t and t - 1. Next one shows that for any system of neighborhoods $\dot{V}_{t,k}$ of $\sigma_{t,k}$ on π_k with $h_s \dot{V}_{t,k} \subset \dot{V}_{t,ks}$,

$$\mathfrak{M}_m(J_t, J_{t-}) \approx \lim_k \{H_m(\dot{V}_{t,k}^*, \dot{V}_{t,k}^-); h\}$$

(here the * and $\bar{}$ superscript have the same meaning as in Theorem A). Finally for a sufficiently small family $\dot{V}_{t,k}$, the pair $(\dot{V}_{t,k}^*, \dot{V}_{t,k})$ can be J deformed into $(\dot{V}_{t,k}^*, \dot{V}_{t,k}) \cap u_k = (V_{t,k}^*, V_{t,k}).$

All the deformations needed to carry this program out are given by Morse in [1]. (p. 250–270.) I am planning to give a more detailed expository account of this phase of the Morse theory in the near future.

With the aid of Theorem A the computation of E_1 is reduced to the study of the groups $H_m(V_{t,k}^*, V_{t,k}^-)$ and the homomorphisms

$$h_{s^*}: H_m(V_{t,k}^*, V_{t,k}^-) \to H_m(V_{t,ks}^*, V_{t,ks}^-)$$

induced by the restriction of h_s to $V_{t,k}$.

It is clear that if we choose $V_{t,k}$ small enough $\sigma_{t,k}$ will be the only critical set of the function J on $V_{t,k}$. We propose to show that $\sigma_{t,k}$ is a nondegenerate critical manifold of J on $V_{t,k}$ in the sense of §1.

If we assign to each oriented closed geodesic $x \in \sigma_{t,k}$ the oriented two plane in which it lies, we obtain a representation of $\sigma_{t,k}$ as a 1-sphere bundle over G_{n-1}^2 the Grassmann manifold of oriented 2 planes in E^{n+1} . The dimension of $\sigma_{t,k}$ is therefore 2n - 1.

Now let x be a point of $\sigma_{t,k}$. As a critical point of J on $V_{t,k}$ x has a well defined index and nullity.

LEMMA. The nullity and index of any point $x \in \sigma_{t,k}$ (t < 3k) are respectively given by

$$N(x) = 2n - 1$$

$$\lambda(x) = (n - 1) (2t - 1).$$

It follows that $\sigma_{t,k}$ is a nondegenerate critical manifold of $V_{t,k}$ and hence by (I*)

$$H_m(V_{t,k}^*, V_{t,k}^-) \approx H_{m-(2t-1)(n-1)}(\sigma_{t,k})$$

for sufficiently small neighborhood $V_{t,k}$.

PROOF. In the neighborhood of $\sigma_{t,k}$, π_k is an analytic manifold on which J is well defined. μ_k is smoothly imbedded in π_k . Every point x of $\sigma_{t,k}$ is a critical point of J on π_k as well as of J on μ_k . It follows that N(x) and $\lambda(x)$ can be interpreted as the nullity and number of negative characteristic roots of the Hessian of J as a function on π_k restricted to the tangent space of μ_k in π_k at x.

At each vertex p_i $(i = 1 \cdots k)$ of x we introduce a local analytic coordinate system $v_1^i \cdots, v_n^i$ covering a neighborhood U_i of p_i in such a manner that the point p on x, whose signed distance from p_i is equal to d (recall that x is oriented) has coordinates $v_1^i = d$; $v_2^i = v_3^i \cdots v_n^i = 0$, for sufficiently small |d|. To emphasize this choice of coordinates notationally we shall write d^i for v_1^i . Clearly a neighborhood U of x in π_k is given by those polygons whose i^{th} vertex lies on U_i $(i = 1 \cdots k)$. (As we are working locally and away from the diagonal in π_k it is clearly "all right" to number our vertices, starting at any vertex we please and then adhering to the choice made.) Consider now the expansion of J in terms of the local coordinates introduced. As J is independent of d^i when $v_2^i =$ $\cdots = v_n^i = 0$, all partials of J with respect to the d^i 's are 0. Also as x is a closed geodesic for any choice of the d^i when $v_2^i = \cdots = v_n^i = 0$, $\partial J/\partial v_j^i = \partial^2 J/\partial d^i \partial v_j^i =$ 0 for $(i, i' = 1 \cdots k, j = 2 \cdots n)$. Hence J has an expansion which starts off with $J = J(x) + Q_x$, where Q_x is a quadratic form in the variables $(v_2^i, \cdots v_n^i)$ $(i = 1 \cdots k)$ only.

This fact can also be expressed in the following fashion. Let $W \subset U$ be the submanifold of U given by the equations $d^i = 0$ $(i = 1 \cdots k)$. W is called by Morse a proper section of x. Let $f: U \to W$ be the projection of U onto W defined by: $f(v_1^1, v_2^1 \cdots; v_2^2, v_2^2 \cdots; \cdots; v_1^k, v_2^k \cdots v_n^k) = (0, v_2^1, \cdots; 0, v_2^2 \cdots; \cdots; 0, v_2^k \cdots)$. The above result is then equivalent to the assertion that J(f(x)) and J(x) coincide on U up to terms of the second order.

The index and nullity respectively of a geodesic x with respect to a proper section W is defined by Morse (see [1], pg. 288) as

$$N_W(x) =$$
nullity of Q_x

(as a function of $v_2^i \cdots v_n^i$ only!) and

$$\lambda_w(x) = \text{index of } Q_x$$
.

Now the numbers $N_W(x)$; $\lambda_W(x)$; N(x), $\lambda(x)$ are related in the following fashion

(5.1)
$$N(x) = N_W(x) + 1$$

(5.2)
$$\lambda(x) = \lambda_W(x).$$

This is seen as follows. To compute N(x) we have to consider the nullity of Q_x in the tangent space of $\mu_k \subset \pi_k$, at x. However in the local coordinate systems

which we have chosen, the tangent space of u_k at x is clearly characterized by the equations $d^1 = d^2 = \cdots = d^k$. As the d^i 's do not occur at all in Q_x the formulae (5.1) and (5.2) become self evident.

A fundamental result of the Morse theory is an invariance theorem (see [1], p. 289) which in our context reads as follows.

THEOREM 6. (Morse) Let $x \in \sigma_{t,k}$, and let $x' \in \sigma_{t,ks}$ be the image of x under the subdivision map h_s (for s = 1, h_1 is the identity map). Let W be a proper section of x and W' a proper section of x'. Then

(5.3)
$$N_{W}(x) = N_{W}(x') \text{ and } \lambda_{W}(x) = \lambda_{W'}(x').$$

It follows that N(x) and $\lambda(x)$ are independent of the particular proper section used and invariant under subdivision; i.e. independent of k.

We can therefore choose k = 4t for the explicit computations. This choice of k makes the length between consecutive points of x equal to $\frac{1}{4}$. We now compute $N_{W}(x)$ and $\lambda_{W}(x)$ for a special proper section W, which in effect has already been constructed by Seifert and Threlfall in another connection (see [2], p. 61).

We recall their construction briefly for the case of S^2 . Let $x \\ \epsilon \\ \sigma_{t,4t}$ be represented by the polygon which starting at p_0 (the point where the x_3 axis intersects S^2), circles the sphere t times in the (2, 3) plane. The second vertex of x will then lie on the first intersection of x with the (12) plane, p_3 will be the antipode of p_1 , p_4 the antipode of p_2 etc. To construct the proper section W we take open intervals about the vertices of x on geodesics cutting these vertices at right angles to x, and use as local coordinates $\{v_2^i\}$ on these, the geodesic length. The well known cosine law of spherical trigonometry then easily yields for Q_x the quadratic form

$$Q_x = -(v_2^1 v_2^2 + v_2^2 v_2^3 + \cdots + v_2^{4t-1} v_2^{4t} + v_2^{4t} v_2^1)$$

whose nullity is seen to be 2 and whose index is (2t - 1). Using the analogous construction for S^n , Q_x is seen to take the form

$$Q_x = -(v_2^1 v_2^2 + v_2^2 v_3^2 + \dots + v_2^{4t-1} v_2^{4t} + v_2^{4t} v_2^1) -(v_3^1 v_3^2 + \dots + v_3^{4t-1} v_3^{4t} + v_3^{4t} v_3^1) -(v_n^1 v_n^2 + \dots + v_n^{4t-1} v_n^{4t} + v_n^{4t} v_n^1)$$

whence its index is (2t - 1) (n - 1) and its nullity 2(n - 1). Hence

$$N(x) = 2(n-1) + 1 = 2n - 1$$

and

$$\lambda(x) = (2t - 1) (n - 1).$$

This completes the proof of the lemma.

The next step in the computation of E_1 is the investigation of the subdivision homomorphism

$$h_{s*}: H(V_{t,k}^*, V_{t,k}^-) \to H(V_{t,ks}^*, V_{t,ks}^-).$$

The Thom construction of the isomorphism ϕ_* of (I*) makes it evident that ϕ_* can be so chosen that commutativity holds in the following diagram:

Hence it is sufficient to study the limit group of the directed system $\{H_m(\sigma_{t,k}), (h \mid \sigma_{t,k})_*\}$ (t fixed !).

It is convenient at this point to transfer the computation to cohomology in order to exploit the multiplicative properties of the cohomology ring of $\sigma_{t,k}$. As the underlying coefficient system is the compact field of integers mod 2 the direct limit homology groups and inverse limit cohomology groups will be isomorphic, assuming the limit groups to be finite.

We recall that $\sigma_{t,k}$ is a 1-sphere bundle over the simply connected manifold G_{n-1}^2 . Let \hat{G}_{n-1}^2 be the Stiefel manifold of two-frames in E^{n+1} (i.e. the tangent bundle of S^n). \hat{G}_{n-1}^2 is also a 1-sphere bundle over G_{n-1}^2 . We further have a bundle map of

 $\hat{G}_{n-1}^2 \to \sigma_{t,k}$

which is constructed by assigning to each tangent vector v on S^n , the $x \in \sigma_{t,k}$ which has one of its vertices at the base point of v and which lies in the two-plane determined by v and the origin of E^{n+1} . This map is fiber preserving and induces the identity map of the base spaces. (It is in fact a covering projection.)

If p^1, \dots, p^k are the vertices of $x \in \sigma_{t,k}$ then only $a_{t,k}$ of the points p^i will be distinct, where $a_{t,k}$ is defined as the numerator of k/t written in its lowest terms. Hence in the mapping $\hat{G}_{n-1}^2 \to \sigma_{t,k}$ the fiber S^1 of \hat{G}_{n-1}^2 is mapped onto the fiber S^1 of $\sigma_{t,k}$ with degree $a_{t,k}$.

Consider now the Leray spectral sequence of the bundle $\sigma_{t,k} \to G_{n-1}^2$. (See Serre [6] in particular p. 470.) Using cohomology with mod 2 coefficients and Serre's notation and terminology throughout, we have the canonical ring isomorphism

$$E_2^{p,q}(\sigma_{t,k}) \approx H^p(G_{n-1}^2) \otimes H^q(S^1).$$

We define $c_{t,k} \in H^2(G_{n-1}^2)$ as the class d_2a where a generates $H^1(S^1)$. We let $c \in H^2(G_{n-1}^2)$ be d_2a in $E_2(\widehat{G}_{n-1}^2)$. The mapping

$$\sigma_{t,k} \to \widehat{G}_{n-1}^2$$

induces a homomorphism of $E_2^{p,q}(G_{n-1}^2) \to E_2^{p,q}(\sigma_{t,k})$ which is the identity on $H(G_{n-1}^2)$, has degree $a_{t,k}$ on $H(S^1)$, and commutes with d_2 . (See [10], p. 130, also [11].) It follows that $c_{t,k} = a_{t,k}c$. In particular if $a_{t,k}$ is even $c_{t,k} = 0$. Hence in that case d_2 is trivial in all of $E_2(\sigma_{t,k})$ and, as the higher differentials are trivial

for dimensional reasons, $E_{\infty}(\sigma_{t,k}) \approx E_2(\sigma_{t,k})$. We have therefore established the following proposition:

PROPOSITION. If $a_{t,k}$ is even

$$E^{p,q}_{\infty}(\sigma_{t,k}) \approx H^p(G^2_{n-1}) \otimes H^q(S^1) \qquad (ring \ isomorphism).$$

REMARK. Recall that the additive structures of $E_{\infty}(\sigma_{t,k})$ and of $H(\sigma_{t,k})$ are (unnaturally) isomorphic as we are dealing with a field of coefficients.

The evaluation of the inverse limit $\lim_{k \in I} H(\sigma_{t,k})$ now proceeds in two steps. First we replace the directed system I (the integers ordered by divisibility) over which k ranges, by I_t , the subset of I for which $a_{t,k}$ is even. For any fixed $t I_t$ is cofinal with I.

Next we compute the inverse limit $\lim_{I_t} E_{\infty}(\sigma_{t,k})$, and show that the additive structure of this group and $\lim_{I_t} H(\sigma_{t,k})$ are, again unnaturally, isomorphic. (The groups $E_{\infty}(\sigma_{t,k}); k \subset I_t$ become an inverse system by means of the homomorphisms $E_{\infty}^{p,q}(\sigma_{t,ks}) \to E_{\infty}^{p,q}(\sigma_{t,k})$ induced by the fiber mappings $h_s:\sigma_{t,k} \to \sigma_{t,ks}$.) Thanks to the proposition above this inverse system can be described in detail: Each group of the family is isomorphic to $H(G_2^{n-1}) \otimes H(S^1)$ and h_s^* is the identity on $H(G_2^{n-1}) \otimes H^0(S^1)$, while $h_s^*(e \otimes a) = s(e \otimes a)$. (Here *e* generates $H^0(G_2^{n-1})$ and *a* generates $H^1(S^1)$ as above.) As we are working mod 2 it follows that no element of the inverse limit can have a component in any group which is of the form $e \otimes a$. As the homomorphisms involved are ring homomorphisms, neither can a component of the type $u \otimes a$ ($u \neq 0 \in H(G_2^{n-1})$) occur. It follows that only components contained in $H(G_2^{n-1}) \otimes H^0(S')$ occur, whence $\lim_{I} E_{\infty}^{p,0}(\sigma_{t,k}) \approx H^p(G_2^{n-1})$ and $\lim_{I} E_{\infty}^{p,q}(\sigma_{t,k}) = 0$ for q > 0. It remains to prove the additive isomorphism of the two groups. $\lim_{I} E_{\infty}(\sigma_{t,k})$ and $\lim_{I} H(\sigma_{t,k})$.

Recall that the fibering of $\sigma_{t,k}$ over G_2^{n-1} induces a filtering of $H(\sigma_{t,k})$ whose associated graded group $\mathfrak{G}H(\sigma_{t,k})$ is canonically isomorphic to $E_{\infty}(\sigma_{t,k})$. The homomorphisms h_s preserve the filtration and thereby induce the maps on E_{∞} which we have been discussing above. We can introduce a filtration in $\lim_I H(\sigma_{t,k})$ by taking the filtration of a limit element to be $\leq p$ if the filtration of every one of its components is $\leq p$. We denote the associated graded group $\mathfrak{G} \lim_I H(\sigma_{t,k})$. As all groups in question are finite dimensional vector spaces over a field, Theorem 5.7, p. 226 of [9]⁵ yields a canonical isomorphism of $\lim_k E_{\infty}(\sigma_{t,k}) = \lim_k \mathfrak{G}H(\sigma_{t,k})$ onto $\mathfrak{G} \lim_I H(\sigma_{t,k})$. It follows that the additive structure of $H(\sigma_{t,k})$ is (unnaturally) isomorphic to $\lim_I E_{\infty}(\sigma_{t,k})$. Assembling these results with Lemma I and reverting to homology we obtain the evaluation:

$$\lim_{I} H_{m}(V_{t,k}^{*}, V_{t,k}^{-}) \approx H_{m-(2t-1)(n-1)}(G_{n-1}^{2}).^{6}$$

⁵ This theorem states that inverse limit operation preserves exactness in the category of finite dimensional vector spaces over a field.

⁶ I am indebted to the referee for pointing out an error in the original derivation of this formula, as well as suggesting a remedy—which has been essentially followed.

Hence by the Morse result (Theorem A)

$$\mathfrak{M}_{m}(J_{t}, J_{t-1}) \approx H_{m-(2t-1)(n-1)}(G_{n-1}^{2}).$$

6. The step from E_1 to E_{∞}

The group E_1 is a direct sum of the groups $\mathfrak{M}_m(J_t, J_{t-1}); t \ge 1$. We set

$$E_1^{p,q} = \mathfrak{M}_{p+q}(J_p, J_{p-1}) \qquad p = 1, 2, \cdots$$

$$E_1^{p,q} = 0 \text{ for } p \leq 0.$$

Thus E_1 becomes the direct sum of the groups $E_1^{p,q}$ over all p, q. From the evaluation of $\mathfrak{M}_m(J_t, J_{t-1})$ and the fact that G_{n-1}^2 has dimension (2n - 2) it is seen that $E_1^{p,q}$ is trivial except possibly when

(6.1)
$$0 \le p + q - (2p - 1)(n - 1) \le 2(n - 1)$$

or equivalently for fixed p > 0 when q lies in the range:

(6.2)
$$(2n-3)p - (n-1) \leq q \leq (2n-3)p + (n-1).$$

Let Q_p be the interval of integers characterized by the above inequality. We observe that $Q_p \cap Q_{p-1}$ consists of just two elements, namely q = (2n-3)p - (n-1) and q = (2n-3)p - (n-1) + 1. In general the difference of the lowest point of Q_p and the highest point of Q_{p-r} is given by r(2n-3) - 2(n-1). Hence if we raise the interval Q_p by r-1 units it fails to intersect Q_{p-r} , for r > 1.

An obvious transcription of this result is the following proposition with r > 1.

PROPOSITION. If d_r is an endomorphism of E_1 which maps $E_1^{p,q}$ into $E_1^{p-r,q+r-1}$, then d_r annihilates all elements of $E_1^{p,q}$.

Actually this proposition is equally valid for r = 1 because of the simple connectedness of G_{n-1}^2 . The vanishing of the first and therefore by the duality in manifolds the vanishing of the $(2n-3)^{rd}$ homology groups of G_{n-1}^2 implies that $E_1^{p,(2n-3)p-(n+1)+1}$ and $E_1^{p-1,(2n-3)p-(n-1)}$ are trivial. Hence in the only range of q where d_1 could be nontrivial (i.e. for the two values of q where Q_p and Q_{p-1} overlap) once the image and once the range of d_1 is trivial. Therefore d_1 again annihilates every element of $E_1^{p,q}$.

Now let A denote the limit singular chain groups of the directed system $\{\pi_k, h\}$. Let $D \subset A$ be the limit chain group of the set $\{\Delta, h\}$. If we now define $A_p \subset A$ as the limit chain groups of the system $\{(\pi_k^p, h\}, we get a grading of <math>A/D$ in the sense of Leray. (We are again using the notations and conventions as given in Serre [6].) In the resulting spectral sequence Serre's $E_1^{p,q}$ $[E_{\infty}^{p,q}]$ is easily identified with our $E_1^{p,q}$ $[E_{\infty}^{p,q}]$. Above we have just shown that d_1 must be trivial in $E_1^{p,q}$. Hence $E_1 \approx E_2$. But d_2 must again be trivial by our dimension argument. Hence $E_3 \approx E_1$. Proceeding in this fashion we see that $E_{\infty} \approx E_1$.

258

Hence, as pointed out earlier, the rank of $H_m(A/D)$ equals the rank of the *m*-dimensional elements of $E_1^{p,q}$ (that is, the sum of all the ranks of $E_1^{p,q}$ with p + q = m). It follows that

$$\mathfrak{M}(S^{n}; z) = P(G_{n-1}^{2}; z) z^{(n-1)} (1 - z^{2(n-1)})^{-1}.$$

REMARK 1. For readers not acquainted with the Leray spectral sequence it should be pointed out that the use of this powerful tool in the last section can be circumvented in a sense just because all the differentials d_r are trivial. Essentially, the "Lückenverfahren" of Seifert and Threlfall (see [2], p. 73) applies. Thus the statement that all the d_r 's are trivial is equivalent to the statement in the Morse terminology that the critical manifolds are of "increasing type", or again, equivalently, the "linking cycles" can be extended "below" the critical level.

REMARK 2. The circular connectivities (as distinguished from the *sensed* circular connectivities!) of the *n*-sphere, are obtained in a precisely analogous fashion. If we bar all the letters in \$ and 5 these sections can as a matter of fact be read in that context.

The only essential difference occurs in §6. Here the d_r 's are trivial for r > 1 as before, but as \tilde{G}_{n-1}^2 is not simply connected it is not a priori obvious that d_1 is trivial. Nevertheless d_1 turns out to be trivial, and we therefore get the following formula for $\overline{\mathfrak{M}}(S^n; z)$:

$$\overline{\mathfrak{M}}(S^{n};z) = P(\overline{G}_{n-1}^{2};z) \ z^{n-1}(1 \ - \ z^{2(n-1)})^{-1}.$$

REMARK 3. It might be of interest to compare the procedure given here with that of Morse in [1].

Morse uses for his model of S^n ellipsoids on which certain closed geodesics are completely isolated. We on the other hand exploit precisely the degeneracy of the *n* sphere in its usual representation. This avoids first of all the study of geodesics on ellipsoids. In particular, in the Morse treatment no *fixed* model will do for the computation in all dimensions. (On any fixed ellipsoid, for instance, the indices of the iterates of certain geodesics behave regularly only for a finite number of iterations.)

Secondly we avoid the explicit construction of the linking cycles λ_{ij}^r (see [1], p. 323. As mentioned earlier their existence is equivalent to the vanishing of all the d_r 's.

Thirdly, the computation of $\mathfrak{M}(J_c, J_{c-})$ differs from that of Morse. The notion of a nondegenerate critical manifold enters essentially here, as even if $\sigma \ \epsilon \ \mu_k$ represents only a single isolated geodesic it will still be a 1-sphere of critical points.

These then are roughly the main deviations. They are of a more or less technical nature. The extent to which the present derivation of $\overline{\mathfrak{M}}(S^n; z)$ is based on the Morse theory is amply given by the basic theorems which we quote from Morse's treatise.

RAOUL BOTT

7. The general theory

It is evident that in certain parts of our computation no reference was made to the special case of the n-sphere which we had in mind. The purpose of this section is to abstract from the preceding pages those features which are valid in general. We state the result as a theorem and corollary.

THEOREM II. Let R be a Riemann manifold. Let $\Pi_k(R) \cdots$ etc. and J be defined as in §2 and let $\mu_k \cdots$ etc. be defined as in §5. Let $\{\tilde{\sigma}_{t,k}\}$ be critical sets of J on $\Pi_k(R)$ (for k sufficiently large) such that $h_s: \tilde{\sigma}_{t,k} \to \tilde{\sigma}_{t,ks}$, on which J = t. Suppose further that (a) $\sigma_{t,k}$ is a nondegenerate critical manifold of $Ju_k(R)$, of index λ_0 . (b) $\sigma_{t,k}$ can be fibered by one spheres S^1 in the obvious fashion. (S^1 acts on $\sigma_{t,k}$ by rotating the closed geodesics.) Let the base-space of this fibering be $\sigma_t^* = \sigma_{t,k}^*$.

Then if $\{V_{t,k}\}$ is a sufficiently small family of neighborhoods of $\sigma_{t,k}(h_s V_{t,k} \subset V_{t,ks})$ in $\mu_k(R)$ and $V_{t,k}^* = \pi_k^t(R) \cap V_{t,k}$; $V_{k,k}^- = \pi_k^{t-1}(R) \cap V_{t,k}$

 $\lim_{k} H_m(V_{t,k}^*, V_{t,k}^-) \approx H_{m-\lambda_{\sigma}}(\sigma_t^*)$

COROLLARY. Let R be a Riemann manifold such that:

a. The critical values of J (on any $\tilde{\pi}_k$) form a countable sequence of numbers $n_1 < n_2 < n_3$ etc.

b. The critical set corresponding to n_r on μ_k is a nondegenerate critical manifold $\sigma_{n_r,k}$ of index λ_{n_r} , satisfying condition b above.

Then $\mathfrak{M}(R)$ has an associated graded group $\mathfrak{GM}(R)$ (see [6]) which is the terminal group E_{∞} of a spectral sequence $E_1, E_2 \cdots E_{\infty}$ of which the first term E_1 is determined by the formula:

$$E_{1}^{p,q} = H_{p+q-\lambda_{n_{p}}}(\sigma_{n_{p}}^{*}), \qquad p > 0$$

$$E_{1}^{0,q} = \phi,$$

provided $E_1^{p,q}$ is finitely generated in each dimension (i.e., for p + q = constant).

REMARK. A given Riemann manifold may of course give rise to quite different groups E_1 when in different metrics. The passage from E_1 to E_{∞} differs correspondingly. Thus for instance if we take a nonspherical ellipsoid for our model of S^n the vanishing of all the d_r 's in the resulting E_1 is by no means a priori obvious, and is, as a matter of fact, *never* true. The proof of this proposition is connected with the study of the "Iteration of closed geodesics" (see [7], and will be treated at another time.

University of Michigan

BIBLIOGRAPHY

- 1. MORSE, M. The calculus of variation in the large. Amer. Math. Soc. Colloquium Publications, vol. XVIII.
- 2. H. SEIFERT and W. THRELFALL. Variations rechnung im Grossen. Chelsea Publ. Co. 1948
- 3. W. GYSIN. Zur Homologie theorie der Abbildungen und Faserungen der Manigfaltigkeiten. Comm. Math. Helv. 14 (1941) p. 61–122.
- R. THOM. Classes caracteristiques et i-carrés. C. R. Acad. Sci. Paris 230, (1950) p. 427-429.

- 5. S. S. CHERN and E. SPANIER. The homology structure of sphere bundles. Proc. Nat. Acad. Sci. U. S. A., 36 (1950) p. 3-27.
- J. P. SERRE. Homologie singulière des espaces fibrés. Ann. of Math., 54, (1951) p. 425– 505.
- MORSE and PITCHER. On certain invariants of closed extremals. Proc. Nat. Acad. Sci. U. S. A., 20, (1934) p. 282-287.
- 8. N. STEENROD. The topology of fiber bundles. Princeton University Press, 1951.
- 9. S. EILENBERG and N. STEENROD. Foundations of algebraic topology. Princeton University Press 1952.
- A. BOREL. Sur la Cohomologie des espaces fibres principaux et des espaces homogènes, de groupes de Lie Compact. Ann. of Math., 57 (1953), 115-207.
- 11. H. CARTAN. Seminaire de Topologie Algébrique de l'E. N. S. (1950-51), Notes polycopies.