

# Hodge Theory on Metric Spaces

Laurent Bartholdi\*

Georg-August-Universität Göttingen  
Deutschland

Thomas Schick\*

Georg-August-Universität Göttingen  
Deutschland

Nat Smale

University of Utah

Steve Smale<sup>†</sup>

City University of Hong Kong

*With an appendix by Anthony W. Baker<sup>‡</sup>  
Mathematics and Computing Technology  
The Boeing Company.*

Last compiled December 1, 2009; last edited December 1, by LB

## Abstract

Hodge theory is a beautiful synthesis of geometry, topology, and analysis, which has been developed in the setting of Riemannian manifolds. On the other hand, spaces of images, which are important in the mathematical foundations of vision and pattern recognition, do not fit this framework. This motivates us to develop a version of Hodge theory on metric spaces with a probability measure. We believe that this constitutes a step towards understanding the geometry of vision.

The appendix by Anthony Baker provides a separable, compact metric space with infinite dimensional  $\alpha$ -scale homology.

## 1 Introduction

Hodge Theory [21] studies the relationships of topology, functional analysis and geometry of a manifold. It extends the theory of the Laplacian on domains of Euclidean space or on a manifold.

---

\*email: [laurent@uni-math.gwdg.de](mailto:laurent@uni-math.gwdg.de) and [schick@uni-math.gwdg.de](mailto:schick@uni-math.gwdg.de)  
www: <http://www.uni-math.gwdg.de/schick>

Laurent Bartholdi and Thomas Schick were partially supported by the Courant Research Center “Higher order structures in Mathematics” of the German Initiative of Excellence

<sup>†</sup>Steve Smale was supported in part by the NSF, and the Toyota Technological Institute, Chicago

<sup>‡</sup>email: [anthony.w.baker@boeing.com](mailto:anthony.w.baker@boeing.com)

However, there are a number of spaces, not manifolds, which could benefit from an extension of Hodge theory, and that is the motivation here. In particular we believe that a deeper analysis in the theory of vision could be led by developments of Hodge type. Spaces of images are important for developing a mathematics of vision (see e.g. Smale, Rosasco, Bouvrie, Caponnetto, and Poggio [32]); but these spaces are far from possessing manifold structures. Other settings include spaces occurring in quantum field theory, manifolds with singularities and/or non-uniform measures.

A number of previous papers have given us inspiration and guidance. For example there are those in combinatorial Hodge theory of Eckmann [16], Dódziuk [13], Friedman [19], and more recently Jiang, Lim, Yao, and Ye [22]. Recent decades have seen extensions of the Laplacian from its classical setting to that of combinatorial graph theory. See e.g. Fan Chung [9]. Robin Forman [18] has useful extensions from manifolds. Further extensions and relationships to the classical settings are Belkin, Niyogi [2], Belkin, De Vito, and Rosasco [3], Coifman, Maggioni [10], and Smale, Zhou [33].

Our approach starts with a metric space  $X$  (complete, separable), endowed with a probability measure. For  $\ell \geq 0$ , an  $\ell$ -form is a function on  $(\ell + 1)$ -tuples of points in  $X$ . The coboundary operator  $\delta$  is defined from  $\ell$ -forms to  $(\ell + 1)$ -forms in the classical way following Čech, Alexander, and Spanier. Using the  $L^2$ -adjoint  $\delta^*$  of  $\delta$  for a boundary operator, the  $\ell$ th order Hodge operator on  $\ell$ -forms is defined by  $\Delta_\ell = \delta^* \delta + \delta \delta^*$ . The harmonic  $\ell$ -forms on  $X$  are solutions of the equation  $\Delta_\ell(f) = 0$ . The  $\ell$ -harmonic forms reflect the  $\ell$ th homology of  $X$  but have geometric features. The harmonic form is a special representative of the homology class and it may be interpreted as one satisfying an optimality condition. Moreover, the Hodge equation is linear and by choosing a finite sample from  $X$  one can obtain an approximation of this representative by a linear equation in finite dimension.

There are two avenues to develop this Hodge theory. The first is a kernel version corresponding to a Gaussian or a reproducing kernel Hilbert space. Here the topology is trivial but the analysis gives a substantial picture. The second version is akin to the adjacency matrix of graph theory and corresponds to a threshold at a given scale  $\alpha$ . When  $X$  is finite this picture overlaps with that of the combinatorial Hodge theory referred to above.

For passage to a continuous Hodge theory, one encounters:

**Problem 1** (Poisson Regularity Problem). If  $\Delta_\ell(f) = g$  is continuous, under what conditions is  $f$  continuous?

It is proved that a positive solution of the Poisson Regularity Problem implies a complete Hodge decomposition for continuous  $\ell$ -forms in the “adjacency matrix” setting (at any scale  $\alpha$ ), provided the  $L^2$ -cohomology is finite dimensional. The problem is solved affirmatively for some cases as  $\ell = 0$ , or  $X$  is finite. One special case is

**Problem 2.** Under what conditions are harmonic  $\ell$ -forms continuous?

Here we have a solution for  $\ell = 0$  and  $\ell = 1$ .

The solution of these regularity problems would be progress toward the important cohomology identification problem: To what extent does the  $L^2$ -cohomology coincide with the classical cohomology? We have an answer to this question, as well as a full Hodge theory in the special, but important case of Riemannian manifolds. The following theorem is proved in Section 9 of this paper.

**Theorem 1.** *Suppose that  $M$  is a compact Riemannian manifold, with strong convexity radius  $r$  and that  $k > 0$  is an upper bound on the sectional curvatures. Then, if  $0 < \alpha < \max\{r, \sqrt{\pi}/2k\}$ , our Hodge theory holds. That is, we have a Hodge decomposition, the kernel of  $\Delta_\ell$  is isomorphic to the  $L^2$ -cohomology, and to the de Rham cohomology of  $M$  in degree  $\ell$ .*

More general conditions on a metric space  $X$  are given in Section 9.

Certain previous studies show how topology questions can give insight into the study of images. Lee, Pedersen, and Mumford [24] have investigated  $3 \times 3$  pixel images from real world data bases to find the evidence for the occurrence of homology classes of degree 1. Moreover, Carlsson, Ishkhanov, de Silva, and Zomorodian [5] have found evidence for homology of surfaces in the same data base. Here we are making an attempt to give some foundations to these studies. Moreover, this general Hodge theory could yield optimal representatives of the homology classes and provide systematic algorithms.

Related in spirit to our  $L^2$ -cohomology, but in a quite different setting, is the  $L^2$ -cohomology as introduced by Atiyah [1]. This is defined either via  $L^2$ -differential forms [1] or combinatorially [14], but again with an  $L^2$  condition. Questions like the Hodge decomposition problem also arise in this setting, and its failure gives rise to additional invariants, the Novikov-Shubin invariants. This theory has been extensively studied, compare e.g. [8, 26, 30, 25] for important properties and geometric as well as algebraic applications. In [27, 31, 15] approximation of the  $L^2$ -Betti numbers for infinite simplicial complexes in terms of associated finite simplicial complexes is discussed in increasing generality. Complete calculations of the spectrum of the associated Laplacian are rarely possible, but compare [11] for one of these cases. The monograph [28] provides rather complete information about this theory.

Here is some background to the writing of this paper. Essentially Sections 2 through 8 were in a finished paper by Nat Smale and Steve Smale, February 20, 2009. That version stated that the coboundary operator of Theorem 4, Section 4 must have a closed image. Thomas Schick pointed out that this assertion was wrong, and in fact produced a counterexample, now Section 10 of this paper. Moreover, Schick and Laurent Bartholdi set in motion the proofs that give the sufficient conditions for the finite dimensionality of the  $L^2$ -cohomology groups in Section 9 of this paper, and hence the property that the image of the coboundary is closed. In particular Theorems 7 and 8 were proved by them.

Some conversations with Shmuel Weinberger were helpful.

## 2 An $L^2$ -Hodge Theory

In this section we construct a general Hodge Theory for certain  $L^2$ -spaces. The amount of structure needed for this theory is minimal. First, let us introduce some notation used throughout the section.  $X$  will denote a set endowed with a probability measure  $\mu$  ( $\mu(X) = 1$ ). The  $\ell$ -fold cartesian product of  $X$  will be denoted as  $X^\ell$  and  $\mu_\ell$  will denote the product measure on  $X^\ell$ . Furthermore, we will assume the existence of a kernel function  $K: X^2 \rightarrow \mathbb{R}$ , a non-negative, measurable, symmetric function which we will assume is in  $L^\infty(X \times X)$ , and for certain results, we will impose additional assumptions on  $K$ . A useful example to keep in mind is this.  $X$  is a compact domain in Euclidean space,  $\mu$  a Borel measure, but not necessarily the Euclidean measure, and  $K$  a Gaussian kernel  $K(x, y) = e^{-\frac{\|x-y\|^2}{\sigma}}$ ,  $\sigma > 0$ . A simpler example is  $K \equiv 1$ , but the Gaussian example contains the notion of locality ( $K(x, y)$  is close to 1 just when  $x$  is near  $y$ ).

Recall that a chain complex of vector spaces is a sequence of vector spaces  $V_j$  and linear maps  $d_j: V_j \rightarrow V_{j-1}$  such that the composition  $d_{j-1} \circ d_j = 0$ . A co-chain complex is the same, except that  $d_j: V_j \rightarrow V_{j+1}$ . The basic spaces in this section are  $L^2(X^\ell)$ , from which we will construct chain and cochain complexes:

$$\dots \xrightarrow{\partial_{\ell+1}} L^2(X^{\ell+1}) \xrightarrow{\partial_\ell} L^2(X^\ell) \xrightarrow{\partial_{\ell-1}} \dots L^2(X) \xrightarrow{\partial_0} 0 \quad (1)$$

and

$$0 \longrightarrow L^2(X) \xrightarrow{\delta_0} L^2(X^2) \xrightarrow{\delta_1} \dots \xrightarrow{\delta_{\ell-1}} L^2(X^{\ell+1}) \xrightarrow{\delta_\ell} \dots \quad (2)$$

Here, both  $\partial_\ell$  and  $\delta_\ell$  will be bounded linear maps, satisfying  $\partial_{\ell-1} \circ \partial_\ell = 0$  and  $\delta_\ell \circ \delta_{\ell-1} = 0$ . When there is no confusion, we will omit the subscripts of these operators.

We first define  $\delta = \delta_{\ell-1}: L^2(X^\ell) \rightarrow L^2(X^{\ell+1})$  by

$$\delta f(x_0, \dots, x_\ell) = \sum_{i=0}^{\ell} (-1)^i \prod_{j \neq i} \sqrt{K(x_i, x_j)} f(x_0, \dots, \hat{x}_i, \dots, x_\ell) \quad (3)$$

where  $\hat{x}_i$  means that  $x_i$  is deleted. This is similar to the co-boundary operator of Alexander-Spanier cohomology (see Spanier [34]). The square root in the formula is unimportant for most of the sequel, and is there so that when we define the Laplacian on  $L^2(X)$ , we recover the operator as defined in Gilboa and Osher [20]. We also note that in the case  $X$  is a finite set,  $\delta_0$  is essentially the same as the gradient operator developed by Zhou and Schölkopf [37] in the context of learning theory.

**Proposition 1.** *For all  $\ell \geq 0$ ,  $\delta: L^2(X^\ell) \rightarrow L^2(X^{\ell+1})$  is a bounded linear map.*

*Proof.* Clearly  $\delta f$  is measurable, as  $K$  is measurable. Since  $\|K\|_\infty < \infty$ , it

follows from the Schwartz inequality in  $\mathbb{R}^\ell$  that

$$\begin{aligned} |\delta f(x_0, \dots, x_\ell)|^2 &\leq C \left( \sum_{i=0}^{\ell} |f(x_0, \dots, \hat{x}_i, \dots, x_\ell)| \right)^2 \\ &\leq C(\ell+1) \sum_{i=0}^{\ell} |f(x_0, \dots, \hat{x}_i, \dots, x_\ell)|^2 \end{aligned}$$

where  $C = \|K\|_\infty^\ell$ . Now, integrating both sides of the inequality with respect to  $d\mu_{\ell+1}$ , using Fubini's Theorem on the right side and the fact that  $\mu(X) = 1$  gives us

$$\|\delta f\|_{L^2(X^{\ell+1})} \leq \sqrt{C}(\ell+1) \|f\|_{L^2(X^\ell)},$$

completing the proof.  $\square$

Essentially the same proof shows that  $\delta$  is a bounded linear map on  $L^p$ ,  $p \geq 1$ .

**Proposition 2.** *For all  $\ell \geq 1$ ,  $\delta_\ell \circ \delta_{\ell-1} = 0$ .*

*Proof.* The proof is standard when  $K \equiv 1$ . For  $f \in L^2(X^\ell)$  we have

$$\begin{aligned} &\delta_\ell(\delta_{\ell-1}f)(x_0, \dots, x_{\ell+1}) \\ &= \sum_{i=0}^{\ell+1} (-1)^i \prod_{j \neq i} \sqrt{K(x_i, x_j)} (\delta_{\ell-1}f)(x_0, \dots, \hat{x}_i, \dots, x_{\ell+1}) \\ &= \sum_{i=0}^{\ell+1} (-1)^i \prod_{j \neq i} \sqrt{K(x_i, x_j)} \sum_{k=0}^{i-1} (-1)^k \prod_{n \neq k, i} \sqrt{K(x_k, x_n)} f(x_0, \dots, \hat{x}_k, \dots, \hat{x}_i, \dots, x_{\ell+1}) \\ &\quad + \sum_{i=0}^{\ell+1} (-1)^i \prod_{j \neq i} \sqrt{K(x_i, x_j)} \sum_{k=i+1}^{\ell+1} (-1)^{k-1} \prod_{n \neq k, i} \sqrt{K(x_k, x_n)} f(x_0, \dots, \hat{x}_i, \dots, \hat{x}_k, \dots, x_{\ell+1}) \end{aligned}$$

Now we note that on the right side of the second equality for given  $i, k$ ,  $k < i$ , the corresponding term in the first sum

$$(-1)^{i+k} \prod_{j \neq i} \sqrt{K(x_i, x_j)} \prod_{n \neq k, i} \sqrt{K(x_k, x_n)} f(x_0, \dots, \hat{x}_k, \dots, \hat{x}_i, \dots, x_{\ell+1})$$

cancels the term in the second sum where  $i$  and  $k$  are reversed

$$(-1)^{k+i-1} \prod_{j \neq k} \sqrt{K(x_k, x_j)} \prod_{n \neq k, i} \sqrt{K(x_k, x_n)} f(x_0, \dots, \hat{x}_k, \dots, \hat{x}_i, \dots, x_{\ell+1})$$

because, using the symmetry of  $K$ ,

$$\prod_{j \neq i} \sqrt{K(x_i, x_j)} \prod_{n \neq k, i} \sqrt{K(x_k, x_n)} = \prod_{j \neq k} \sqrt{K(x_k, x_j)} \prod_{n \neq k, i} \sqrt{K(x_i, x_n)}. \quad \square$$

It follows that (2) and (3) define a co-chain complex. We now define, for  $\ell > 0$ ,  $\partial_\ell: L^2(X^{\ell+1}) \rightarrow L^2(X^\ell)$  by

$$\partial_\ell g(x) = \sum_{i=0}^{\ell} (-1)^i \int_X \left( \prod_{j=0}^{\ell-1} \sqrt{K(t, x_j)} \right) g(x_0, \dots, x_{i-1}, t, x_i, \dots, x_{\ell-1}) d\mu(t) \quad (4)$$

where  $x = (x_0, \dots, x_{\ell-1})$  and for  $\ell = 0$  we define  $\partial_0: L^2(X) \rightarrow 0$ .

**Proposition 3.** *For all  $\ell \geq 0$ ,  $\partial_\ell: L^2(X^{\ell+1}) \rightarrow L^2(X^\ell)$  is a bounded linear map.*

*Proof.* For  $g \in L^2(X^{\ell+1})$ , we have

$$\begin{aligned} |\partial_\ell g(x_0, \dots, x_{\ell-1})| &\leq \|K\|_\infty^{\ell-1} \sum_{i=0}^{\ell} \int_X |g(x_0, \dots, x_{i-1}, t, \dots, x_{\ell-1})| d\mu(t) \\ &\leq \|K\|_\infty^{\ell-1} \sum_{i=0}^{\ell} \left( \int_X |g(x_0, \dots, x_{i-1}, t, \dots, x_{\ell-1})|^2 d\mu(t) \right)^{\frac{1}{2}} \\ &\leq \|K\|_\infty^{\ell-1} \sqrt{\ell+1} \left( \sum_{i=0}^{\ell} \int_X |g(x_0, \dots, x_{i-1}, t, \dots, x_{\ell-1})|^2 d\mu(t) \right)^{\frac{1}{2}} \end{aligned}$$

where we have used the Schwartz inequalities for  $L^2(X)$  and  $\mathbb{R}^{\ell+1}$  in the second and third inequalities respectively. Now, square both sides of the inequality and integrate over  $X^\ell$  with respect to  $\mu_\ell$  and use Fubini's Theorem arriving at the following bound to finish the proof:

$$\|\partial_\ell g\|_{L^2(X^\ell)} \leq \|K\|_\infty^{\ell-1} (\ell+1) \|g\|_{L^2(X^{\ell+1})}. \quad \square$$

**Remark 1.** As in Proposition 1, we can replace  $L^2$  by  $L^p$ , for  $p \geq 1$ .

We now show that (for  $p = 2$ )  $\partial_\ell$  is actually the adjoint of  $\delta_{\ell-1}$  (which gives a second proof of Proposition 3).

**Proposition 4.**  $\delta_{\ell-1}^* = \partial_\ell$ . *That is  $\langle \delta_{\ell-1} f, g \rangle_{L^2(X^{\ell+1})} = \langle f, \partial_\ell g \rangle_{L^2(X^\ell)}$  for all  $f \in L^2(X^\ell)$  and  $g \in L^2(X^{\ell+1})$ .*

*Proof.* For  $f \in L^2(X^\ell)$  and  $g \in L^2(X^{\ell+1})$  we have, by Fubini's Theorem

$$\begin{aligned} \langle \delta_{\ell-1} f, g \rangle &= \sum_{i=0}^{\ell} (-1)^i \int_{X^{\ell+1}} \prod_{j \neq i} \sqrt{K(x_i, x_j)} f(x_0, \dots, \hat{x}_i, \dots, x_\ell) g(x_0, \dots, x_\ell) d\mu_{\ell+1} \\ &= \sum_{i=0}^{\ell} (-1)^i \int_{X^\ell} f(x_0, \dots, \hat{x}_i, \dots, x_\ell) \cdot \int_X \prod_{j \neq i} \sqrt{K(x_i, x_j)} g(x_0, \dots, x_\ell) d\mu(x_i) d\mu(x_0) \cdots \widehat{d\mu(x_i)} \cdots d\mu(x_\ell) \end{aligned}$$

In the  $i$ -th term on the right, relabeling the variables  $x_0, \dots, \hat{x}_i, \dots, x_\ell$  with  $y = (y_0, \dots, y_{\ell-1})$  (that is  $y_j = x_{j+1}$  for  $j \geq i$ ) and putting the sum inside the integral gives us

$$\int_{X^\ell} f(y) \sum_{i=0}^{\ell} (-1)^i \int_X \prod_{j=0}^{\ell-1} \sqrt{K(x_i, y_j)} g(y_0, \dots, y_{i-1}, x_i, y_i, \dots, y_{\ell-1}) d\mu(x_i) d\mu_\ell(y)$$

which is just  $\langle f, \partial_\ell g \rangle$ .  $\square$

We note, as a corollary, that  $\partial_{\ell-1} \circ \partial_\ell = 0$ , and thus (1) and (4) define a chain complex. We can thus define the homology and cohomology spaces (real coefficients) of (1) and (2) as follows. Since  $\text{Im } \partial_\ell \subset \text{Ker } \partial_{\ell-1}$  and  $\text{Im } \delta_{\ell-1} \subset \text{Ker } \delta_\ell$  we define the quotient spaces

$$H_\ell(X) = H_\ell(X, K, \mu) = \frac{\text{Ker } \partial_\ell}{\text{Im } \partial_{\ell-1}} \quad H^\ell(X) = H^\ell(X, K, \mu) = \frac{\text{Ker } \delta_\ell}{\text{Im } \delta_{\ell-1}} \quad (5)$$

which will be referred to the  $L^2$ -homology and cohomology of degree  $\ell$ , respectively. In later sections, with additional assumptions on  $X$  and  $K$ , we will investigate the relation between these spaces and the topology of  $X$ , for example, the Alexander-Spanier cohomology. In order to proceed with the Hodge Theory, we consider  $\delta$  to be the analogue of the exterior derivative  $d$  on  $\ell$ -forms from differential topology, and  $\partial = \delta^*$  as the analogue of  $d^*$ . We then define the Laplacian (in analogy with the Hodge Laplacian) to be  $\Delta_\ell = \delta_\ell^* \delta_\ell + \delta_{\ell-1} \delta_{\ell-1}^*$ . Clearly  $\Delta_\ell: L^2(X^{\ell+1}) \rightarrow L^2(X^{\ell+1})$  is a bounded, self adjoint, positive semi-definite operator since for  $f \in L^2(X^{\ell+1})$

$$\langle \Delta f, f \rangle = \langle \delta^* \delta f, f \rangle + \langle \delta \delta^* f, f \rangle = \|\delta f\|^2 + \|\delta^* f\|^2 \quad (6)$$

where we have left off the subscripts on the operators. The Hodge Theorem will give a decomposition of  $L^2(X^{\ell+1})$  in terms of the image spaces under  $\delta$ ,  $\delta^*$  and the kernel of  $\Delta$ , and also identify the kernel of  $\Delta$  with  $H^\ell(X, K, \mu)$ . Elements of the kernel of  $\Delta$  will be referred to as harmonic. For  $\ell = 0$ , one easily computes that

$$\frac{1}{2} \Delta_0 f(x) = D(x) f(x) - \int_X K(x, y) f(y) d\mu(y) \quad \text{where } D(x) = \int_X K(x, y) d\mu(y)$$

which, in the case  $K$  is a positive definite kernel on  $X$ , is the Laplacian defined in Smale and Zhou [33] (see section 5 below).

**Remark 2.** It follows from (6) that  $\Delta f = 0$  if and only if  $\delta_\ell f = 0$  and  $\delta_\ell^* f = 0$ , and so  $\text{Ker } \Delta_\ell \subset \text{Ker } \delta_\ell$ .

The main goal of this section is the following  $L^2$ -Hodge theorem.

**Theorem 2.** Assume that  $0 < \sigma \leq K(x, y) \leq \|K\|_\infty < \infty$  almost everywhere. Then we have the orthogonal, direct sum decomposition

$$L^2(X^{\ell+1}) = \text{Im } \delta_{\ell-1} \oplus \text{Im } \delta_\ell^* \oplus \text{Ker } \Delta_\ell$$

and the cohomology space  $H^\ell(X, K, \mu)$  is isomorphic to  $\text{Ker } \Delta_\ell$ , with each equivalence class in the former having a unique representative in the latter.

In this case  $H^\ell(X) = 0$  for  $\ell > 0$  and  $H^0(X) = \mathbb{R}$ . Indeed, the theorem holds as long as  $\delta_\ell$  (or equivalently  $\partial_\ell$ ) has closed range for all  $\ell$ .

In subsequent sections we will have occasion to use the  $L^2$ -spaces of alternating functions:

$$L_a^2(X^{\ell+1}) = \{f \in L^2(X^{\ell+1}) : f(x_0, \dots, x_\ell) = (-1)^{\text{sign } \sigma} f(x_{\sigma(x_0)}, \dots, x_{\sigma(x_\ell)}), \\ \sigma \text{ a permutation}\}$$

Due to the symmetry of  $K$ , it is easy to check that the coboundary  $\delta$  preserves the alternating property, and thus Propositions 1 through 4, as well as formulas (1), (2), (5) and (6) hold with  $L_a^2$  in place of  $L^2$ . We note that the alternating map

$$\text{Alt} : L^2(X^{\ell+1}) \rightarrow L_a^2(X^{\ell+1})$$

defined by

$$\text{Alt}(f)(x_0, \dots, x_\ell) := \frac{1}{(\ell+1)!} \sum_{\sigma \in S_{\ell+1}} (-1)^{\text{sign } \sigma} f(x_{\sigma(x_0)}, \dots, x_{\sigma(x_\ell)})$$

is a projection relating the two definitions of  $\ell$ -forms. It is easy to compute that this is actually an orthogonal projection, its inverse is just the inclusion map.

**Remark 3.** It follows from homological algebra that these maps induce inverse to each other isomorphisms of the cohomology groups we defined. Indeed, there is a standard chain homotopy between a variant of the projection  $\text{Alt}$  and the identity, given by  $hf(x_0, \dots, x_n) = \frac{1}{n} \sum_{i=0}^n f(x_i, x_0, \dots, x_n)$ . Because many formulas simplify, from now on we will therefore most of the time work with the subcomplex of alternating functions.

We first recall some relevant facts in a more abstract setting in the following

**Lemma 1** (Hodge Lemma). *Suppose we have the cochain and corresponding dual chain complexes*

$$\begin{array}{ccccccc} 0 & \longrightarrow & V_0 & \xrightarrow{\delta_0} & V_1 & \xrightarrow{\delta_1} & \dots & \xrightarrow{\delta_{\ell-1}} & V_\ell & \xrightarrow{\delta_\ell} & \dots \\ & & & & & & & & & & \\ \dots & & \xrightarrow{\delta_\ell^*} & V_\ell & \xrightarrow{\delta_{\ell-1}^*} & V_{\ell-1} & \xrightarrow{\delta_{\ell-2}^*} & \dots & \xrightarrow{\delta_0^*} & V_0 & \longrightarrow & 0 \end{array}$$

where for  $\ell = 0, 1, \dots$ ,  $V_\ell, \langle, \rangle_\ell$  is a Hilbert space,  $\delta_\ell$  (and thus  $\delta_\ell^*$ , the adjoint of  $\delta_\ell$ ) is a bounded linear map with  $\delta^2 = 0$ . Let  $\Delta_\ell = \delta_\ell^* \delta_\ell + \delta_{\ell-1} \delta_{\ell-1}^*$ . Then the following are equivalent

- (1)  $\delta_\ell$  has closed range for all  $\ell$ ;
- (2)  $\delta_\ell^*$  has closed range for all  $\ell$ .



Furthermore, if one of the above conditions hold, we have the orthogonal, direct sum decomposition into closed subspaces

$$V_\ell = \text{Im } \delta_{\ell-1} \oplus \text{Im } \delta_\ell^* \oplus \text{Ker } \Delta_\ell$$

and the quotient space  $\frac{\text{Ker } \delta_\ell}{\text{Im } \delta_{\ell-1}}$  is isomorphic to  $\text{Ker } \Delta_\ell$ , with each equivalence class in the former having a unique representative in the latter.

*Proof.* We first assume conditions 1 and 2 above and prove the decomposition. For all  $f \in V_{\ell-1}$  and  $g \in V_{\ell+1}$  we have

$$\langle \delta_{\ell-1} f, \delta_\ell^* g \rangle_\ell = \langle \delta_\ell \delta_{\ell-1} f, g \rangle_{\ell+1} = 0.$$

Also, as in (6),  $\Delta_\ell f = 0$  if and only if  $\delta_\ell f = 0$  and  $\delta_{\ell-1}^* f = 0$ . Therefore, if  $f \in \text{Ker } \Delta_\ell$ , then for all  $g \in V_{\ell-1}$  and  $h \in V_{\ell+1}$

$$\langle f, \delta_{\ell-1} g \rangle_\ell = \langle \delta_{\ell-1}^* f, g \rangle_{\ell-1} = 0 \quad \text{and} \quad \langle f, \delta_\ell^* h \rangle_\ell = \langle \delta_\ell f, h \rangle_{\ell+1} = 0$$

and thus  $\text{Im } \delta_{\ell-1}$ ,  $\text{Im } \delta_\ell^*$  and  $\text{Ker } \Delta_\ell$  are mutually orthogonal. Now, since  $\text{Im } \delta_{\ell-1} \oplus \text{Im } \delta_\ell^*$  is closed, to prove the decomposition it suffices to show that  $\text{Ker } \Delta_\ell \supseteq (\text{Im } \delta_{\ell-1} \oplus \text{Im } \delta_\ell^*)^\perp$ . Let  $v \in (\text{Im } \delta_{\ell-1} \oplus \text{Im } \delta_\ell^*)^\perp$ . Then, for all  $w \in V_\ell$ ,

$$\langle \delta_\ell v, w \rangle = \langle v, \delta_\ell^* w \rangle = 0 \quad \text{and} \quad \langle \delta_{\ell-1}^* v, w \rangle = \langle v, \delta_{\ell-1} w \rangle = 0,$$

which implies that  $\delta_\ell v = 0$  and  $\delta_{\ell-1}^* v = 0$  and as noted above this implies that  $\Delta_\ell v = 0$ , proving the decomposition.

We define an isomorphism

$$\tilde{P}: \frac{\text{Ker } \delta_\ell}{\text{Im } \delta_{\ell-1}} \rightarrow \text{Ker } \Delta_\ell$$

as follows. Let  $P: V_\ell \rightarrow \text{Ker } \Delta_\ell$  be the orthogonal projection. Then, for an equivalence class  $[f] \in \frac{\text{Ker } \delta_\ell}{\text{Im } \delta_{\ell-1}}$  define  $\tilde{P}([f]) = P(f)$ . Note that if  $[f] = [g]$  then  $f = g + h$  with  $h \in \text{Im } \delta_{\ell-1}$ , and therefore  $P(f) - P(g) = P(h) = 0$  by the orthogonal decomposition, and so  $\tilde{P}$  is well defined, and linear as  $P$  is linear. If  $\tilde{P}([f]) = 0$  then  $P(f) = 0$  and so  $f \in \text{Im } \delta_{\ell-1} \oplus \text{Im } \delta_\ell^*$ . But  $f \in \text{Ker } \delta_\ell$ , and so, for all  $g \in V_{\ell+1}$  we have  $\langle \delta_\ell^* g, f \rangle = \langle g, \delta_\ell f \rangle = 0$ , and thus  $f \in \text{Im } \delta_{\ell-1}$  and therefore  $[f] = 0$  and  $\tilde{P}$  is injective. On the other hand,  $\tilde{P}$  is surjective because, if  $w \in \text{Ker } \Delta_\ell$ , then  $w \in \text{Ker } \delta_\ell$  and so  $\tilde{P}([w]) = P(w) = w$ .

Finally, the equivalence of conditions 1 and 2 is a general fact about Hilbert spaces. If  $\delta: V \rightarrow H$  is a bounded linear map between Hilbert spaces, and  $\delta^*$  is its adjoint, and if  $\text{Im } \delta$  is closed in  $H$ , then  $\text{Im } \delta^*$  is closed in  $V$ . We include the proof for completeness. Since  $\text{Im } \delta$  is closed, the bijective map

$$\delta: (\text{Ker } \delta)^\perp \rightarrow \text{Im } \delta$$

is an isomorphism by the open mapping theorem. It follows that

$$\inf \{ \|\delta(v)\| : v \in (\text{Ker } \delta)^\perp, \|v\| = 1 \} > 0$$

Since  $\text{Im } \delta \subset (\text{Ker } \delta^*)^\perp$ , it suffices to show that

$$\delta^* \delta: (\text{Ker } \delta)^\perp \rightarrow (\text{Ker } \delta)^\perp$$

is an isomorphism, for then  $\text{Im } \delta^* = (\text{Ker } \delta)^\perp$  which is closed. However, this is established by noting that  $\langle \delta^* \delta v, v \rangle = \|\delta v\|^2$  and the above inequality imply that

$$\inf\{\langle \delta^* \delta v, v \rangle : v \in (\text{Ker } \delta)^\perp, \|v\| = 1\} > 0.$$

This finishes the proof of the lemma.  $\square$

**Corollary 1.** *For all  $\ell \geq 0$  the following are isomorphisms, provided  $\text{Im}(\delta)$  is closed.*

$$\delta_\ell: \text{Im } \delta_\ell^* \rightarrow \text{Im } \delta_\ell \quad \text{and} \quad \delta_\ell^*: \text{Im } \delta_\ell \rightarrow \text{Im } \delta_\ell^*$$

*Proof.* The first map is injective because if  $\delta(\delta^* f) = 0$  then  $0 = \langle \delta \delta^* f, f \rangle = \|\delta^* f\|^2$  and so  $\delta f = 0$ . It is surjective because of the decomposition (leaving out the subscripts)

$$\delta(V) = \delta(\text{Im } \delta \oplus \text{Im } \delta^* \oplus \text{Ker } \Delta) = \delta(\text{Im } \delta^*)$$

since  $\delta$  is zero on the first and third summands of the left side of the second equality. The argument for the second map is the same.  $\square$

The difficulty in applying the Hodge Lemma is in verifying that either  $\delta$  or  $\delta^*$  has closed range. A sufficient condition is the following, first pointed out to us by Shmuel Weinberger.

**Proposition 5.** *Suppose that in the context of Lemma 1, the  $L^2$ -cohomology space  $\text{Ker } \delta_\ell / \text{Im } \delta_{\ell-1}$  is finite dimensional. Then  $\delta_{\ell-1}$  has closed range.*

*Proof.* We show more generally, that if  $T: B \rightarrow V$  is a bounded linear map of Banach spaces, with  $\text{Im } T$  having finite codimension in  $V$  then  $\text{Im } T$  is closed in  $V$ . We can assume without loss of generality that  $T$  is injective, by replacing  $B$  with  $B / \text{Ker } T$  if necessary. Thus  $T: B \rightarrow \text{Im } T \oplus F = V$  where  $\dim F < \infty$ . Now define  $G: B \oplus F \rightarrow V$  by  $G(x, y) = Tx + y$ .  $G$  is bounded, surjective and injective, and thus an isomorphism by the open mapping theorem. Therefore  $G(B) = T(B)$  is closed in  $V$ .  $\square$

We now finish the proof of Theorem 2. Consider first the special case where  $K(x, y) = 1$  for all  $x, y$  in  $X$ . Let  $\partial_\ell^0$  be the corresponding operator in (4). We have

**Lemma 2.** *For  $\ell > 1$ ,  $\text{Im } \partial_\ell^0 = \text{Ker } \partial_{\ell-1}^0$ , and  $\text{Im } \partial_1^0 = \{1\}^\perp$  the orthogonal complement of the constants in  $L^2(X)$ .*

Of course this implies that  $\text{Im } \partial_\ell$  is closed for all  $\ell$  since null spaces and orthogonal complements are closed, and in fact shows that the homology (5) in this case is trivial for  $\ell > 0$  and one dimensional for  $\ell = 0$ .

*Proof of Lemma 2.* Let  $h \in \{1\}^\perp \subset L^2(X)$ . Define  $g \in L^2(X^2)$  by  $g(x, y) = h(y)$ . Then from (4)

$$\partial_1^0 g(x_0) = \int_X (g(t, x_0) - g(x_0, t)) d\mu(t) = \int_X (h(x_0) - h(t)) d\mu(t) = h(x_0)$$

since  $\mu(X) = 1$  and  $\int_X h d\mu = 0$ . It can be easily checked that  $\partial_1^0$  maps  $L^2(X^2)$  into  $\{1\}^\perp$ , thus proving the lemma for  $\ell = 1$ . For  $\ell > 1$  let  $h \in \text{Ker } \partial_{\ell-1}^0$ . Define  $g \in L^2(X^{\ell+1})$  by  $g(x_0, \dots, x_\ell) = (-1)^\ell h(x_0, \dots, x_{\ell-1})$ . Then, by (4)

$$\begin{aligned} \partial_\ell^0 g(x_0, \dots, x_{\ell-1}) &= \sum_{i=0}^{\ell} (-1)^i \int_X g(x_0, \dots, x_{i-1}, t, x_i, \dots, x_{\ell-1}) d\mu(t) \\ &= (-1)^\ell \sum_{i=0}^{\ell-1} (-1)^i \int_X h(x_0, \dots, x_{i-1}, t, x_i, \dots, x_{\ell-2}) d\mu(t) \\ &\quad + (-1)^{2\ell} h(x_0, \dots, x_{\ell-1}) \\ &= (-1)^\ell \partial_{\ell-1}^0 h(x_0, \dots, x_{\ell-2}) + h(x_0, \dots, x_{\ell-1}) \\ &= h(x_0, \dots, x_{\ell-1}) \end{aligned}$$

since  $\partial_{\ell-1}^0 h = 0$ , finishing the proof.  $\square$

The next lemma give some general conditions on  $K$  that guarantee  $\partial_\ell$  has closed range.

**Lemma 3.** *Assume that  $K(x, y) \geq \sigma > 0$  for all  $x, y \in X$ . Then  $\text{Im } \partial_\ell$  is closed for all  $\ell$ . In fact,  $\text{Im } \partial_\ell = \text{Ker } \partial_{\ell-1}$  for  $\ell > 1$  and has co-dimension one in  $L^2(X)$  for  $\ell = 1$ .*

*Proof.* Let  $M_\ell: L^2(X^\ell) \rightarrow L^2(X^\ell)$  be the multiplication operator

$$M_\ell(f)(x_0, \dots, x_\ell) = \prod_{j \neq k} \sqrt{K(x_j, x_k)} f(x_0, \dots, x_\ell)$$

Since  $K \in L^\infty(X^2)$  and is bounded below by  $\sigma$ ,  $M_\ell$  clearly defines an isomorphism. The Lemma then follows from Lemma 2, and the observation that

$$\partial_\ell = M_{\ell-1}^{-1} \circ \partial_\ell^0 \circ M_\ell. \quad \square$$

Theorem 2 now follows from the Hodge Lemma, and Lemma 3.

Note that in these, the cohomology is trivial. We also note that Lemma 2, Lemma 3 and Theorem 2 hold in the alternating setting, when  $L^2(X^\ell)$  is replaced with  $L_a^2(X^\ell)$ .

For background, one could see Munkres [22] for the algebraic topology, Lang [23] for the analysis, and Warner [35] for the geometry.

### 3 Metric spaces

For the rest of the paper, we assume that  $X$  is a complete, separable metric space, and that  $\mu$  is a Borel probability measure on  $X$ , and  $K$  is a continuous function on  $X^2$  (as well as symmetric, non-negative and bounded as in Section 2). We will also assume throughout the rest of the paper that  $\mu(U) > 0$  for  $U$  any nonempty open set.

The goal of this section is a Hodge Decomposition for continuous alternating functions. Let  $C(X^{\ell+1})$  denote the continuous functions on  $X^{\ell+1}$ . We will use the following notation:

$$C^{\ell+1} = C(X^{\ell+1}) \cap L_a^2(X^{\ell+1}) \cap L^\infty(X^{\ell+1}).$$

Note that

$$\delta: C^{\ell+1} \rightarrow C^{\ell+2} \quad \text{and} \quad \partial: C^{\ell+1} \rightarrow C^\ell$$

are well defined linear maps. The only thing to check is that  $\delta(f)$  and  $\partial(f)$  are continuous and bounded if  $f \in C^{\ell+1}$ . In the case of  $\delta(f)$  this is obvious from (3). The following proposition from analysis, (4) and the fact that  $\mu$  is Borel imply that  $\partial(f)$  is bounded and continuous.

**Proposition 6.** *Let  $Y$  and  $X$  be metric spaces,  $\mu$  a Borel measure on  $X$ , and  $M, g \in C(Y \times X) \cap L^\infty(Y \times X)$ . Then  $dg \in C(X) \cap L^\infty(X)$ , where*

$$dg(x) = \int_X M(x, t)g(x, t) d\mu(t).$$

*Proof.* The fact that  $dg$  is bounded follows easily from the definition and properties of  $M$  and  $g$ , and continuity follows from a simple application of the Dominated Convergence Theorem, proving the proposition.  $\square$

Therefore we have the chain complexes:

$$\dots \xrightarrow{\partial_{\ell+1}} C^{\ell+1} \xrightarrow{\partial_\ell} C^\ell \xrightarrow{\partial_{\ell-1}} \dots C^1 \xrightarrow{\partial_0} 0$$

and

$$0 \longrightarrow C^1 \xrightarrow{\delta_0} C^2 \xrightarrow{\delta_1} \dots \xrightarrow{\delta_{\ell-1}} C^{\ell+1} \xrightarrow{\delta_\ell} \dots$$

In this setting we will prove

**Theorem 3.** *Assume that  $K$  satisfies the hypotheses of Theorem 2, and is continuous. Then we have the orthogonal (with respect to  $L^2$ ), direct sum decomposition*

$$C^{\ell+1} = \delta(C^\ell) \oplus \partial(C^{\ell+2}) \oplus \text{Ker}_C \Delta$$

where  $\text{Ker}_C \Delta$  denotes the subspace of elements in  $\text{Ker} \Delta$  that are in  $C^{\ell+1}$ .

As in Theorem 2, the third summand is trivial except when  $\ell = 0$  in which case it consists of the constant functions. We first assume that  $K \equiv 1$ . The proof follows from a few propositions. In the remainder of the section,  $\text{Im} \delta$  and  $\text{Im} \partial$  will refer to the image spaces of  $\delta$  and  $\partial$  as operators on  $L_a^2$ . The next proposition gives formulas for  $\partial$  and  $\Delta$  on alternating functions.

**Proposition 7.** For  $f \in L_a^2(X^{\ell+1})$  we have

$$\partial f(x_0, \dots, x_{\ell-1}) = (\ell + 1) \int_X f(t, x_0, \dots, x_{\ell-1}) d\mu(t)$$

and

$$\Delta f(x_0, \dots, x_\ell) = (\ell + 2)f(x_0, \dots, x_\ell) - \frac{1}{\ell + 1} \sum_{i=0}^{\ell} \partial f(x_0, \dots, \hat{x}_i, \dots, x_\ell).$$

*Proof.* The first formula follows immediately from (4) and the fact that  $f$  is alternating. The second follows from a simple calculation using (3), (4) and the fact that  $f$  is alternating.  $\square$

Let  $P_1$ ,  $P_2$ , and  $P_3$  be the orthogonal projections implicit in Theorem 2

$$P_1: L_a^2(X^{\ell+1}) \rightarrow \text{Im } \delta, \quad P_2: L_a^2(X^{\ell+1}) \rightarrow \text{Im } \partial, \quad \text{and } P_3: L_a^2(X^{\ell+1}) \rightarrow \text{Ker } \Delta$$

**Proposition 8.** Let  $f \in C^{\ell+1}$ . Then  $P_1(f) \in C^{\ell+1}$ .

*Proof.* It suffices to show that  $P_1(f)$  is continuous and bounded. Let  $g = P_1(f)$ . It follows from Theorem 2 that  $\partial f = \partial g$ , and therefore  $\partial g$  is continuous and bounded. Since  $\delta g = 0$ , we have, for  $t, x_0, \dots, x_\ell \in X$

$$0 = \delta g(t, x_0, \dots, x_\ell) = g(x_0, \dots, x_\ell) - \sum_{i=0}^{\ell} (-1)^i g(t, x_0, \dots, \hat{x}_i, \dots, x_\ell).$$

Integrating over  $t \in X$  gives us

$$\begin{aligned} g(x_0, \dots, x_\ell) &= \int_X g(x_0, \dots, x_\ell) d\mu(t) = \sum_{i=0}^{\ell} (-1)^i \int_X g(t, x_0, \dots, \hat{x}_i, \dots, x_\ell) d\mu(t) \\ &= \frac{1}{\ell + 1} \sum_{i=0}^{\ell} (-1)^i \partial g(x_0, \dots, \hat{x}_i, \dots, x_\ell). \end{aligned}$$

As  $\partial g$  is continuous and bounded, this implies  $g$  is continuous and bounded.  $\square$

**Corollary 2.** If  $f \in C^{\ell+1}$ , then  $P_2(f) \in C^{\ell+1}$ .

This follows from the Hodge decomposition (Theorem 2) and the fact that  $P_3(f)$  is continuous and bounded (being a constant).

The following proposition can be thought of as analogous to a regularity result in elliptic PDE's. It states that solutions to  $\Delta u = f$ ,  $f$  continuous, which are *a priori* in  $L^2$  are actually continuous.

**Proposition 9.** If  $f \in C^{\ell+1}$  and  $\Delta u = f$ ,  $u \in L_a^2(X^{\ell+1})$  then  $u \in C^{\ell+1}$ .

*Proof.* From Proposition 7, (with  $u$  in place of  $f$ ) we have

$$\begin{aligned}\Delta u(x_0, \dots, x_\ell) &= (\ell + 2)u(x_0, \dots, x_\ell) - \frac{1}{\ell + 1} \sum_{i=0}^{\ell} \partial u(x_0, \dots, \hat{x}_i, \dots, x_\ell) \\ &= f(x_0, \dots, x_\ell)\end{aligned}$$

and solving for  $u$ , we get

$$u(x_0, \dots, x_\ell) = \frac{1}{\ell + 2} f(x_0, \dots, x_\ell) + \frac{1}{(\ell + 2)(\ell + 1)} \sum_{i=0}^{\ell} \partial u(x_0, \dots, \hat{x}_i, \dots, x_\ell).$$

It therefore suffices to show that  $\partial u$  is continuous and bounded. However, it is easy to check that  $\Delta \circ \partial = \partial \circ \Delta$  and thus

$$\Delta(\partial u) = \partial \Delta u = \partial f$$

is continuous and bounded. But then, again using Proposition 7,

$$\begin{aligned}\Delta(\partial u)(x_0, \dots, x_{\ell-1}) &= (\ell + 1)\partial u(x_0, \dots, x_{\ell-1}) \\ &\quad - \frac{1}{\ell} \sum_{i=0}^{\ell-1} (-1)^i \partial(\partial u)(x_0, \dots, \hat{x}_i, \dots, x_{\ell-1})\end{aligned}$$

and so, using  $\partial^2 = 0$  we get

$$(\ell + 1)\partial u = \partial f$$

which implies that  $\partial u$  is continuous and bounded, finishing the proof.  $\square$

**Proposition 10.** *If  $g \in C^{\ell+1} \cap \text{Im } \delta$ , then  $g = \delta h$  for some  $h \in C^\ell$ .*

*Proof.* From the corollary of the Hodge Lemma, let  $h$  be the unique element in  $\text{Im } \partial$  with  $g = \delta h$ . Now  $\partial g$  is continuous and bounded, and

$$\partial g = \partial \delta h = \partial \delta h + \delta \partial h = \Delta h$$

since  $\partial h = 0$ . But now  $h$  is continuous and bounded from Proposition 9.  $\square$

**Proposition 11.** *If  $g \in C^{\ell+1} \cap L_a^2(X^{\ell+1})$ , then  $g = \partial h$  for some  $h \in C^{\ell+2}$ .*

The proof is identical to the one for Proposition 10.

Theorem 3, in the case  $K \equiv 1$  now follows from Propositions 8 through 11. The proof easily extends to general  $K$  which is bounded below by a positive constant.

## 4 Hodge Theory at Scale $\alpha$

As seen in Sections 2 and 3, the chain and cochain complexes constructed on the whole space yield trivial cohomology groups. In order to have a theory that gives us topological information about  $X$ , we define our complexes on a neighborhood of the diagonal, and restrict the boundary and coboundary operator to these complexes. The corresponding cohomology can be considered a cohomology of  $X$  at a scale, with the scale being the size of the neighborhood. We will assume throughout this section that  $(X, d)$  is a compact metric space. For  $x, y \in X^\ell$ ,  $\ell > 1$ , this induces a metric compatible with the product topology

$$d_\ell(x, y) = \max\{d(x_0, y_0), \dots, d(x_{\ell-1}, y_{\ell-1})\}$$

The diagonal  $D_\ell$  of  $X^\ell$  is just  $\{x \in X^\ell : x_i = x_j, i, j = 0, \dots, \ell-1\}$ . For  $\alpha > 0$  we define the  $\alpha$ -neighborhood of the diagonal to be

$$\begin{aligned} U_\alpha^\ell &= \{x \in X^\ell : d_\ell(x, D_\ell) \leq \alpha\} \\ &= \{x \in X^\ell : \exists t \in X \text{ such that } d(x_i, t) \leq \alpha, i = 0, \dots, \ell-1\}. \end{aligned}$$

Observe that  $U_\alpha^\ell$  is closed and that for  $\alpha \geq \text{diameter } X$ ,  $U_\alpha^\ell = X^\ell$ .

The measure  $\mu_\ell$  induces a Borel measure on  $U_\alpha^\ell$  which we will simply denote by  $\mu_\ell$  (not a probability measure). For simplicity, we will take  $K \equiv 1$  throughout this section, and consider only alternating functions in our complexes. We first discuss the  $L^2$ -theory, and thus our basic spaces will be  $L_a^2(U_\alpha^\ell)$ , the space of alternating functions on  $U_\alpha^\ell$  that are in  $L^2$  with respect to  $\mu_\ell$ ,  $\ell > 0$ . Note that if  $(x_0, \dots, x_\ell) \in U_\alpha^{\ell+1}$ , then  $(x_0, \dots, \hat{x}_i, \dots, x_\ell) \in U_\alpha^\ell$  for  $i = 0, \dots, \ell$ . It follows that if  $f \in L_a^2(U_\alpha^\ell)$ , then  $\delta f \in L_a^2(U_\alpha^{\ell+1})$ . We therefore have the well defined cochain complex

$$0 \longrightarrow L_a^2(U_\alpha^1) \xrightarrow{\delta} L_a^2(U_\alpha^2) \cdots \xrightarrow{\delta} L_a^2(U_\alpha^\ell) \xrightarrow{\delta} L_a^2(U_\alpha^{\ell+1}) \cdots$$

Since  $\partial = \delta^*$  depends on the integral, the expression for it will be different from (4). We define a “slice” by

$$S_{x_0 \cdots x_{\ell-1}} = \{t \in X : (x_0, \dots, x_{\ell-1}, t) \in U_\alpha^{\ell+1}\}.$$

We note that, for  $S_{x_0 \cdots x_{\ell-1}}$  to be nonempty,  $(x_0, \dots, x_{\ell-1})$  must be in  $U_\alpha^\ell$ . Furthermore

$$U_\alpha^{\ell+1} = \{(x_0, \dots, x_\ell) : (x_0, \dots, x_{\ell-1}) \in U_\alpha^\ell, \text{ and } x_\ell \in S_{x_0 \cdots x_{\ell-1}}\}.$$

It follows from the proof of Proposition 1 of Section 2 and the fact that  $K \equiv 1$ , that  $\delta : L_a^2(U_\alpha^\ell) \rightarrow L_a^2(U_\alpha^{\ell+1})$  is bounded and that  $\|\delta\| \leq \ell + 1$ , and therefore  $\delta^*$  is bounded. The adjoint of the operator  $\delta : L_a^2(U_\alpha^\ell) \rightarrow L_a^2(U_\alpha^{\ell+1})$  will be denoted, as before, by either  $\partial$  or  $\delta^*$  (without the subscript  $\ell$ ).

**Proposition 12.** *For  $f \in L_a^2(U_\alpha^{\ell+1})$  we have*

$$\partial f(x_0, \dots, x_{\ell-1}) = (\ell + 1) \int_{S_{x_0 \cdots x_{\ell-1}}} f(t, x_0, \dots, x_{\ell-1}) d\mu(t).$$

*Proof.* The proof is essentially the same as the proof of Proposition 4, using the fact that  $K \equiv 1$ ,  $f$  is alternating, and the above remark.  $\square$

It is worth noting that the domain of integration depends on  $x \in U_\alpha^\ell$ , and this makes the subsequent analysis more difficult than in Section 3. We thus have the corresponding chain complex

$$\dots \xrightarrow{\partial} L_a^2(U_\alpha^{\ell+1}) \xrightarrow{\partial} L_a^2(U_\alpha^\ell) \xrightarrow{\partial} \dots L_a^2(U_\alpha^1) \xrightarrow{\partial} 0.$$

Of course,  $U_\alpha^1 = X$ . The corresponding Hodge Laplacian is the operator  $\Delta: L_a^2(U_\alpha^\ell) \rightarrow L_a^2(U_\alpha^\ell)$ ,  $\Delta = \partial\delta + \delta\partial$ , where all of these operators depend on  $\ell$  and  $\alpha$ . When we want to emphasize this dependence, we will list  $\ell$  and (or)  $\alpha$  as subscripts. We will use the following notation for the cohomology and harmonic functions of the above complexes:

$$H_{L^2,\alpha}^\ell(X) = \frac{\text{Ker } \delta_{\ell,\alpha}}{\text{Im } \delta_{\ell-1,\alpha}} \quad \text{and} \quad \text{Harm}_\alpha^\ell(X) = \text{Ker } \Delta_{\ell,\alpha}.$$

**Remark 4.** If  $\alpha \geq \text{diam}(X)$ , then  $U_\alpha^\ell = X^\ell$ , so the situation is as in Theorem 2 of Section 2, so  $H_{L^2,\alpha}^\ell(X) = 0$  for  $\ell > 0$  and  $H_{L^2,\alpha}^0(X) = \mathbb{R}$ . Also, if  $X$  is a finite union of connected components  $X_1, \dots, X_k$ , and  $\alpha < d(X_i, X_j)$  for all  $i \neq j$ , then  $H_{L^2,\alpha}^\ell(X) = \oplus_{i=1}^k H_{L^2,\alpha}^\ell(X_i)$ .

**Definition 1.** We say that Hodge theory for  $X$  at scale  $\alpha$  holds if we have the orthogonal direct sum decomposition into closed subspaces

$$L_a^2(U_\alpha^\ell) = \text{Im } \delta_{\ell-1} \oplus \text{Im } \delta_\ell^* \oplus \text{Harm}_\alpha^\ell(X) \quad \text{for all } \ell$$

and furthermore,  $H_{\alpha,L^2}^\ell(X)$  is isomorphic to  $\text{Harm}_\alpha^\ell(X)$ , with each equivalence class in the former having a unique representative in the latter.

**Theorem 4.** If  $X$  is a compact metric space,  $\alpha > 0$ , and the  $L^2$ -cohomology spaces  $\text{Ker } \delta_{\ell,\alpha} / \text{Im } \delta_{\ell-1,\alpha}$ ,  $\ell \geq 0$  are finite dimensional, then Hodge theory for  $X$  at scale  $\alpha$  holds.

*Proof.* This is immediate from the Hodge Lemma (Lemma 1), using Proposition 5 from Section 2.  $\square$

We record the formulas for  $\delta\partial f$  and  $\partial\delta f$  for  $f \in L_a^2(U_\alpha^{\ell+1})$

$$\begin{aligned} \delta(\partial f)(x_0, \dots, x_\ell) &= (\ell+1) \sum_{i=0}^{\ell} (-1)^i \int_{S_{x_0, \dots, \hat{x}_i, \dots, x_\ell}} f(t, x_0, \dots, \hat{x}_i, \dots, x_\ell) d\mu(t) \\ \partial(\delta f)(x_0, \dots, x_\ell) &= (\ell+2) \mu(S_{x_0, \dots, x_\ell}) f(x_0, \dots, x_\ell) \\ &\quad + (\ell+2) \sum_{i=0}^{\ell} (-1)^{i+1} \int_{S_{x_0, \dots, x_\ell}} f(t, x_0, \dots, \hat{x}_i, \dots, x_\ell) d\mu(t) \end{aligned} \quad (7)$$

Of course, the formula for  $\Delta f$  is found by adding these two.



**Remark 5.** Harmonic forms are solutions of the optimization problem: Minimize the “Dirichlet norm”  $\|\delta f\|^2 + \|\partial f\|^2 = \langle \Delta f, f \rangle = \langle \Delta^{1/2} f, \Delta^{1/2} f \rangle$  over  $f \in L_a^2(U_\alpha^{\ell+1})$ .

**Remark 6.** There is a second notion of  $U_\alpha^{\ell+1}$  called the Rips complex (see Chazal and Oudot [7]) defined by  $(x_0, \dots, x_\ell) \in U_\alpha^{\ell+1}(\text{Rips})$  if and only if  $d(x_i, x_j) \leq \alpha$  for all  $i, j$ . This corresponds to the theory developed in Section 2 with  $K(x, y)$  equal to the characteristic function of  $U_\alpha^2(\text{Rips})$ . A version of Theorem 4 holds in this case.

## 5 $L^2$ -Theory of $\alpha$ -Harmonic 0-Forms

In this section we assume that we are in the setting of Section 4, with  $\ell = 0$ . Thus  $X$  is a compact metric space with a probability measure and with a fixed scale  $\alpha > 0$ .

Recall that  $f \in L^2(X)$  is  $\alpha$ -harmonic if  $\Delta_\alpha f = 0$ . Moreover, if  $\delta: L^2(X) \rightarrow L_a^2(U_\alpha^2)$  denotes the coboundary, then  $\Delta_\alpha f = 0$  if and only if  $\delta f = 0$ ; also  $\delta f(x_0, x_1) = f(x_1) - f(x_0)$  for all pairs  $(x_0, x_1) \in U_\alpha^2$ .

Recall that for any  $x \in X$ , the slice  $S_{x,\alpha} = S_x \subset X^2$  is the set

$$S_x = S_{x,\alpha} = \{t \in X : \exists p \in X \text{ such that } x, t \in B_\alpha(p)\}.$$

Note that  $B_\alpha(x) \subset S_{x,\alpha} \subset B_{2\alpha}(x)$ . It follows that  $x_1 \in S_{x_0,\alpha}$  if and only if  $x_0 \in S_{x_1,\alpha}$ . We conclude

**Proposition 13.** *Let  $f \in L^2(X)$ . Then  $\Delta_\alpha f = 0$  if and only if  $f$  is locally constant in the sense that  $f$  is constant on  $S_{x,\alpha}$  for every  $x \in X$ . Moreover if  $\Delta_\alpha f = 0$ , then*

- (a) *If  $X$  is connected, then  $f$  is constant.*
- (b) *If  $\alpha$  is greater than the maximum distance between components of  $X$ , then  $f$  is constant.*
- (c) *For any  $x \in X$ ,  $f(x)$  = average of  $f$  on  $S_{x,\alpha}$  and on  $B_\alpha(x)$ .*
- (d) *Harmonic functions are continuous.*

We note that continuity of  $f$  follows from the fact that  $f$  is constant on each slice  $S_{x,\alpha}$ , and thus locally constant.

**Remark 7.** We will show that (d) is also true for harmonic 1-forms with an additional assumption on  $\mu$ , (Section 8) but are unable to prove it for harmonic 2-forms.

Consider next an extension of (d) to the Poisson regularity problem. If  $\Delta_\alpha f = g$  is continuous, is  $f$  continuous? In general the answer is no, and we will give an example.

Since  $\partial_0$  on  $L^2(X)$  is zero, the  $L^2$ - $\alpha$ -Hodge theory (Section 9) takes the form

$$L^2(X) = \text{Im } \partial \oplus \text{Harm}_\alpha,$$

where  $\partial: L^2(U_\alpha^2) \rightarrow L^2(X)$  and  $\Delta f = \partial \delta f$ . Thus for  $f \in L^2(X)$ , by (7)

$$\Delta_\alpha f(x) = 2\mu(S_{x,\alpha})f(x) - 2 \int_{S_{x,\alpha}} f(t) d\mu(t) \quad (8)$$

The following example shows that an additional assumption is needed for the Poisson regularity problem to have an affirmative solution. Let  $X$  be the closed interval  $[-1, 1]$  with the usual metric  $d$  and let  $\mu$  be the Lebesgue measure on  $X$  with an atom at 0,  $\mu(\{0\}) = 1$ . Fix any  $\alpha < 1/4$ . We will define a piecewise linear function on  $X$  with discontinuities at  $-2\alpha$  and  $2\alpha$  as follows. Let  $a$  and  $b$  be any real numbers  $a \neq b$ , and define

$$f(x) = \begin{cases} \frac{a-b}{8\alpha} + a, & -1 \leq x < -2\alpha \\ \frac{b-a}{4\alpha}(x - 2\alpha) + b, & -2\alpha \leq x \leq 2\alpha \\ \frac{a-b}{8\alpha} + b, & 2\alpha < x \leq 1. \end{cases}$$

Using (8) above one readily checks that  $\Delta_\alpha f$  is continuous by computing left hand and right hand limits at  $\pm 2\alpha$ . (The constant values of  $f$  outside  $[-2\alpha, 2\alpha]$  are chosen precisely so that the discontinuities of the two terms on the right side of (8) cancel out.)

With an additional “regularity” hypothesis imposed on  $\mu$ , the Poisson regularity property holds. In the rest of this section assume that  $\mu(S_x \cap A)$  is a continuous function of  $x \in X$  for each measurable set  $A$ . One can show that if  $\mu$  is Borel regular, then this will hold provided  $\mu(S_x \cap A)$  is continuous for all closed sets  $A$  (or all open sets  $A$ ).

**Proposition 14.** *Assume that  $\mu(S_x \cap A)$  is a continuous function of  $x \in X$  for each measurable set  $A$ . If  $\Delta_\alpha f = g$  is continuous for  $f \in L^2(X)$  then  $f$  is continuous.*

*Proof.* From (8) we have

$$f(x) = \frac{g(x)}{2\mu(S_x)} + \frac{1}{\mu(S_x)} \int_{S_x} f(t) d\mu(t)$$

The first term on the right is clearly continuous by our hypotheses on  $\mu$  and the fact that  $g$  is continuous. It suffices to show that the function  $h(x) = \int_{S_x} f(t) d\mu(t)$  is continuous. If  $f = \chi_A$  is the characteristic function of any measurable set  $A$ , then  $h(x) = \mu(S_x \cap A)$  is continuous, and therefore  $h$  is continuous for  $f$  any simple function (linear combination of characteristic functions of measurable sets). From general measure theory, if  $f \in L^2(X)$ , we can find a sequence of simple functions  $f_n$  such that  $f_n(t) \rightarrow f(t)$  a.e. and  $|f_n(t)| \leq |f(t)|$  for all  $t \in X$ . Thus  $h_n(x) = \int_{S_x} f_n(t) d\mu(t)$  is continuous and

$$|h_n(x) - h(x)| \leq \int_{S_x} |f_n(t) - f(t)| d\mu(t) \leq \int_X |f_n(t) - f(t)| d\mu(t)$$

Since  $|f_n - f| \rightarrow 0$  a.e. and  $|f_n - f| \leq 2|f|$  with  $f$  being in  $L^1(X)$ , it follows from the dominated convergence theorem that  $\int_X |f_n - f| d\mu \rightarrow 0$ . Thus  $h_n$  converges uniformly to  $h$  and so continuity of  $h$  follows from continuity of  $h_n$ .  $\square$

We don't have a similar result for 1-forms.

Partly to relate our framework of  $\alpha$ -harmonic theory to some previous work, we combine the setting of Section 2 with Section 4. Thus we now put back the function  $K$ . Assume  $K > 0$  is a symmetric and continuous function  $K: X \times X \rightarrow \mathbb{R}$ , and  $\delta$  and  $\partial$  are defined as in Section 2, but use a similar extension to general  $\alpha > 0$ , of Section 4, all in the  $L^2$ -theory.

Let  $D: L^2(X) \rightarrow L^2(X)$  be the operator defined as multiplication by the function

$$D(x) = \int_X G(x, y) d\mu(y) \quad \text{where } G(x, y) = K(x, y)\chi_{U_\alpha^2}$$

using the characteristic function  $\chi_{U_\alpha^2}$  of  $U_\alpha^2$ . So  $\chi_{U_\alpha^2}(x_0, x_1) = 1$  if  $(x_0, x_1) \in U_\alpha^2$  and 0 otherwise. Furthermore, let  $L_G: L^2(X) \rightarrow L^2(X)$  be the integral operator defined by

$$L_G f(x) = \int_X G(x, y) f(y) d\mu(y).$$

Note that  $L_G(1) = D$  where 1 is the constant function. When  $X$  is compact  $L_G$  is a Hilbert-Schmidt operator (this was first noted to us by Ding-Xuan Zhou). Thus  $L_G$  is trace class and self adjoint. It is not difficult to see now that (8) takes the form

$$\frac{1}{2}\Delta_\alpha f = Df - L_G f. \quad (9)$$

(For the special case  $\alpha = \infty$ , i.e.  $\alpha$  is irrelevant as in Section 2, this is the situation as in Smale and Zhou [33] for the case  $K$  is a reproducing kernel.) As in the previous proposition:

**Proposition 15.** *The Poisson Regularity Property holds for the operator of (9).*

To get a better understanding of (9) it is useful to define a normalization of the kernel  $G$  and the operator  $L_G$  as follows. Let  $\hat{G}: X \times X \rightarrow \mathbb{R}$  be defined by

$$\hat{G}(x, y) = \frac{G(x, y)}{(D(x)D(y))^{1/2}}$$

and  $L_{\hat{G}}: L^2(X) \rightarrow L^2(X)$  be the corresponding integral operator. Then  $L_{\hat{G}}$  is trace class, self adjoint, with non-negative eigenvalues, and has a complete orthonormal system of continuous eigenfunctions.

A normalized  $\alpha$ -Laplacian may be defined on  $L^2(X)$  by

$$\frac{1}{2}\hat{\Delta} = I - L_{\hat{G}}$$

so that the spectral theory of  $L_{\hat{G}}$  may be transferred to  $\hat{\Delta}$ . (Also, one might consider  $\frac{1}{2}\Delta^* = I - D^{-1}L_G$  as in Belkin, De Vito, and Rosasco [3].)

In Smale and Zhou [33], for  $\alpha = \infty$ , error estimates are given (reproducing kernel case) for the spectral theory of  $L_{\hat{G}}$  in terms of finite dimensional approximations. See especially Belkin and Niyogi [2] for limit theorems as  $\alpha \rightarrow 0$ .

## 6 Harmonic forms on constant curvature manifolds

In this section we will give an explicit description of harmonic forms in a special case. Let  $X$  be a compact, connected, oriented manifold of dimension  $n > 0$ , with a Riemannian metric  $g$  of constant sectional curvature. Also, assume that  $g$  is normalized so that  $\mu(X) = 1$  where  $\mu$  is the measure induced by the volume form associated with  $g$ , and let  $d$  be the metric on  $X$  induced by  $g$ . Let  $\alpha > 0$  be sufficiently small so that for all  $p \in X$ , the ball  $B_{2\alpha}(p)$  is geodesically convex. That is, for  $x, y \in B_{2\alpha}(p)$  there is a unique, length minimizing geodesic  $\gamma$  from  $x$  to  $y$ , and  $\gamma$  lies in  $B_{2\alpha}(p)$ . Note that if  $(x_0, \dots, x_n) \in U_{\alpha}^{n+1}$ , then  $d(x_i, x_j) \leq 2\alpha$  for all  $i, j$ , and thus all  $x_i$  lie in a common geodesically convex ball. Such a point defines an  $n$ -simplex with vertices  $x_0, \dots, x_n$  whose faces are totally geodesic submanifolds, which we will denote by  $\sigma(x_0, \dots, x_n)$ . We will also denote the  $k$ -dimensional faces by  $\sigma(x_{i_0}, \dots, x_{i_k})$  for  $k < n$ . Thus  $\sigma(x_i, x_j)$  is the geodesic segment from  $x_i$  to  $x_j$ ,  $\sigma(x_i, x_j, x_k)$  is the union of geodesic segments from  $x_i$  to points on  $\sigma(x_j, x_k)$  and higher dimensional simplices are defined inductively. (Since  $X$  has constant curvature, this construction is symmetric in  $x_0, \dots, x_n$ .) A  $k$ -dimensional face will be called degenerate if one of its vertices is contained in one of its  $(k-1)$ -dimensional faces.

For  $(x_0, \dots, x_n) \in U_{\alpha}^{n+1}$ , the orientation on  $X$  induces an orientation on  $\sigma(x_0, \dots, x_n)$  (assuming it is non-degenerate). For example, if  $v_1, \dots, v_n$  denote the tangent vectors at  $x_0$  to the geodesics from  $x_0$  to  $x_1, \dots, x_n$ , we can define  $\sigma(x_0, \dots, x_n)$  to be positive (negative) if  $\{v_1, \dots, v_n\}$  is a positive (respectively negative) basis for the tangent space at  $x_0$ . Of course, if  $\tau$  is a permutation, the orientation of  $\sigma(x_0, \dots, x_n)$  is equal to  $(-1)^{\text{sign } \tau}$  times the orientation of  $\sigma(x_{\tau(0)}, \dots, x_{\tau(n)})$ . We now define  $f: U_{\alpha}^{n+1} \rightarrow \mathbb{R}$  by

$$\begin{aligned} f(x_0, \dots, x_n) &= \mu(\sigma(x_0, \dots, x_n)) \quad \text{for } \sigma(x_0, \dots, x_n) \text{ positive} \\ &= -\mu(\sigma(x_0, \dots, x_n)) \quad \text{for } \sigma(x_0, \dots, x_n) \text{ negative} \\ &= 0 \quad \text{for } \sigma(x_0, \dots, x_n) \text{ degenerate.} \end{aligned}$$

Thus  $f$  is the signed volume of oriented geodesic  $n$ -simplices. Clearly  $f$  is continuous as non-degeneracy is an open condition and the volume of a simplex varies continuously in the vertices. The main result of this section is

**Theorem 5.** *Let  $X$  be a oriented Riemannian  $n$ -manifold of constant sectional curvature and  $f$ ,  $\alpha$  as above. Then  $f$  is harmonic. In fact  $f$  is the unique harmonic  $n$ -form in  $L_a^2(U_{\alpha}^{n+1})$  up to scaling.*

*Proof.* Uniqueness follows from Section 9. We will show that  $\partial f = 0$  and  $\delta f = 0$ . Let  $(x_0, \dots, x_{n-1}) \in U_\alpha^n$ . To show  $\partial f = 0$ , it suffices to show, by Proposition 12, that

$$\int_{S_{x_0 \dots x_{n-1}}} f(t, x_0, \dots, x_{n-1}) d\mu(t) = 0. \quad (10)$$

We may assume that  $\sigma(x_0, \dots, x_{n-1})$  is non-degenerate, otherwise the integrand is identically zero. Recall that  $S_{x_0 \dots x_{n-1}} = \{t \in X : (t, x_0, \dots, x_{n-1}) \subset U_\alpha^{n+1}\} \subset B_{2\alpha}(x_0)$  where  $B_{2\alpha}(x_0)$  is the geodesic ball of radius  $2\alpha$  centered at  $x_0$ . Let  $\Gamma$  be the intersection of the totally geodesic  $n-1$  dimensional submanifold containing  $x_0, \dots, x_{n-1}$  with  $B_{2\alpha}(x_0)$ . Thus  $\Gamma$  divides  $B_{2\alpha}(x_0)$  into two pieces  $B^+$  and  $B^-$ . For  $t \in \Gamma$ , the simplex  $\sigma(t, x_0, \dots, x_{n-1})$  is degenerate and therefore the orientation is constant on each of  $B^+$  and  $B^-$ , and we can assume that the orientation of  $\sigma(t, x_0, \dots, x_{n-1})$  is positive on  $B^+$  and negative on  $B^-$ . For  $x \in B_{2\alpha}(x_0)$  define  $\phi(x)$  to be the reflection of  $x$  across  $\Gamma$ . Thus the geodesic segment from  $x$  to  $\phi(x)$  intersects  $\Gamma$  perpendicularly at its midpoint. Because  $X$  has constant curvature,  $\phi$  is a local isometry and since  $x_0 \in \Gamma$ ,  $d(x, x_0) = d(\phi(x), x_0)$ . Therefore  $\phi: B_{2\alpha}(x_0) \rightarrow B_{2\alpha}(x_0)$  is an isometry which maps  $B^+$  isometrically onto  $B^-$  and  $B^-$  onto  $B^+$ . Denote  $S_{x_0 \dots x_{n-1}}$  by  $S$ . It is easy to see that  $\phi: S \rightarrow S$ , and so defining  $S^\pm = S \cap B^\pm$  it follows that  $\phi: S^+ \rightarrow S^-$  and  $\phi: S^- \rightarrow S^+$  are isometries. Now

$$\begin{aligned} & \int_{S_{x_0 \dots x_{n-1}}} f(t, x_0, \dots, x_{n-1}) d\mu(t) \\ &= \int_{S^+} f(t, x_0, \dots, x_{n-1}) d\mu(t) + \int_{S^-} f(t, x_0, \dots, x_{n-1}) d\mu(t) \\ &= \int_{S^+} \mu(\sigma(t, x_0, \dots, x_{n-1})) d\mu(t) - \int_{S^-} \mu(\sigma(t, x_0, \dots, x_{n-1})) d\mu(t). \end{aligned}$$

Since  $\mu(\sigma(t, x_0, \dots, x_{n-1})) = \mu(\sigma(\phi(t), x_0, \dots, x_{n-1}))$  for  $t \in S^+$ , the last two terms on the right side cancel establishing (10).

We now show that  $\delta f = 0$ . Let  $(t, x_0, \dots, x_n) \in U_\alpha^{n+2}$ . Thus

$$\delta f(t, x_0, \dots, x_n) = f(x_0, \dots, x_n) + \sum_{i=0}^n (-1)^{i+1} f(t, x_0, \dots, \hat{x}_i, \dots, x_n)$$

and we must show that

$$f(x_0, \dots, x_n) = \sum_{i=0}^n (-1)^i f(t, x_0, \dots, \hat{x}_i, \dots, x_n). \quad (11)$$

Without loss of generality, we will assume that  $\sigma(x_0, \dots, x_n)$  is positive. The demonstration of (11) depends on the location of  $t$ . Suppose that  $t$  is in the interior of the simplex  $\sigma(x_0, \dots, x_n)$ . Then for each  $i$ , the orientation of  $\sigma(x_0, \dots, x_{i-1}, t, x_{i+1}, \dots, x_n)$  is the same as the orientation of  $\sigma(x_0, \dots, x_n)$

since  $t$  and  $x_i$  lie on the same side of the face  $\sigma(x_0, \dots, \hat{x}_i, \dots, x_n)$ , and is thus positive. On the other hand, the orientation of  $\sigma(t, x_0, \dots, \hat{x}_i, \dots, x_n)$  is  $(-1)^i$  times the orientation of  $\sigma(x_0, \dots, x_{i-1}, t, x_{i+1}, \dots, x_n)$ . Therefore the right side of (11) becomes

$$\sum_{i=0}^n \mu(\sigma(x_0, \dots, x_{i-1}, t, x_{i+1}, \dots, x_n)).$$

This however equals  $\mu(\sigma(x_0, \dots, x_n))$  which is the left side of (11), since

$$\sigma(x_0, \dots, x_n) = \bigcup_{i=0}^n \sigma(x_0, \dots, x_{i-1}, t, x_{i+1}, \dots, x_n)$$

when  $t$  is interior to  $\sigma(x_0, \dots, x_n)$ .

There are several cases when  $t$  is exterior to  $\sigma(x_0, \dots, x_n)$  (or on one of the faces), depending on which side of the various faces it lies. We just give the details of one of these, the others being similar. Simplifying notation, let  $F_i$  denote the face “opposite”  $x_i$ ,  $\sigma(x_0, \dots, \hat{x}_i, \dots, x_n)$ , and suppose that  $t$  is on the opposite side of  $F_0$  from  $x_0$ , but on the same side of  $F_i$  as  $x_i$  for  $i \neq 0$ . As in the above argument, the orientation of  $\sigma(x_0, \dots, x_{i-1}, t, x_{i+1}, \dots, x_n)$  is positive for  $i \neq 0$  and is negative for  $i = 0$ . Therefore the right side of (11) is equal to

$$\sum_{i=1}^n \mu(\sigma(x_0, \dots, x_{i-1}, t, x_{i+1}, \dots, x_n)) - \mu(\sigma(t, x_1, \dots, x_n)). \quad (12)$$

Let  $s$  be the point where the geodesic from  $x_0$  to  $t$  intersects  $F_0$ . Then for each  $i > 0$

$$\begin{aligned} \sigma(x_0, \dots, x_{i-1}, t, x_{i+1}, \dots, x_n) &= \sigma(x_0, \dots, x_{i-1}, s, x_{i+1}, \dots, x_n) \\ &\quad \cup \sigma(s, \dots, x_{i-1}, t, x_{i+1}, \dots, x_n). \end{aligned}$$

Taking  $\mu$  of both sides and summing over  $i$  gives

$$\begin{aligned} \sum_{i=1}^n \mu(\sigma(x_0, \dots, x_{i-1}, t, x_{i+1}, \dots, x_n)) &= \sum_{i=1}^n \mu(\sigma(x_0, \dots, x_{i-1}, s, x_{i+1}, \dots, x_n)) \\ &\quad + \sum_{i=1}^n \mu(\sigma(s, \dots, x_{i-1}, t, x_{i+1}, \dots, x_n)). \end{aligned}$$

However, the first term on the right is just  $\mu(\sigma(x_0, \dots, x_n))$  and the second term is  $\mu(\sigma(t, x_1, \dots, x_n))$ . Combining this with (12) gives us (11), finishing the proof of  $\delta f = 0$ .  $\square$

**Remark 8.** The proof that  $\partial f = 0$  strongly used the fact that  $X$  has constant curvature. In the case where  $X$  is an oriented Riemannian surface of variable

curvature, totally geodesic  $n$  simplices don't generally exist, although geodesic triangles  $\sigma(x_0, x_1, x_2)$  are well defined for  $(x_0, x_1, x_2) \in U_\alpha^3$ . In this case, the proof above shows that  $\delta f = 0$ . More generally, for an  $n$ -dimensional connected oriented Riemannian manifold, using the order of a tuple  $(x_0, \dots, x_n)$  one can iteratively form convex combinations and in this way assign an oriented  $n$ -simplex to  $(x_0, \dots, x_n)$  and then define the volume cocycle as above (if  $\alpha$  is small enough).

Using a chain map to simplicial cohomology which evaluates at the vertices' points, it is easy to check that these cocycles represent a generator of the cohomology in degree  $n$  (which by the results of Section 9 is exactly 1-dimensional).

## 7 Cohomology

Traditional cohomology theories on general spaces are typically defined in terms of limits as in Čech theory, with nerves of coverings. However, an algorithmic approach suggests a development via a scaled theory, at a given scale  $\alpha > 0$ . Then, as  $\alpha \rightarrow 0$  one recovers the classical setting. A closely related point of view is that of persistent homology, see Edelsbrunner, Letscher, and Zomorodian [17], Zomorodian and Carlsson [38], and Carlsson [6].

We give a setting for such a scaled theory, with a fixed scaling parameter  $\alpha > 0$ .

Let  $X$  be a separable, complete metric space with metric  $d$ , and  $\alpha > 0$  a "scale". We will define a (generally infinite) simplicial complex  $C_{X,\alpha}$  associated to  $(X, d, \alpha)$ . Toward that end let  $X^{\ell+1}$ , for  $\ell \geq 0$ , be the  $(\ell + 1)$ -fold Cartesian product, with metric still denoted by  $d$ ,  $d: X^{\ell+1} \times X^{\ell+1} \rightarrow \mathbb{R}$  where  $d(x, y) = \max_{i=0, \dots, \ell} d(x_i, y_i)$ . As in Section 4, let

$$U_\alpha^{\ell+1}(X) = U_\alpha^{\ell+1} = \{x \in X^{\ell+1} : d(x, D_{\ell+1}) \leq \alpha\}$$

where  $D_{\ell+1} \subset X^{\ell+1}$  is the diagonal, so  $D_{\ell+1} = \{(t, \dots, t) \text{ } \ell + 1 \text{ times}\}$ . Then let  $C_{X,\alpha}^\ell = U_\alpha^{\ell+1}$ . This has the structure of a simplicial complex whose  $\ell$ -simplices consist of points of  $U_\alpha^{\ell+1}$ . This is well defined since if  $x \in U_\alpha^{\ell+1}$ , then  $y = (x_0, \dots, \hat{x}_i, \dots, x_\ell) \in U_\alpha^\ell$ , for each  $i = 0, \dots, \ell$ . We will write  $\alpha = \infty$  to mean that  $U_\alpha^\ell = X^\ell$ . Following e.g. Munkres [29], there is a well-defined cohomology theory, simplicial cohomology, for this simplicial complex, with cohomology vector spaces (always over  $\mathbb{R}$ ), denoted by  $H_\alpha^\ell(X)$ . We especially note that  $C_{X,\alpha}$  is not necessarily a finite simplicial complex. For example, if  $X$  is an open non-empty subset of Euclidean space, the vertices of  $C_{X,\alpha}$  are the points of  $X$  and of course infinite in number. The complex  $C_{X,\alpha}$  will be called the simplicial complex at scale  $\alpha$  associated to  $X$ .

**Example 1.**  $X$  is finite. Fix  $\alpha > 0$ . In this case, for each  $\ell$ , the set of  $\ell$ -simplices is finite, the  $\ell$ -chains form a finite dimensional vector space and the  $\alpha$ -cohomology groups (i.e. vector spaces)  $H_\alpha^\ell(X)$  are all finite dimensional. One can check that for  $\alpha = \infty$ , one has  $\dim H_\alpha^0(X) = 1$  and  $H_\alpha^i(X)$  are trivial for all  $i > 0$ . Moreover, for  $\alpha$  sufficiently small ( $\alpha < \min\{d(x, y) : x, y \in X, x \neq y\}$ )

$\dim H_\alpha^0(X) = \text{cardinality of } X$ , with  $H_\alpha^i(X) = 0$  for all  $i > 0$ . For intermediate  $\alpha$ , the  $\alpha$ -cohomology can be rich in higher dimensions, but  $C_{X,\alpha}$  is a finite simplicial complex.

**Example 2.** First let  $A \subset \mathbb{R}^2$  be the annulus  $A = \{x \in \mathbb{R}^2 : 1 \leq \|x\| \leq 2\}$ . Form  $A^*$  by deleting the finite set of points with rational coordinates  $(p/q, r/s)$ , with  $|q|, |s| \leq 10^{10}$ . Then one may check that for  $\alpha > 4$ ,  $H_\alpha^\ell(A^*)$  has the cohomology of a point, for certain intermediate values of  $\alpha$ ,  $H_\alpha^\ell(A^*) = H_\alpha^\ell(A)$ , and for  $\alpha$  small enough  $H_\alpha^\ell(A^*)$  has enormous dimension. Thus the scale is crucial to see the features of  $A^*$  clearly.

Returning to the case of general  $X$ , note that if  $0 < \beta < \alpha$  one has a natural inclusion  $J: U_\beta^\ell \rightarrow U_\alpha^\ell$ ,  $J: C_{X,\beta} \rightarrow C_{X,\alpha}$  and the restriction  $J^*: L_a^2(U_\alpha^\ell) \rightarrow L_a^2(U_\beta^\ell)$  commuting with  $\delta$  (a chain map).

Now assume  $X$  is compact. For fixed scale  $\alpha$ , consider the covering  $\{B_\alpha(x) : x \in X\}$ , where  $B_\alpha(x)$  is the ball  $B_\alpha(x) = \{y \in X : d(x, y) < \alpha\}$ , and the nerve of the covering is  $C_{X,\alpha}$ , giving the ‘‘Čech construction at scale  $\alpha$ ’’. Thus from Čech cohomology theory, we see that the limit as  $\alpha \rightarrow 0$  of  $H_\alpha^\ell(X) = H^\ell(X) = H_{\text{Čech}}^\ell(X)$  is the  $\ell$ -th Čech cohomology group of  $X$ .

The next observation is to note that our construction of the scaled simplicial complex  $C_{X,\alpha}$  of  $X$  follows the same path as Alexander-Spanier theory (see Spanier [34]). Thus the scaled cohomology groups  $H_\alpha^\ell(X)$  will have the direct limit as  $\alpha \rightarrow 0$  which maps to the Alexander-Spanier group  $H_{\text{Alex-Sp}}^\ell(X)$  (and in many cases will be isomorphic). Thus  $H^\ell(X) = H_{\text{Alex-Sp}}^\ell(X) = H_{\text{Čech}}^\ell(X)$ . In fact in much of the literature this is recognized by the use of the term Alexander-Spanier-Čech cohomology. What we have done is describe a finite scale version of the classical cohomology.

Now that we have defined the scale  $\alpha$  cohomology groups,  $H_\alpha^\ell(X)$  for a metric space  $X$ , our Hodge theory suggests this modification. From Theorem 4, we have considered instead of arbitrary cochains (i.e. arbitrary functions on  $U_\alpha^{\ell+1}$  which give the definition here of  $H_\alpha^\ell(X)$ ), cochains defined by  $L^2$ -functions on  $U_\alpha^{\ell+1}$ . Thus we have constructed cohomology groups at scale  $\alpha$  from  $L^2$ -functions on  $U_\alpha^{\ell+1}$ ,  $H_{\alpha,L^2}^\ell(X)$ , when  $\alpha > 0$ , and  $X$  is a metric space equipped with Borel probability measure.

**Question 1** (Cohomology Identification Problem (CIP)). To what extent are  $H_{L^2,\alpha}^\ell(X)$  and  $H_\alpha^\ell(X)$  isomorphic?

This is important via Theorem 4 which asserts that  $H_{\alpha,L^2}^\ell(X) \rightarrow \text{Harm}_\alpha^\ell(X)$  is an isomorphism, in case  $H_{\alpha,L^2}^\ell(X)$  is finite dimensional.

One may replace  $L^2$ -functions in the construction of the  $\alpha$ -scale cohomology theory by continuous functions. As in the  $L^2$ -theory, this gives rise to cohomology groups  $H_{\alpha,\text{cont}}^\ell(X)$ . Analogous to CIP we have the simpler question: To what extent is the natural map  $H_{\alpha,\text{cont}}^\ell(X) \rightarrow H_\alpha^\ell(X)$  an isomorphism?

We will give answers to these questions for special  $X$  in Section 9.

Note that in the case  $X$  is finite, or  $\alpha = \infty$ , we have an affirmative answer to this question, as well as CIP (see Sections 2 and 3).



**Proposition 16.** *There is a natural injective linear map*

$$\text{Harm}_{cont,\alpha}^\ell(X) \rightarrow H_{cont,\alpha}^\ell(X).$$

*Proof.* The inclusion, which is injective

$$J: \text{Im}_{cont,\alpha} \delta \oplus \text{Harm}_{cont,\alpha}^\ell(X) \rightarrow \text{Ker}_{cont,\alpha}$$

induces an injection

$$J^*: \text{Harm}_{cont,\alpha}^\ell(X) = \frac{\text{Im}_{cont,\alpha} \delta \oplus \text{Harm}_{cont,\alpha}^\ell(X)}{\text{Im}_{cont,\alpha} \delta} \rightarrow \frac{\text{Ker}_{cont,\alpha}}{\text{Im}_{cont,\alpha}} = H_{cont,\alpha}^\ell(X)$$

and the proposition follows.  $\square$

## 8 Continuous Hodge theory on the neighborhood of the diagonal

As in the last section,  $(X, d)$  will denote a compact metric space equipped with a Borel probability measure  $\mu$ . For topological reasons (see Section 6) it would be nice to have a Hodge decomposition for continuous functions on  $U_\alpha^{\ell+1}$ , analogous to the continuous theory on the whole space (Section 4). We will use the following notation.  $C_\alpha^{\ell+1}$  will denote the continuous alternating real valued functions on  $U_\alpha^{\ell+1}$ ,  $\text{Ker}_{\alpha,cont} \Delta_\ell$  will denote the functions in  $C_\alpha^{\ell+1}$  that are harmonic, and  $\text{Ker}_{\alpha,cont} \delta_\ell$  will denote those elements of  $C_\alpha^{\ell+1}$  that are closed. Also,  $H_{\alpha,cont}^\ell(X)$  will denote the quotient space (cohomology space)  $\text{Ker}_{\alpha,cont} \delta_\ell / \delta(C_\alpha^\ell)$ . We raise the following question, analogous to Theorem 4.

**Question 2** (Continuous Hodge Decomposition). Under what conditions on  $X$  and  $\alpha > 0$  is it true that there is the following orthogonal (with respect to the  $L^2$ -inner product) direct sum decomposition

$$C_\alpha^{\ell+1} = \delta(C_\alpha^\ell) \oplus \partial(C_\alpha^{\ell+2}) \oplus \text{Ker}_{\alpha,cont} \Delta_\ell$$

where  $\text{Ker}_{cont,\alpha} \Delta_\ell$  is isomorphic to  $H_{\alpha,cont}^\ell(X)$ , with every element in  $H_{\alpha,cont}^\ell(X)$  having a unique representative in  $\text{Ker}_{\alpha,cont} \Delta_\ell$ ?

There is a related analytical problem that is analogous to elliptic regularity for partial differential equations, and in fact elliptic regularity features prominently in classical Hodge theory.

**Question 3** (The Poisson Regularity Problem). For  $\alpha > 0$ , and  $\ell > 0$ , suppose that  $\Delta f = g$  where  $g \in C_\alpha^{\ell+1}$  and  $f \in L_a^2(U_\alpha^{\ell+1})$ . Under what conditions on  $(X, d, \mu)$  is  $f$  continuous?

**Theorem 6.** *An affirmative answer to the Poisson Regularity problem, together with closed image  $\delta(L_a^2(U_\alpha^\ell))$  implies an affirmative solution to the continuous Hodge decomposition question.*

*Proof.* Assume that the Poisson regularity property holds, and let  $f \in C_\alpha^{\ell+1}$ . From Theorem 4 we have the  $L^2$ -Hodge decomposition

$$f = \delta f_1 + \partial f_2 + f_3$$

where  $f_1 \in L_a^2(U_\alpha^\ell)$ ,  $f_2 \in L_a^2(U_\alpha^{\ell+2})$  and  $f_3 \in L_a^2(U_\alpha^{\ell+1})$  with  $\Delta f_3 = 0$ . It suffices to show that  $f_1$  and  $f_2$  can be taken to be continuous, and  $f_3$  is continuous. Since  $\Delta f_3 = 0$  is continuous,  $f_3$  is continuous by Poisson regularity. We will show that  $\partial f_2 = \partial(\delta h_2)$  where  $\delta h_2$  is continuous (and thus  $f_2$  can be taken to be continuous). Recall (corollary of the Hodge Lemma in Section 2) that the following maps are isomorphisms

$$\delta: \partial(L_a^2(U_\alpha^{\ell+2})) \rightarrow \delta(L_a^2(U_\alpha^{\ell+1})) \text{ and } \partial: \delta(L_a^2(U_\alpha^\ell)) \rightarrow \partial(L_a^2(U_\alpha^{\ell+1}))$$

for all  $\ell \geq 0$ . Thus

$$\partial f_2 = \partial(\delta h_2) \text{ for some } h_2 \in L_a^2(U_\alpha^{\ell+1}).$$

Now,

$$\Delta(\delta(h_2)) = \delta(\partial(\delta(h_2))) + \partial(\delta(\delta(h_2))) = \delta(\partial(\delta(h_2))) = \delta(\partial(f_2)) \quad (13)$$

since  $\delta^2 = 0$ . However, from the decomposition for  $f$  we have, since  $\delta f_3 = 0$

$$\delta f = \delta(\partial f_2)$$

and since  $f$  is continuous  $\delta f$  is continuous, and therefore  $\delta(\partial f_2)$  is continuous. It then follows from Poisson regularity and (13) that  $\delta h_2$  is continuous as to be shown. A dual argument shows that  $\delta f_1 = \delta(\partial h_1)$  where  $\partial h_1$  is continuous, completing the proof.  $\square$

Notice that a somewhat weaker result than Poisson regularity would imply that  $f_3$  above is continuous, namely regularity of harmonic functions.

**Question 4** (Harmonic Regularity Problem). For  $\alpha > 0$ , and  $\ell > 0$ , suppose that  $\Delta f = 0$  where  $f \in L_a^2(U_\alpha^{\ell+1})$ . What conditions on  $(X, d, \mu)$  would imply  $f$  is continuous?

Under some additional conditions on the measure, we have answered this for  $\ell = 0$  (see Section 5) and can do so for  $\ell = 1$ , which we now consider.

We assume in addition that the inclusion of continuous functions into  $L^2$ -functions induces an epimorphism of the associated Alexander-Spanier-Čech cohomology groups, i.e. that every cohomology class in the  $L^2$ -theory has a continuous representative. In Section 9 we will see that this is often the case.

Let now  $f \in L_a^2(U_\alpha^2)$  be harmonic. Let  $g$  be a continuous function in the same cohomology class. Then there is  $x \in L_a^2(U_\alpha^1)$  such that  $f = g + dx$ . As  $\delta^* f = 0$  it follows that  $\delta^* dx = -\delta^* g$  is continuous. If the Poisson regularity property in degree zero holds (compare Proposition 14 of Section 5) then  $x$  is continuous and therefore also  $f = g + dx$  is continuous.

Thus we have the following proposition.

**Proposition 17.** *Assume that  $\mu(S_x \cap A)$  are continuous for  $x \in X$  and all  $A$  measurable. Assume that every cohomology class of degree 1 has a continuous representative. If  $f$  is an  $\alpha$ -harmonic 1-form in  $L_a^2(U_\alpha^2)$ , then  $f$  is continuous.*

As in Section 5, if  $\mu$  is Borel regular, it suffices that the hypotheses hold for all  $A$  closed (or all  $A$  open).

## 9 Finite dimensional cohomology

In this section, we will establish conditions on  $X$  and  $\alpha > 0$  that imply that the  $\alpha$  cohomology is finite dimensional. In particular, in the case of the  $L^2$ - $\alpha$  cohomology, they imply that the image of  $\delta$  is closed, and that Hodge theory for  $X$  at scale  $\alpha$  holds. Along the way, we will compute the  $\alpha$ -cohomology in terms of ordinary Čech cohomology of a covering and that the different variants of our Alexander-Spanier-Čech cohomology at fixed scale ( $L^2$ , continuous, ...) are all isomorphic. We then show that the important class of metric spaces, Riemannian manifolds satisfy these conditions for  $\alpha$  small. In particular, in this case the  $\alpha$ -cohomology will be isomorphic to ordinary cohomology with  $\mathbb{R}$ -coefficients.

Throughout this section,  $(X, d)$  will denote a compact metric space,  $\mu$  a Borel probability measure on  $X$  such that  $\mu(U) > 0$  for all nonempty open sets  $U \subset X$ , and  $\alpha > 0$ . As before  $U_\alpha^\ell$  will denote the closed  $\alpha$ -neighborhood of the diagonal in  $X^\ell$ . We will denote by  $F_a(U_\alpha^\ell)$  the space of all alternating real valued functions on  $U_\alpha^\ell$ , by  $C_a(U_\alpha^\ell)$  the continuous alternating real valued functions on  $U_\alpha^\ell$ , and by  $L_a^p(U_\alpha^\ell)$  the  $L^p$  alternating real valued functions on  $U_\alpha^\ell$  for  $p \geq 1$  (in particular, the case  $p = 2$  was discussed in the preceding sections). If  $X$  is a smooth Riemannian manifold,  $C_a^\infty(U_\alpha^\ell)$  will be the smooth alternating real valued functions on  $U_\alpha^\ell$ . We will be interested in the following cochain complexes:

$$0 \longrightarrow L_a^p(X) \xrightarrow{\delta_0} L_a^p(U_\alpha^2) \xrightarrow{\delta_1} \dots \xrightarrow{\delta_{\ell-1}} L_a^p(U_\alpha^{\ell+1}) \xrightarrow{\delta_\ell} \dots$$

$$0 \longrightarrow C_a(X) \xrightarrow{\delta_0} C_a(U_\alpha^2) \xrightarrow{\delta_1} \dots \xrightarrow{\delta_{\ell-1}} C_a(U_\alpha^{\ell+1}) \xrightarrow{\delta_\ell} \dots$$

$$0 \longrightarrow F_a(X) \xrightarrow{\delta_0} F_a(U_\alpha^2) \xrightarrow{\delta_1} \dots \xrightarrow{\delta_{\ell-1}} F_a(U_\alpha^{\ell+1}) \xrightarrow{\delta_\ell} \dots$$

And if  $X$  is a smooth Riemannian manifold,

$$0 \longrightarrow C_a^\infty(X) \xrightarrow{\delta_0} C_a^\infty(U_\alpha^2) \xrightarrow{\delta_1} \dots \xrightarrow{\delta_{\ell-1}} C_a^\infty(U_\alpha^{\ell+1}) \xrightarrow{\delta_\ell} \dots$$

The corresponding cohomology spaces  $\text{Ker } \delta_\ell / \text{Im } \delta_{\ell-1}$  will be denoted by  $H_{\alpha, L^p}^\ell(X)$ , or briefly  $H_{\alpha, L^p}^\ell$ ,  $H_{\alpha, cont}^\ell$ ,  $H_\alpha^\ell$  and  $H_{\alpha, smooth}^\ell$  respectively. The proof

of finite dimensionality of these spaces, under certain conditions, involves the use of bicomplexes, some facts about which we collect here.

A bicomplex  $C^{*,*}$  will be a rectangular array of vector spaces  $C^{j,k}$ ,  $j, k \geq 0$ , and linear maps (coboundary operators)  $c_{j,k}: C^{j,k} \rightarrow C^{j+1,k}$ , and  $d_{j,k}: C^{j,k} \rightarrow C^{j,k+1}$  such that the rows and columns are chain complexes, that is  $c_{j+1,k}c_{j,k} = 0$ ,  $d_{j,k+1}d_{j,k} = 0$ , and  $c_{j,k+1}d_{j,k} = d_{j+1,k}c_{j,k}$ . Given such a bicomplex, we associate the total complex  $E^*$ , a chain complex

$$0 \longrightarrow E^0 \xrightarrow{D_0} E^1 \xrightarrow{D_1} \dots \xrightarrow{D_{\ell-1}} E^\ell \xrightarrow{D_\ell} \dots$$

where  $E^\ell = \bigoplus_{j+k=\ell} C^{j,k}$  and where on each term  $C^{j,k}$  in  $E^\ell$ ,  $D_\ell = c_{j,k} + (-1)^k d_{j,k}$ . Using commutativity of  $c$  and  $d$ , one can easily check that  $D_{\ell+1}D_\ell = 0$ , and thus the total complex is a chain complex. We recall a couple of definitions from homological algebra. If  $E^*$  and  $F^*$  are cochain complexes of vector spaces with coboundary operators  $e$  and  $f$  respectively, then a chain map  $g: E^* \rightarrow F^*$  is a collection of linear maps  $g_j: E^j \rightarrow F^j$  that commute with  $e$  and  $f$ . A chain map induces a map on cohomology. A cochain complex  $E^*$  is said to be exact at the  $k$ th term if the kernel of  $e_k: E_k \rightarrow E_{k+1}$  is equal to the image of  $e_{k-1}: E_{k-1} \rightarrow E_k$ . Thus the cohomology at that term is zero.  $E^*$  is defined to be exact if it is exact at each term. A chain contraction  $h: E^* \rightarrow E^*$  is a family of linear maps  $h_j: E^j \rightarrow E^{j-1}$  such that  $e_{j-1}h_j + h_{j+1}e_j = \text{Id}$ . The existence of a chain contraction on  $E^*$  implies that  $E^*$  is exact. The following fact from homological algebra is fundamental in proving finite dimensionality of our cohomology spaces.

**Lemma 4.** *Suppose that  $C^{*,*}$  is a bicomplex as above, and  $E^*$  is the associated total complex. Suppose that we augment the bicomplex with a column on the left which is a chain complex  $C^{-1,*}$ ,*

$$C^{-1,0} \xrightarrow{d_{-1,0}} C^{-1,1} \xrightarrow{d_{-1,1}} \dots \xrightarrow{d_{-1,\ell-1}} C^{-1,\ell} \xrightarrow{d_{-1,\ell}} \dots$$

and linear maps  $c_{-1,k}: C^{-1,k} \rightarrow C^{0,k}$ , such that the augmented rows

$$0 \longrightarrow C^{-1,k} \xrightarrow{c_{-1,k}} C^{0,k} \xrightarrow{c_{0,k}} \dots \xrightarrow{c_{\ell-1,k}} C^{\ell,k} \xrightarrow{c_{\ell,k}} \dots$$

are chain complexes with  $d_{0,k}c_{-1,k} = c_{-1,k+1}d_{-1,k}$ . Then, the maps  $c_{-1,k}$  induce a chain map  $c_{-1,*}: C^{-1,*} \rightarrow E^*$ . Furthermore, if the first  $K$  rows of the augmented complex are exact, then  $c_{-1,*}$  induces an isomorphism on the homology of the complexes  $c_{-1,*}: H^k(C^{-1,*}) \rightarrow H^k(E^*)$  for  $k \leq K$  and an injection for  $k = K+1$ . In fact, one only needs exactness of the first  $K$  rows up to the  $K$ th term  $C^{K,j}$ .

A simple proof of this is given in Bott and Tu [4, pages 95–97], in the case of the Čech-de Rham complex, but the the proof generalizes to the abstract setting. Of course, if we augmented the bicomplex with a row  $C^{*, -1}$  with the same properties, the conclusions would hold. In fact, we will show the cohomologies of two chain complexes are isomorphic by augmenting a bicomplex as above with one such row and one such column.

**Corollary 3.** *Suppose that  $C^{*,*}$  is a bicomplex as in the Lemma, and that  $C^{*,*}$  is augmented with a column  $C^{-1,*}$  as in the Lemma, and a row  $C^{*, -1}$  also a chain complex with coboundary operators  $c_{j,-1}: C^{j,-1} \rightarrow C^{j+1,-1}$  and linear maps  $d_{j,-1}: C^{j,-1} \rightarrow C^{j,0}$  such that the augmented columns*

$$0 \longrightarrow C^{j,-1} \xrightarrow{d_{j,-1}} C^{0,k} \xrightarrow{d_{j,-1}} \dots \xrightarrow{d_{j,\ell-1}} C^{j,\ell} \xrightarrow{d_{j,\ell}} \dots$$

*are chain complexes, and  $c_j, 0d_{j,-1} = d_{j+1,-1}c_{j,-1}$ . Then, if the first  $K$  rows are exact and the first  $K+1$  columns are exact, up to the  $K+1$  term, it follows that the cohomology  $H^\ell(C^{-1,*})$  of  $C^{-1,*}$  and  $H^\ell(C^{*, -1})$  of  $C^{*, -1}$  are isomorphic for  $0 \leq K$ , and  $H^{K+1}(C^{-1,*})$  is isomorphic to a subspace of  $H^{K+1}(C^{*, -1})$ .*

*Proof.* This follows immediately from the lemma, as the cohomology up to order  $K$  of both  $C^{-1,*}$  and  $C^{*, -1}$  are isomorphic to the cohomology of the total complex. Also,  $H^{K+1}(C^{-1,*})$  is isomorphic to a subspace of  $H^{K+1}(E^*)$  which is isomorphic to  $H^{K+1}(C^{*, -1})$ .  $\square$

**Remark 9.** If all of the spaces  $C^{j,k}$  in the Lemma and Corollary are Banach spaces, and the coboundaries,  $c_{j,k}$  and  $d_{j,k}$  are bounded, then the isomorphisms of cohomology can be shown to be topological isomorphisms, where the topologies on the cohomology spaces are induced by the quotient semi-norms.

Let  $\{V_i, i \in S\}$  be a finite covering of  $X$  by Borel sets (usually taken to be balls). We construct the corresponding Čech- $L^p$ -Alexander bicomplex at scale  $\alpha$  as follows.

$$C^{k,\ell} = \bigoplus_{I \in S^{k+1}} L_a^p(U_\alpha^{\ell+1} \cap V_I^{\ell+1}) \quad \text{for } k, \ell \geq 0$$

where we use the abbreviation  $V_I = V_{i_0, \dots, i_k} = \bigcap_{j=0}^k V_{i_j}$ . The vertical coboundary  $d_{k,\ell}$  is just the usual coboundary  $\delta_\ell$  as in Section 4, acting on each  $L_a^p(U_\alpha^{\ell+1} \cap V_{I^{\ell+1}})$ . The horizontal coboundary  $c_{k,\ell}$  is the “Čech differential”. More explicitly, if  $f \in C^{k,\ell}$ , then it has components  $f_I$  which are functions on  $U_\alpha^{\ell+1} \cap V_I^{\ell+1}$  for each  $(k+1)$ -tuple  $I$ , and for any  $k+2$  tuple  $J = (j_0, \dots, j_{k+1})$ ,  $cf$  is defined on  $U_\alpha^{\ell+1} \cap V_J^{\ell+1}$  by

$$(c_{k,\ell}f)_J = \sum_{i=0}^{k+1} (-1)^i f_{j_0, \dots, \hat{j}_i, \dots, j_{k+1}} \quad \text{restricted to } V_J^{\ell+1}.$$

It is not hard to check that the coboundaries commute  $c\delta = \delta c$ . We augment the complex on the left with the column (chain complex)  $C^{-1,\ell} = L_a^p(U_\alpha^{\ell+1})$  with horizontal map  $c_{-1,\ell}$  equal to restriction on each  $V_i$  and vertical map the usual coboundary. We augment the complex on the bottom with the chain complex  $C^{*, -1}$  which is the Čech complex of the cover  $\{V_i\}$ . That is an element  $f \in C^{k,-1}$  is a function that assigns to each  $V_I$  a real number or equivalently  $C^{k,-1} = \bigoplus_{I \in S^{k+1}} \mathbb{R}V_I$ . The vertical maps are just inclusions into  $C^{*,0}$ , and the horizontal maps are the Čech differential as defined above.

**Remark 10.** We can similarly define the Čech-Alexander bicomplex, the Čech-Continuous Alexander bicomplex and the Čech-Smooth Alexander bicomplex (in case  $X$  is a smooth Riemannian manifold) by replacing  $L_a^p$  everywhere in the above complex with  $F_a$ ,  $C_a$  and  $C_a^\infty$  respectively.

**Remark 11.** The cohomology spaces of  $C^{*, -1}$  are finite dimensional since the cover  $\{V_i\}$  is finite. This is called the Čech cohomology of the cover, and is the same as the simplicial cohomology of the simplicial complex that is the nerve of the cover  $\{V_i\}$ .

We will use the above complex to show, under some conditions, that  $H_{\alpha, L^p}^\ell$ ,  $H_\alpha^\ell$  and  $H_{\alpha, \text{cont}}^\ell$  are isomorphic to the Čech cohomology of an appropriate finite open cover of  $X$  and thus finite dimensional.

**Theorem 7.** *Let  $\{V_i\}_{i \in S}$  be a finite cover of  $X$  by Borel sets as above, and assume that  $\{V_i^{K+1}\}_{i \in S}$  is a cover for  $U_\alpha^{K+1}$  for some  $K \geq 0$ . Assume also that the first  $K+1$  columns of the corresponding Čech- $L^p$ -Alexander complex are exact up to the  $K+1$  term. Then  $H_{\alpha, L^p}^\ell$  is isomorphic to  $H^\ell(C^{*, -1})$  for  $\ell \leq K$  and is thus finite dimensional. Also  $H_{\alpha, L^p}^{K+1}$  is isomorphic to a subspace of  $H^{K+1}(C^{*, -1})$ . If  $\{V_i\}_{i \in S}$  is an open cover, then the same conclusion holds for  $H_\alpha^\ell$ ,  $H_{\alpha, \text{cont}}^\ell$  and  $H_{\alpha, \text{smooth}}^\ell$  (in case  $X$  is a smooth Riemannian manifold), and hence all are isomorphic to each other. Those isomorphisms are induced by the natural inclusion maps of smooth functions into continuous functions into  $L^q$ -functions into  $L^p$ -functions ( $q \geq p$ ) into arbitrary real valued functions.*

*Proof.* In light of the corollary above, it suffices to show that the first  $K$  rows of the bicomplex are exact. Indeed, we are computing the sheaf cohomology of  $U_\alpha^{k+1}$  for a flabby sheaf (the sheaf of smooth or continuous or  $L^p$  or arbitrary functions) which vanishes. We write out the details: Note that for  $\ell \leq K$ ,  $\{V_i^{\ell+1}\}$  covers  $U_\alpha^{\ell+1}$  and therefore  $c_{-1, \ell}: L_a^p(U_\alpha^{\ell+1}) \rightarrow \bigoplus_{i \in S} L_a^p(U_\alpha^{\ell+1} \cap V_i^{\ell+1})$  is injective (as  $c_{-1, \ell}$  is restriction), and therefore we have exactness at the first term. In general, we construct a chain contraction  $h$  on the  $\ell$ th row. Let  $\{\phi_i\}$  be a measurable partition of unity for  $U_\alpha^{\ell+1}$  subordinate to the cover  $\{U_\alpha^{\ell+1} \cap V_i^{\ell+1}\}$  (thus support  $\phi_i \subset U_\alpha^{\ell+1} \cap V_i^{\ell+1}$  and  $\sum_i \phi_i(x) = 1$  for all  $x$ ). Then define

$$h: \bigoplus_{I \in S^{k+1}} L_a^p(U_\alpha^{\ell+1} \cap V_I^{\ell+1}) \rightarrow \bigoplus_{I \in S^k} L_a^p(U_\alpha^{\ell+1} \cap V_I^{\ell+1})$$

for each  $k$  by  $(hf)_{i_0, \dots, i_{k-1}} = \sum_{j \in S} \phi_j f_{j, i_0, \dots, i_{k-1}}$ . We show that  $h$  is a chain contraction, that is  $ch + hc = \text{Id}$ :

$$(c(hf))_{i_0, \dots, i_{k-1}} = \sum_{n=0}^{k-1} (-1)^n (hf)_{i_0, \dots, \hat{i}_n, \dots, i_{k-1}} = \sum_{j, n} (-1)^n \phi_j f_{j, i_0, \dots, \hat{i}_n, \dots, i_{k-1}}.$$

Now,

$$\begin{aligned}
(h(cf))_{i_0, \dots, i_{k-1}} &= \sum_{j \in S} \phi_j(cf)_{j, i_0, \dots, i_{k-1}} \\
&= \sum_j \phi_j(f_{i_0, \dots, i_{k-1}} - \sum_n^{k-1} (-1)^n f_{j, i_0, \dots, \hat{i}_n, \dots, i_{k-1}}) \\
&= f_{i_0, \dots, i_{k-1}} - (c(hf))_{i_0, \dots, i_{k-1}}.
\end{aligned}$$

Thus  $h$  is a chain contraction for the  $\ell$ th row, proving exactness (note that exactness follows, since if  $cf = 0$  then from above  $c(hf) = f$ ). If  $\{V_i\}$  is an open cover, then the partition of unity  $\{\phi_i\}$  can be chosen to be continuous, or even smooth in case  $X$  is a smooth Riemannian manifold. Then  $h$  as defined above is a chain contraction on the corresponding complexes with  $L_a^p$  replaced by  $F_a$ ,  $C_a$  or  $C_a^\infty$ .

Observe that the inclusions  $C^\infty \hookrightarrow C^0 \hookrightarrow L^q \hookrightarrow L^p \hookrightarrow F$  (where  $F$  stands for arbitrary real valued functions) extend to inclusions of the augmented bi-complexes, whose restriction to the Čech column  $C^{*, -1}$  is the identity. As the identity clearly induces an isomorphism in cohomology, and the inclusion of this augmented bottom row into the (non-augmented) bicomplex also does, by naturality the various inclusions of the bicomplexes induce isomorphisms in cohomology. The same argument applied backwards to the inclusions of the Alexander-Spanier-Čech rows into the bicomplexes shows that the inclusions of the smaller function spaces into the larger function spaces induce isomorphisms in  $\alpha$ -cohomology.

This finishes the proof of the theorem.  $\square$

We can use Theorem 7 to prove finite dimensionality of the cohomologies in general, for  $\ell = 0$  and 1.

**Theorem 8.** *For any compact  $X$  and any  $\alpha > 0$ ,  $H_{\alpha, L^p}^\ell$ ,  $H_\alpha^\ell$ ,  $H_{\alpha, \text{cont}}^\ell$ , and  $H_{\alpha, \text{smooth}}^\ell$  ( $X$  a smooth manifold) are finite dimensional and are isomorphic, for  $\ell = 0, 1$ .*

Let  $\{V_i\}$  be a covering of  $X$  by open balls of radius  $\alpha/3$ . Then the first row ( $\ell = 0$ ) of the Čech- $L^p$ -Alexander Complex is exact from the proof of Theorem 7 (taking  $K = 0$ ). It suffices to show that the columns are exact. Note that  $V_I^{\ell+1} \subset U_\alpha^{\ell+1}$  trivially, for each  $\ell$  and  $I \in S^{k+1}$  because  $\text{diam}(V_I) < \alpha$ . For  $k$  fixed, and  $I \in S^{k+1}$  we define  $g: L_a^p(V_I^{\ell+1}) \rightarrow L_a^p(V_I^\ell)$  by

$$gf(x_0, \dots, x_{\ell-1}) = \frac{1}{\mu(V_I)} \int_{V_I} f(t, x_0, \dots, x_{\ell-1}) d\mu(t).$$

We check that  $g$  defines a chain contraction:

$$\begin{aligned}
\delta(gf)(x_0, \dots, x_\ell) &= \sum_i (-1)^i (gf)(x_0, \dots, \hat{x}_i, \dots, x_\ell) \\
&= \sum_i (-1)^i \frac{1}{\mu(V_I)} \int_{V_I} f(t, x_0, \dots, \hat{x}_i, \dots, x_\ell) d\mu(t).
\end{aligned}$$

But,

$$\begin{aligned} g(\delta f)(x_0, \dots, x_\ell) &= \frac{1}{\mu(V_I)} \int_{V_I} \delta f(t, x_0, \dots, x_\ell) d\mu(t) \\ &= \frac{1}{\mu(V_I)} \left( \int_{V_I} f((x_0, \dots, x_\ell) d\mu(t) - \sum_i (-1)^i \int_{V_I} f(t, \dots, \hat{x}_i, \dots, x_\ell) d\mu(t) \right) \\ &= f(x_0, \dots, x_\ell) - \delta(gf)(x_0, \dots, x_\ell). \end{aligned}$$

Thus  $g$  defines a chain contraction on the  $k$ th column and the columns are exact. For the corresponding Alexander, continuous and smooth bicomplexes, a chain contraction can be defined by fixing for each  $V_I$ ,  $I \in S^{k+1}$  a point  $p \in V_I$  and setting  $gf(x_0, \dots, x_{\ell-1}) = f(p, x_0, \dots, x_{\ell-1})$ . This is easily verified to be a chain contraction, finishing the proof of the theorem.

Recall that for  $x = (x_0, \dots, x_{\ell-1}) \in U_\alpha^\ell$  we define the slice  $S_x = \{t \in X : (t, x_0, \dots, x_{\ell-1}) \in U_\alpha^{\ell+1}\}$ . We consider the following hypothesis on  $X$ ,  $\alpha > 0$  and non-negative integer  $K$ :

**Definition 2. Hypothesis (\*):** There exists a  $\delta > 0$  such that whenever  $V = \cap_i V_i$  is a non-empty intersection of finitely many open balls of radius  $\alpha + \delta$ , then there is a Borel set  $W$  of positive measure such that for each  $\ell \leq K + 1$

$$W \subset V \cap \left( \bigcap_{x \in U_\alpha^\ell \cap V^\ell} S_x \right).$$

**Theorem 9.** Assume that  $X$ ,  $\alpha > 0$  and  $K$  satisfy hypothesis (\*). Then, for  $\ell \leq K$ ,  $H_{\alpha, L^p}^\ell$ ,  $H_\alpha^\ell$ ,  $H_{\alpha, cont}^\ell$ , and  $H_{\alpha, smooth}^\ell$  (in the case  $X$  is a smooth Riemannian manifold) are all finite dimensional, and are isomorphic to the Čech cohomology of some finite covering of  $X$  by open balls of radius  $\alpha + \delta$ . Furthermore, the Hodge theorem for  $X$  at scale  $\alpha$  holds (Theorem 4 of Section 4).

*Proof.* Let  $\{V_i\}$ ,  $i \in S$  be a finite open cover of  $X$  by balls of radius  $\alpha + \delta$  such that  $\{V_i^{K+1}\}$  is a covering for  $U_\alpha^{K+1}$ . This can always be done since  $U_\alpha^{K+1}$  is compact. We first consider the case of the Čech- $L^p$ -Alexander bicomplex corresponding to the cover. By Theorem 7, it suffices to show that there is a chain contraction of the columns up to the  $K$ th term. For each  $I \in S^{k+1}$ , and  $\ell \leq K + 1$  let  $W$  be the Borel set of positive measure assumed to exist in (\*) with  $V_I$  playing the role of  $V$  in (\*). Then we define  $g: L_a^p(U_\alpha^{\ell+1} \cap V_I^{\ell+1}) \rightarrow L_a^p(U_\alpha^\ell \cap V_I^\ell)$  by

$$gf(x_0, \dots, x_{\ell-1}) = \frac{1}{\mu(W)} \int_W f(t, x_0, \dots, x_{\ell-1}) d\mu(t).$$

The hypothesis (\*) implies that  $g$  is well defined. The proof that  $g$  defines a chain contraction on the  $k$ th column (up to the  $K$ th term) is identical to the one in the proof of Theorem 8. As in the proof of Theorem 8, the chain



contraction for the case when  $L_a^p$  is replaced by  $F_a$ ,  $C_a$  and  $C_a^\infty$  can be taken to be  $gf(x_0, \dots, x_{\ell-1}) = f(p, x_0, \dots, x_{\ell-1})$  for some fixed  $p \in W$ . Note that in these cases, we don't require that  $\mu(W) > 0$ , only that  $W \neq \emptyset$ .  $\square$

**Remark 12.** If  $X$  satisfies certain local conditions as in Wilder [36], then the Čech cohomology of the cover, for small  $\alpha$ , is isomorphic to the Čech cohomology of  $X$ .

Our next goal is to give somewhat readily verifiable conditions on  $X$  and  $\alpha$  that will imply (\*). This involves the notion of midpoint and radius of a closed set in  $X$ .

Let  $\Lambda \subset X$  be closed. We define the radius  $r(\Lambda)$  by  $r(\Lambda) = \inf\{\beta : \cap_{x \in \Lambda} B_\beta(x) \neq \emptyset\}$  where  $B_\beta(x)$  denotes the closed ball of radius  $\beta$  centered at  $x$ .

**Proposition 18.**  $\cap_{x \in \Lambda} B_{r(\Lambda)}(x) \neq \emptyset$ . Furthermore, if  $p \in \cap_{x \in \Lambda} B_{r(\Lambda)}(x)$ , then  $\Lambda \subset B_{r(\Lambda)}(p)$ , and if  $\Lambda \subset B_\beta(q)$  for some  $q \in \Lambda$ , then  $r(\Lambda) \leq \beta$ .

Such a  $p$  is called a midpoint of  $\Lambda$ .

*Proof.* Let  $J = \{\beta \in \mathbb{R} : \cap_{x \in \Lambda} B_\beta(x) \neq \emptyset\}$ . For  $\beta \in J$  define  $R_\beta = \cap_{x \in \Lambda} B_\beta(x)$ . Note that if  $\beta \in J$  and  $\beta < \beta'$ , then  $\beta' \in J$ , and  $R_\beta \subset R_{\beta'}$ .  $R_\beta$  is compact, and therefore  $\cap_{\beta \in J} R_\beta \neq \emptyset$ . Let  $p \in \cap_{\beta \in J} R_\beta$ . Then, for  $x \in \Lambda$ ,  $p \in B_\beta(x)$  for all  $\beta \in J$  and so  $d(p, x) \leq \beta$ . Taking the infimum of this over  $\beta \in J$  yields  $d(p, x) \leq r(\Lambda)$  or  $p \in R_{r(\Lambda)}$  proving the first assertion of the proposition. Now, if  $x \in \Lambda$  then  $p \in B_{r(\Lambda)}(x)$  which implies  $x \in B_{r(\Lambda)}(p)$  and thus  $\Lambda \subset B_{r(\Lambda)}(p)$ . Now suppose that  $\Lambda \subset B_\beta(q)$  for some  $q \in \Lambda$ . Then for every  $x \in \Lambda$ ,  $q \in B_\beta(x)$  and thus  $\cap_{x \in \Lambda} B_\beta(x) \neq \emptyset$  which implies  $\beta \geq r(\Lambda)$  finishing the proof.  $\square$

We define  $\mathcal{K}(X) = \{\Lambda \subset X : \Lambda \text{ is compact}\}$ , and we endow  $\mathcal{K}(X)$  with the Hausdorff metric  $D(A, B) = \max\{\sup_{t \in B} d(t, A), \sup_{t \in A} d(t, B)\}$ . We also define, for  $x = (x_0, \dots, x_\ell) \in U_\alpha^{\ell+1}$ , the witness set of  $x$  by  $w_\alpha(x) = \cap_i B_\alpha(x_i)$  (we are suppressing the dependence of  $w_\alpha$  on  $\ell$ ). Thus  $w_\alpha : U_\alpha^{\ell+1} \rightarrow \mathcal{K}(X)$ . We have

**Theorem 10.** Let  $X$  be compact, and  $\alpha > 0$ . Suppose that  $w_\alpha : U_\alpha^{\ell+1} \rightarrow \mathcal{K}(X)$  is continuous for  $\ell \leq K+1$ , and suppose there exists  $\delta_0 > 0$  such that whenever  $\Lambda = \cap_{i=0}^k B_i$  is a finite intersection of closed balls of radius  $\alpha + \delta$ ,  $\delta \in (0, \delta_0]$  then  $r(\Lambda) \leq \alpha + \delta$ . Then Hypothesis (\*) holds.

The proof will follow from

**Proposition 19.** Under the hypotheses of Theorem 10, given  $\epsilon > 0$ , there exists  $\delta > 0$ ,  $\delta \leq \delta_0$  such that for all  $\beta \in [\alpha, \alpha + \delta]$  we have  $D(w_\alpha(\sigma), w_\beta(\sigma)) \leq \epsilon$  for all simplices  $\sigma \in U_\alpha^{\ell+1} \subset U_\beta^{\ell+1}$ .

*Proof of Theorem 10.* Fix  $\epsilon < \alpha$ , and let  $\delta > 0$  be as in Proposition 19. Let  $\{V_i\}$  be a finite collection of open balls of radius  $\alpha + \delta$  such that  $\cap_i V_i \neq \emptyset$ , and

let  $\{B_i\}$  be the corresponding collection of closed balls of radius  $\alpha + \delta$ . Define  $\Lambda$  to be the closure of  $\cap_i V_i$  and thus

$$\Lambda = \overline{\cap_i V_i} \subset \cap_i \overline{V_i} \subset \cap_i B_i.$$

Let  $p$  be a midpoint of  $\Lambda$ . We will show that  $d(p, w_\alpha(\sigma)) \leq \epsilon$  for any  $\sigma = (x_0, \dots, x_{\ell+1}) \in \Lambda^{\ell+1}$ . We have

$$p \in \cap_{x \in \Lambda} B_{r(\Lambda)}(x) \subset \cap_{i=0}^{\ell+1} B_{r(\Lambda)}(x_i) = w_{r(\Lambda)}(\sigma) \subset w_{\alpha+\delta}(\sigma)$$

since  $r(\Lambda) \leq \alpha + \delta$ . But  $D(w_\alpha(\sigma), w_{\alpha+\delta}(\sigma)) \leq \epsilon$  from Proposition 19, and so  $d(p, w_\alpha(\sigma)) \leq \epsilon$ . In particular, there exists  $q \in w_\alpha(\sigma)$  with  $d(p, q) \leq \epsilon$ . Now, if  $x \in B_{\alpha-\epsilon}(p) \cap \Lambda$ , then  $d(x, q) \leq d(x, p) + d(p, q) \leq \alpha - \epsilon + \epsilon = \alpha$ . Therefore  $(x, x_0, \dots, x_\ell) \in U_\alpha^{\ell+2}$  and so  $x \in S_\sigma \cap \Lambda$ . Thus  $B_{\alpha-\epsilon}(p) \cap \Lambda \subset \cap_{\sigma \in U_\alpha^{\ell+1} \cap \Lambda^{\ell+1}} S_\sigma$ . Let  $B'_s(p)$  denote the open ball of radius  $s$  and let  $V = \cap_i V_i$ . Then define  $W = B'_{\alpha-\epsilon}(p) \cap V$ . Then  $W$  is a nonempty open set (since  $p \in \overline{V}$ ),  $\mu(W) > 0$  and  $W \subset \cap_{\sigma \in U_\alpha^{\ell+1} \cap V^{\ell+1}} S_\sigma$  and Hypothesis (\*) is satisfied finishing the proof of Theorem 10.  $\square$

*Proof of Proposition 19.* Let  $\epsilon > 0$ . Note that for  $\beta \geq \alpha$ , and  $\sigma \in U_\alpha^{\ell+1}$ ,  $w_\alpha(\sigma) \subset w_\beta(\sigma)$ . It thus suffices to show that there exists  $\delta > 0$  such that

$$\sup_{x \in w_\beta(\sigma)} d(x, w_\alpha(\sigma)) \leq \epsilon \text{ for all } \beta \in [\alpha, \alpha + \delta].$$

Suppose that this is not the case. Then there exists  $\beta_j \downarrow \alpha$  and  $\sigma_j \in U_\alpha^{\ell+1}$  such that

$$\sup_{x \in w_{\beta_j}(\sigma_j)} d(x, w_\alpha(\sigma_j)) > \epsilon$$

and thus there exists  $x_n \in w_{\beta_n}(\sigma_n)$  with  $d(x_n, w_\alpha(\sigma_n)) \geq \epsilon$ . Let  $\sigma_n = (y_0^n, \dots, y_\ell^n)$ . Thus  $d(x_n, y_i^n) \leq \beta_n$  for all  $i$ . By compactness, after taking a subsequence, we can assume  $\sigma_n \rightarrow \sigma = (y_0, \dots, y_\ell)$  and  $x_n \rightarrow x$ . Thus  $d(x, y_i) \leq \alpha$  for all  $i$  and  $\sigma \in U_\alpha^{\ell+1}$ , and  $x \in w_\alpha(\sigma)$ . However, by continuity of  $w_\alpha$ ,  $w_\alpha(\sigma_n) \rightarrow w_\alpha(\sigma)$  which implies  $d(x, w_\alpha(\sigma)) \geq \epsilon$  (since  $d(x_n, w_\alpha(\sigma_n)) \geq \epsilon$ ) a contradiction, finishing the proof of the proposition.  $\square$

We now turn to the case where  $X$  is a compact Riemannian manifold of dimension  $n$ , with Riemannian metric  $g$ . We will always assume that the metric  $d$  on  $X$  is induced from  $g$ . Recall that a set  $\Lambda \subset X$  is strongly convex if given  $p, q \in \Lambda$ , then the length minimizing geodesic from  $p$  to  $q$  is unique, and lies in  $\Lambda$ . The strong convexity radius at a point  $x \in X$  is defined by  $\rho(x) = \sup\{r : B_r(x) \text{ is strongly convex}\}$ . The strong convexity radius of  $X$  is defined as  $\rho(X) = \inf\{\rho(x) : x \in X\}$ . It is a basic fact of Riemannian geometry that for  $X$  compact,  $\rho(X) > 0$ . Thus for any  $x \in X$  and  $r < \rho(X)$ ,  $B_r(x)$  is strongly convex.

**Theorem 11.** *Assume as above that  $X$  is a compact Riemannian manifold. Let  $k > 0$  be an upper bound for the sectional curvatures of  $X$  and let  $\alpha < \min\{\rho(X), \frac{\pi}{2\sqrt{k}}\}$ . Then Hypothesis (\*) holds.*

**Corollary 4.** *In the situation of Theorem 11, the cohomology groups  $H_{\alpha, L^p}^\ell$ ,  $H_\alpha^\ell$ ,  $H_{\alpha, \text{cont}}^\ell$ , and  $H_{\alpha, \text{smooth}}^\ell$  are finite dimensional and isomorphic to each other and to the ordinary cohomology of  $X$  with real coefficients (and the natural inclusions induce the isomorphisms). Moreover, Hodge theory for  $X$  at scale  $\alpha$  holds.*

*Proof of Theorem 11.* From Theorem 10, it suffices to prove the following propositions.

**Proposition 20.** *Let  $\alpha < \min\{\rho(X), \frac{\pi}{2\sqrt{k}}\}$ . Then  $w_\alpha: U_\alpha^{\ell+1} \rightarrow \mathcal{K}(X)$  is continuous for  $\ell \leq K$ .*

**Proposition 21.** *Let  $\delta > 0$  such that  $\alpha + \delta < \min\{\rho(X), \frac{\pi}{2\sqrt{k}}\}$ . Whenever  $\Lambda$  is a closed, convex set in some  $B_{\alpha+\delta}(z)$ , then  $r(\Lambda) \leq \alpha + \delta$ .*

Of course, the conclusion of Proposition 21 is stronger than the second hypothesis of Theorem 10, since the finite intersection of balls of radius  $\alpha + \delta$  is convex and  $\alpha + \delta < \rho(X)$ .

*Proof of Proposition 20.* We start with

**Claim 1.** Let  $\sigma = (x_0, \dots, x_\ell) \in U_\alpha^{\ell+1}$ , and suppose that  $p, q \in w_\alpha(\sigma)$  and that  $x$  is on the minimizing geodesic from  $p$  to  $q$  (but not equal to  $p$  or  $q$ ). Then  $B_\epsilon(x) \subset w_\alpha(\sigma)$  for some  $\epsilon > 0$ .

*Proof of Claim.* For points  $r, s, t$  in a strongly convex neighborhood in  $X$  we define  $\angle rst$  to be the angle that the minimizing geodesic from  $s$  to  $r$  makes with the minimizing geodesic from  $s$  to  $t$ . Let  $\gamma$  be the geodesic from  $p$  to  $q$ , and for fixed  $i$  let  $\phi$  be the geodesic from  $x$  to  $x_i$ . Now, the angles that  $\phi$  makes with  $\gamma$  at  $x$  satisfy  $\angle pxx_i + \angle x_ixq = \pi$  and therefore one of these angles is greater than or equal to  $\pi/2$ . Assume, without loss of generality that  $\theta = \angle pxx_i \geq \pi/2$ . Let  $c = d(x, x_i)$ ,  $r = d(p, x)$  and  $d = d(p, x_i) \leq \alpha$  (since  $p \in w_\alpha(\sigma)$ ). Now consider a geodesic triangle in the sphere of curvature  $k$  with vertices  $p', x'$ , and  $x'_i$  such that

$$d(p', x') = d(p, x) = r, \quad d(x', x'_i) = d(x, x_i) = c \quad \text{and} \quad \angle p'x'x'_i = \theta,$$

and let  $d' = d(p', x'_i)$ . Then, the hypotheses on  $\alpha$  imply that the Rauch Comparison Theorem (see for example do-Carmo [12]) holds, and we can conclude that  $d' \leq d$ . However, with  $\theta \geq \pi/2$ , it follows that on a sphere, where  $p', x', x'_i$  are inside a ball of radius less than the strong convexity radius, that  $c' < d'$ . Therefore we have  $c = c' < d' \leq d \leq \alpha$  and there is an  $\epsilon > 0$  such that  $y \in B_\epsilon(x)$  implies  $d(y, x_i) \leq \alpha$ . Taking the smallest  $\epsilon > 0$  so that this is true for each  $i = 0, \dots, \ell$  finishes the proof of the claim.  $\square$

**Corollary 5** (Corollary of Claim). *For  $\sigma \in U_\alpha^{\ell+1}$ , either  $w_\alpha(\sigma)$  consists of a single point, or every point of  $w_\alpha(\sigma)$  is an interior point or the limit of interior points.*

Now suppose that  $\sigma_j \in U_\alpha^{\ell+1}$  and  $\sigma_j \rightarrow \sigma$  in  $U_\alpha^{\ell+1}$ . We must show  $w_\alpha(\sigma_j) \rightarrow w_\alpha(\sigma)$ , that is

- (a)  $\sup_{x \in w_\alpha(\sigma_j)} d(x, w_\alpha(\sigma)) \rightarrow 0$ ,
- (b)  $\sup_{x \in w_\alpha(\sigma)} d(x, w_\alpha(\sigma_j)) \rightarrow 0$ .

In fact (a) holds for any metric space and any  $\alpha > 0$ . Suppose that (a) was not true. Then there exists a subsequence (still denoted by  $\sigma_j$ ), and  $\eta > 0$  such that

$$\sup_{x \in w_\alpha(\sigma_j)} d(x, w_\alpha(\sigma)) \geq \eta$$

and therefore there exists  $y_j \in w_\alpha(\sigma_j)$  with  $d(y_j, w_\alpha(\sigma)) \geq \eta/2$ . After taking another subsequence, we can assume  $y_j \rightarrow y$ . But if  $\sigma_j = (x_0^j, \dots, x_\ell^j)$ , and  $\sigma = (x_0, \dots, x_\ell)$ , then  $d(y_j, x_i^j) \leq \alpha$  which implies  $d(y, x_i) \leq \alpha$  for each  $i$  and thus  $y \in w_\alpha(\sigma)$ . But this is impossible given  $d(y_j, w_\alpha(\sigma)) \geq \eta/2$ .

We use the corollary to the Claim to establish (b). First, suppose that  $w_\alpha(\sigma)$  consists of a single point  $p$ . We show that  $d(p, w_\alpha(\sigma_j)) \rightarrow 0$ . Let  $p_j \in w_\alpha(\sigma_j)$  such that  $d(p, p_j) = d(p, w_\alpha(\sigma_j))$ . If  $d(p, p_j)$  does not converge to 0 then, after taking a subsequence, we can assume  $d(p, p_j) \geq \eta > 0$  for some  $\eta$ . But after taking a further subsequence, we can also assume  $p_j \rightarrow y$  for some  $y$ . However, as in the argument above it is easy to see that  $y \in w_\alpha(\sigma)$  and therefore  $y = p$ , a contradiction, and so (b) holds in this case.

Now suppose that every point in  $w_\alpha(\sigma)$  is either an interior point, or the limit of interior points. If (b) did not hold, there would be a subsequence (still denoted by  $\sigma_j$ ) such that

$$\sup_{x \in w_\alpha(\sigma)} d(x, w_\alpha(\sigma_j)) \geq \eta > 0$$

and thus there exists  $p_j \in w_\alpha(\sigma)$  such that  $d(p_j, w_\alpha(\sigma_j)) \geq \eta/2$ . After taking another subsequence, we can assume  $p_j \rightarrow p$  and  $p \in w_\alpha(\sigma)$ , and, for  $j$  sufficiently large  $d(p, w_\alpha(\sigma_j)) \geq \eta/4$ . If  $p$  is an interior point of  $w_\alpha(\sigma)$  then  $d(p, x_i) < \alpha$  for  $i = 0, \dots, \ell$ . But then, for all  $j$  sufficiently large  $d(p, x_i^j) \leq \alpha$  for each  $i$ . But this implies  $p \in w_\alpha(\sigma_j)$ , a contradiction. If  $p$  is not an interior point, then  $p$  is a limit point of interior points  $q_m$ . But then, from above,  $q_m \in w_\alpha(\sigma_{j_m})$  for  $j_m$  large which implies  $d(p, w_\alpha(\sigma_{j_m})) \rightarrow 0$ , a contradiction, thus establishing (b) and finishing the proof of Proposition 20.  $\square$

*Proof of Proposition 21.* Let  $\delta$  be such that  $\alpha + \delta < \min\{\rho(X), \frac{\pi}{2\sqrt{k}}\}$ , and let  $\Lambda$  be any closed convex set in  $B_{\alpha+\delta}(z)$ . We will show  $r(\Lambda) \leq \alpha + \delta$ . If  $z \in \Lambda$ , we are done for then  $\Lambda \subset B_{\alpha+\delta}(z)$  implies  $r(\Lambda) \leq \alpha + \delta$  by Proposition 18. If  $z \notin \Lambda$  let  $z_0 \in \Lambda$  such that  $d(z, z_0) = d(z, \Lambda)$  (the closest point in  $\Lambda$  to  $z$ ). Now let  $y_0 \in \Lambda$  such that  $d(z_0, y_0) = \max_{y \in \Lambda} d(z_0, y)$ . Let  $\gamma$  be the minimizing geodesic from  $z_0$  to  $y_0$ , and  $\phi$  the minimizing geodesic from  $z_0$  to  $z$ . Since  $\Lambda$  is convex  $\gamma$  lies on  $\Lambda$ . If  $\theta$  is the angle between  $\gamma$  and  $\phi$ ,  $\theta = \angle z z_0 y_0$ , then, by the First Variation Formula of Arc Length [12],  $\theta \geq \pi/2$ ; otherwise the distance from  $z$

to points on  $\gamma$  would be initially decreasing. Let  $c = d(z, z_0)$ ,  $d = d(z_0, y_0)$  and  $R = d(z, y_0)$ . In the sphere of constant curvature  $k$ , let  $z'$ ,  $z'_0$ ,  $y'_0$  be the vertices of a geodesic triangle such that  $d(z', z'_0) = d(z, z_0) = c$ ,  $d(z'_0, y'_0) = d(z_0, y_0) = d$  and  $\angle z'z'_0y'_0 = \theta$ . Let  $R' = d(z', y'_0)$ . Then by the Rauch Comparison Theorem,  $R' \leq R$ . However, it can easily be checked that on the sphere of curvature  $k$  holds  $d' < R'$ , since  $z'$ ,  $z'_0$  and  $y'_0$  are all within a strongly convex ball and  $\theta \geq \pi/2$ . Therefore  $d = d' < R' \leq R \leq \alpha + \delta$ . Thus  $\Lambda \subset B_{\alpha+\delta}(z_0)$  with  $z_0 \in \Lambda$ , which implies  $r(\Lambda) \leq \alpha + \delta$  by Proposition 18. This finishes the proof of Proposition 21.  $\square$

The proof of Theorem 11 is finished.  $\square$

## 10 Example with codifferential without closed range

For convenience, we fix the scale  $\alpha = 10$ ; any large enough value is suitable for our construction. We consider a compact metric measure space  $X$  of the following type:

As metric space, it has three cluster points  $x_\infty, y_\infty, z_\infty$  and discrete points  $(x_n)_{n \in \mathbb{N}}, (y_n)_{n \in \mathbb{N}}, (z_n)_{n \in \mathbb{N}}$  converging to  $x_\infty, y_\infty, z_\infty$ , respectively.

We set  $K_x := \{x_k : k \in \mathbb{N} \cup \{\infty\}\}$ ,  $K_y := \{y_k : k \in \mathbb{N} \cup \{\infty\}\}$ , and  $K_z := \{z_k : k \in \mathbb{N} \cup \{\infty\}\}$ . Then  $X$  is the disjoint union of the three “clusters”  $K_x, K_y, K_z$ .

We require:

$$d(x_\infty, y_\infty) = d(y_\infty, z_\infty) = \alpha \quad \text{and} \quad d(x_\infty, z_\infty) = 2\alpha.$$

We also require

$$d(x_k, y_n) < \alpha \quad \text{precisely when } n \in \{2k, 2k+1, 2k+2\}, \quad n \in \mathbb{N}, \quad k \in \mathbb{N} \cup \{\infty\},$$

$$d(z_k, y_n) < \alpha \quad \text{precisely when } n \in \{2k-1, 2k, 2k+1\}, \quad n \in \mathbb{N}, \quad k \in \mathbb{N} \cup \{\infty\}.$$

We finally require that the clusters  $K_x, K_y, K_z$  have diameter  $< \alpha$ , and that the distance between  $K_x$  and  $K_y$  as well as between  $K_z$  and  $K_y$  is  $\geq \alpha$ .

Such a configuration can easily be found in an infinite dimensional Banach space such as  $l^1(\mathbb{N})$ . For example, in  $l^1(\mathbb{N})$  consider the canonical basis vectors  $e_0, e_1, \dots$ , and set

$$x_\infty := -\alpha e_0, \quad y_\infty := 0, \quad z_\infty := \alpha e_0.$$

Define then

$$\begin{aligned} x_k &:= -\left(\alpha + \frac{1}{10k} - \frac{1}{2k} - \frac{1}{2k+1} - \frac{1}{2k+2}\right)e_0 + \frac{1}{2k}e_{2k} + \frac{1}{2k+1}e_{2k+1} + \frac{1}{2k+2}e_{2k+2}, \\ y_k &:= \frac{1}{k}e_k, \\ z_k &:= \left(\alpha + \frac{1}{10k} - \frac{1}{2k-1} - \frac{1}{2k} - \frac{1}{2k+1}\right)e_0 + \frac{1}{2k-1}e_{2k-1} + \frac{1}{2k}e_{2k} + \frac{1}{2k+1}e_{2k+1}. \end{aligned}$$

We can now give a very precise description of the open  $\alpha$ -neighborhood  $U_d$  of the diagonal in  $X^d$ . It contains all the tuples whose entries

- all belong to  $K_x \cup \{y_{2k}, y_{2k+1}, y_{2k+2}\}$  for some  $k \in \mathbb{N}$ ; or
- all belong to  $K_y \cup \{x_k, x_{k+1}, z_{k+1}\}$  for some  $k \in \mathbb{N}$ ; or
- all belong to  $K_y \cup \{x_k, z_k, z_{k+1}\}$  for some  $k \in \mathbb{N}$ ; or
- all belong to  $K_z \cup \{y_{2k-1}, y_{2k}, y_{2k+1}\}$  for some  $k \in \mathbb{N}$ .

For the closed  $\alpha$ -neighborhood, one has to add tuples whose entries all belong to  $K_y \cup \{x_\infty\}$  or to  $K_y \cup \{z_\infty\}$ .

This follows by looking at the possible intersections of  $\alpha$ -balls centered at our points.

In this topology, every set is a Borel set. We give  $x_\infty, y_\infty, z_\infty$  measure zero. When considering  $L^2$ -functions on the  $U_d$  we can therefore ignore all tuples containing one of these points.

We specify  $\mu(x_n) := \mu(z_n) := 2^{-n}$  and  $\mu(y_n) := 2^{-2^n}$ ; in this way, the total mass is finite.

We form the  $L^2$ -Alexander chain complex at scale  $\alpha$  and complement it by  $C^{-1} := \mathbb{R}^3 = \mathbb{R}x \oplus \mathbb{R}y \oplus \mathbb{R}z$ ; the three summands standing for the three clusters. The differential  $c^{-1}: C^{-1} \rightarrow L^2(X)$  is defined by  $(\alpha, \beta, \gamma) \mapsto \alpha\chi_{K_x} + \beta\chi_{K_y} + \gamma\chi_{K_z}$ , where  $\chi_{K_j}$  denotes the characteristic function of the cluster  $K_j$ .

Restriction to functions supported on  $K_x^{*+1}$  defines a bounded surjective cochain map from the  $L^2$ -Alexander complex at scale  $\alpha$  for  $X$  to the one for  $xK_x$ . Note that  $\text{diam}(K_x) < \alpha$ , consequently its Alexander complex at scale  $\alpha$  is contractible.

Looking at the long exact sequence associated to a short exact sequence of Banach cochain complexes, therefore, the cohomology of  $X$  is isomorphic (as topological vector spaces) to the cohomology of the kernel of this projection, i.e. to the cohomology of the Alexander complex of functions vanishing on  $K_x^{k+1}$ .

This can be done two more times (looking at the kernels of the restrictions to  $K_y$  and  $K_z$ ), so that finally we arrive at the chain complex  $C^*$  of  $L^2$ -functions on  $X^{k+1}$  vanishing at  $K_x^{k+1} \cup K_y^{k+1} \cup K_z^{k+1}$ .

In particular,  $C^{-1} = 0$  and  $C^0 = 0$ .

We now construct a sequence in  $C^1$  whose differentials converge in  $C^2$ , but such that the limit point does not lie in the image of  $c^1$ .

Following the above discussion, the  $\alpha$ -neighborhood of the diagonal in  $X^2$  contains in particular the “one-simplices”  $v_k := (x_k, z_k)$  and  $v'_k := (x_k, z_{k+1})$ , and their “inverses”  $\overline{v_k} := (z_k, x_k)$ ,  $\overline{v'_k} := (z_{k+1}, x_k)$ .

We define  $f_\lambda \in C^1$  with  $f_\lambda(\overline{v_k}) := f_\lambda(\overline{v'_k}) := -f_\lambda(v_k)$ ,  $f_\lambda(v'_k) := f_\lambda(v_k) := b_{\lambda,k} := 2^{\lambda k}$  and  $f_n(v) = 0$  for all other simplices.

Note that for  $0 < \lambda < 1$

$$\begin{aligned} \int_{X^2} |f|^2 &= \sum_{k=1}^{\infty} |f(v_k)|^2 \mu(v_k) + |f(v'_k)|^2 \mu(v'_k) + |f(\overline{v_k})|^2 \mu(v_k) + |f(\overline{v'_k})|^2 \mu(v'_k) \\ &= \sum_{k=1}^{\infty} 2 \cdot (2^{2\lambda k} 2^{-2k} + 2^{2\lambda k} 2^{-2k-1}) \end{aligned}$$

which is a finite sum, whereas for  $\lambda = 1$  the sum is not longer finite.

Let us now consider  $g_\lambda := c^1(f_\lambda)$ . It vanishes on all “2-simplices” (points in  $X^2$ ) except those of the form

- $d_k := (x_k, z_k, z_{k+1})$  and more generally  $d_k^\sigma := \sigma(x_k, z_k, z_{k+1})$  for  $\sigma \in S_3$  a permutation of three entries
- $e_k := (x_{k-1}, z_k, x_k)$  or more generally  $d_k^\sigma$  as before
- on degenerate simplices of the form  $(x_k, z_k, x_k)$  etc.  $g$  vanishes because  $f(x_k, z_k) = -f(z_k, x_k)$ .

We obtain

$$g_\lambda(d_k) = -f(v'_k) + f(v_k) = 0, g_\lambda(e_k) = f(\overline{v_k}) + f(v'_{k-1}) = -2^{\lambda k} + 2^{\lambda(k-1)} = 2^{\lambda k} \cdot (2^{-\lambda} - 1).$$

Similarly,  $g_\lambda(d_k^\sigma) = 0$  and  $g_\lambda(e_k^\sigma) = \text{sign}(\sigma) g_\lambda(e_k)$ .

We claim that  $g_1$ , defined with these formulas, belongs to  $L^2(X^3)$  and is the limit in  $L^2$  of  $g_\lambda$  as  $\lambda$  tends to 1.

To see this, we simply compute the  $L^2$ -norm

$$\begin{aligned} \int_{X^3} |g_1 - g_\lambda|^2 &= 6 \sum_{k=1}^{\infty} |2^{k-1} - 2^{\lambda k} (1 - 2^{-\lambda})|^2 2^{1-3k} \\ &\leq 6 \left( \sup_{k \in \mathbb{N}} 2^{-k/2} |2^{-1} - 2^{(\lambda-1)k} (1 - 2^{-\lambda})|^2 \right) \cdot \sum_{k=1}^{\infty} 2^{1-k/2} \\ &\xrightarrow{\lambda \rightarrow 1} 0 \end{aligned}$$

(the factor 6 comes from the six permutations of each non-degenerate simplex which each contribute equally).

The supremum tends to zero because each individual term does so even without the factor  $2^{-k/2}$  and the sequence is bounded.

Now we study which properties an  $f \in C^1$  with  $c^1(f) = g_1$  has to have.

Observe that for an arbitrary  $f \in C^1$ ,  $c^1 f(e_k^\sigma)$  is determined by  $f(v_k)$ ,  $f(\overline{v_k})$ ,  $f(v'_{k-1})$ ,  $f(\overline{v'_{k-1}})$  (as  $f$  vanishes on  $K_x$ ).

If  $c^1 f$  has to vanish on degenerate simplices (and this is the case for  $g_1$ ), then  $f(v_k) = -f(\overline{v_k})$  and  $f(v'_k) = -f(\overline{v'_k})$ .

$c^1 f(d_k^\sigma) = 0$  then implies that  $f(v_k) = f(v'_k)$ .

It is now immediate from the formula for  $c^1 f(d_k)$  and  $c^1 f(e_k)$  that the values of  $f$  at  $v_k$ ,  $v'_k$  are determined by  $c^1 f(d_k)$ ,  $c^1 f(e_k)$  up to addition of a constant.

Finally, observe that (in the Alexander cochain complex without growth conditions)  $f_1$  (which is not in  $L^2$ ) satisfies  $c^1(f_1) = g_1$ .

As constant functions are in  $L^2$ , we observe that  $f_1 + K$  is not in  $L^2$  for any  $K \in \mathbb{R}$ . Nor is any function  $f$  on  $X^2$  which coincides with  $f_1 + K$  on  $v_k, v'_k, \overline{v_k}, \overline{v'_k}$ .

But these are the only candidates which could satisfy  $c^1(f) = g_1$ . It follows that  $g_1$  is not in the image of  $c^1$ . On the other hand, we constructed it in such a way that it is in the closure of the image. Therefore the image is not closed.

### 10.1 A modified example where volumes of open and closed balls coincide

The example given has one drawback: although at the chosen scale  $\alpha$  open and closed balls coincide in volume (and even as sets, except for the balls around  $x_\infty, y_\infty, z_\infty$ ) for other balls this is not the case — and necessarily so, as we construct a zero-dimensional object.

We modify our example as follows, by replacing each of the points  $x_k, y_k, z_k$  by a short interval: inside  $X \times [0, 1]$ , with  $l^1$  metric (that is,  $d_Y((x, t), (y, u)) = d_X(x, y) + |t - u|$ ), consider

$$Y = \bigcup_{k \in \mathbb{N} \cup \{\infty\}} \{x_k, y_k, z_k\} \times [0, 1/(12k)].$$

For conveniency, let us write  $I_{x,k}$  for the interval  $\{x_k\} \times [0, 1/(12k)]$ , and similarly for the  $y_k$  and  $z_k$ . We then put on each of these intervals the standard Lebesgue measure weighted by a suitable factor, so that  $\mu_Y(I_{x,k}) = \mu(x_k)$ , and similarly for the  $y_k$  and  $z_k$ .

Now, if two points  $x_k, y_n$  are at distance less than  $\alpha$  in  $X$ , then they are at distance  $< \alpha - 1/k$ ; the corresponding statement holds for all other pairs of points. On the other hand, because of our choice of metric,  $d((x_k, t), (y_n, s)) \geq d(x_k, y_n)$  and again the corresponding statement holds for all other pairs of points in  $Y$ . It follows that the  $\alpha$ -neighborhood of the diagonal in  $Y^k$  is the union of products of the corresponding intervals, and exactly those intervals show up where the corresponding tuple is contained in the 1-neighborhood of the diagonal in  $X^k$ .

It is now quite hard to explicitly compute the cohomology of the  $L^2$ -Alexander cochain complex at scale  $\alpha$ .

However, we do have a projection  $Y \rightarrow X$ , namely the projection on the first coordinate. By the remark about the  $\alpha$ -neighborhoods, this projection extends to a map from the  $\alpha$ -neighborhoods of  $Y^k$  onto those of  $X^k$ , which is compatible with the projections onto the factors.

It follows that pullback of functions defines a bounded cochain map (in the reverse direction) between the  $L^2$ -Alexander cochain complexes at scale  $\alpha$ . Note that this is an isometric embedding by our choice of the measures.

This cochain map has a one-sided inverse given by integration of a function on (the  $\alpha$ -neighborhood of the diagonal in)  $Y^k$  over a product of intervals



(divided by the measure of this product) and assigning this value to the corresponding tuple in  $X^k$ . By Cauchy-Schwarz, this is bounded with norm 1.

As pullback along projections commutes with the weighted integral we are using, one checks easily that this local integration map also is a cochain map for our  $L^2$ -Alexander complexes at scale  $\alpha$ .

Consequently, the induced maps in cohomology give an isometric inclusion with inverse between the cohomology of  $X$  and of  $Y$ .

We have shown that in  $H^2(X)$  there are non-zero classes of norm 0. Their images (under pullback) are non-zero classes (because of the retraction given by the integration map) of norm 0. Therefore, the cohomology of  $Y$  is non-Hausdorff, and the first codifferential doesn't have closed image, either.

## References

- [1] M. F. Atiyah. Elliptic operators, discrete groups and von Neumann algebras. In *Colloque "Analyse et Topologie" en l'Honneur de Henri Cartan (Orsay, 1974)*, pages 43–72. Astérisque, No. 32–33. Soc. Math. France, Paris, 1976.
- [2] M. Belkin and P. Niyogi. Convergence of laplacian eigenmaps. preprint, May 2008.
- [3] M. Belkin, E. De Vito, and L. Rosasco. Random estimates of operators and their spectral properties for learning. working paper.
- [4] Raoul Bott and Loring W. Tu. *Differential forms in algebraic topology*, volume 82 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, 1982.
- [5] G. Carlsson, T. Ishkhanov, V. de Silva, and A. Zomorodian. On the local behavior of spaces of natural images. *International Journal of Computer Vision*, 76(1):1–12, 2008.
- [6] Gunnar Carlsson. Topology and data. *Bull. Amer. Math. Soc. (N.S.)*, 46(2):255–308, 2009.
- [7] Frédéric Chazal and Steve Y. Oudot. Towards persistence-based reconstruction in Euclidean spaces. In *Computational geometry (SCG'08)*, pages 232–241. ACM, New York, 2008.
- [8] Jeff Cheeger and Mikhael Gromov.  $L_2$ -cohomology and group cohomology. *Topology*, 25(2):189–215, 1986.
- [9] Fan R. K. Chung. *Spectral graph theory*, volume 92 of *CBMS Regional Conference Series in Mathematics*. Published for the Conference Board of the Mathematical Sciences, Washington, DC, 1997.
- [10] Ronald R. Coifman and Mauro Maggioni. Diffusion wavelets. *Appl. Comput. Harmon. Anal.*, 21(1):53–94, 2006.

- [11] Warren Dicks and Thomas Schick. The spectral measure of certain elements of the complex group ring of a wreath product. *Geom. Dedicata*, 93:121–137, 2002.
- [12] Manfredo Perdigão do Carmo. *Riemannian geometry*. Mathematics: Theory & Applications. Birkhäuser Boston Inc., Boston, MA, 1992. Translated from the second Portuguese edition by Francis Flaherty.
- [13] Józef Dodziuk. Finite-difference approach to the Hodge theory of harmonic forms. *Amer. J. Math.*, 98(1):79–104, 1976.
- [14] Józef Dodziuk. de Rham-Hodge theory for  $L^2$ -cohomology of infinite coverings. *Topology*, 16(2):157–165, 1977.
- [15] Józef Dodziuk, Peter Linnell, Varghese Mathai, Thomas Schick, and Stuart Yates. Approximating  $L^2$ -invariants and the Atiyah conjecture. *Comm. Pure Appl. Math.*, 56(7):839–873, 2003. Dedicated to the memory of Jürgen K. Moser.
- [16] Beno Eckmann. Harmonische Funktionen und Randwertaufgaben in einem Komplex. *Comment. Math. Helv.*, 17:240–255, 1945.
- [17] Herbert Edelsbrunner, David Letscher, and Afra Zomorodian. Topological persistence and simplification. *Discrete Comput. Geom.*, 28(4):511–533, 2002. Discrete and computational geometry and graph drawing (Columbia, SC, 2001).
- [18] Robin Forman. A user’s guide to discrete Morse theory. *Sém. Lothar. Combin.*, 48:Art. B48c, 35 pp. (electronic), 2002.
- [19] Joel Friedman. Computing Betti numbers via combinatorial Laplacians. In *Proceedings of the Twenty-eighth Annual ACM Symposium on the Theory of Computing (Philadelphia, PA, 1996)*, pages 386–391, New York, 1996. ACM.
- [20] Guy Gilboa and Stanley Osher. Nonlocal operators with applications to image processing. *Multiscale Model. Simul.*, 7(3):1005–1028, 2008.
- [21] W. V. D. Hodge. *The Theory and Applications of Harmonic Integrals*. Cambridge University Press, Cambridge, England, 1941.
- [22] X. Jiang, L-H. Lim, Y. Yao, and Y. Ye. Statistical ranking and combinatorial hodge theory. [arxiv.org/abs/0811.1067](https://arxiv.org/abs/0811.1067).
- [23] Serge Lang. *Analysis II*. Addison-Wesley, Reading, Mass, 1969.
- [24] A. B. Lee, K. S. Pedersen, and D. Mumford. The non-linear statistics of high-contrast patches in natural images. *International Journal of Computer Vision*, 54:83–103, 2003.

- [25] Peter Linnell and Thomas Schick. Finite group extensions and the Atiyah conjecture. *J. Amer. Math. Soc.*, 20(4):1003–1051 (electronic), 2007.
- [26] John Lott and Wolfgang Lück.  $L^2$ -topological invariants of 3-manifolds. *Invent. Math.*, 120(1):15–60, 1995.
- [27] W. Lück. Approximating  $L^2$ -invariants by their finite-dimensional analogues. *Geom. Funct. Anal.*, 4(4):455–481, 1994.
- [28] Wolfgang Lück.  $L^2$ -invariants: theory and applications to geometry and  $K$ -theory, volume 44 of *Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge. A Series of Modern Surveys in Mathematics [Results in Mathematics and Related Areas. 3rd Series. A Series of Modern Surveys in Mathematics]*. Springer-Verlag, Berlin, 2002.
- [29] James R. Munkres. *Elements of algebraic topology*. Addison-Wesley Publishing Company, Menlo Park, CA, 1984.
- [30] Thomas Schick. Integrality of  $L^2$ -Betti numbers. *Math. Ann.*, 317(4):727–750, 2000.
- [31] Thomas Schick.  $L^2$ -determinant class and approximation of  $L^2$ -Betti numbers. *Trans. Amer. Math. Soc.*, 353(8):3247–3265 (electronic), 2001.
- [32] S. Smale, L. Rosasco, J. Bouvrie, A. Caponnetto, and T. Poggio. Mathematics of the neural response. *Foundations of Computational Mathematics*, 2009.
- [33] Steven Smale and D-X. Zhou. Geometry on probability spaces. *Constructive Approximation*, 2009.
- [34] Edwin H. Spanier. *Algebraic topology*. Springer-Verlag, New York, 1981. Corrected reprint.
- [35] Frank W. Warner. *Foundations of differentiable manifolds and Lie groups*, volume 94 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, 1983. Corrected reprint of the 1971 edition.
- [36] Raymond Louis Wilder. *Topology of manifolds*, volume 32 of *American Mathematical Society Colloquium Publications*. American Mathematical Society, Providence, R.I., 1979. Reprint of 1963 edition.
- [37] D. Zhou and B. Schölkopf. Regularization on discrete spaces. In *Pattern Recognition*, Proc. 27th DAGM Symposium, pages 361–368, Berlin, 2005.
- [38] Afra Zomorodian and Gunnar Carlsson. Computing persistent homology. *Discrete Comput. Geom.*, 33(2):249–274, 2005.

## A An example of a space with infinite dimensional $\alpha$ -scale homology

Appendix written by

Anthony W. Baker<sup>1</sup>  
Mathematics and Computing Technology  
The Boeing Company

The work in in the main body of this paper has inspired the question of the existence of a separable, compact metric space with infinite dimensional  $\alpha$ -scale homology. This appendix provides one such example and further shows the sensitivity of the homology to changes in  $\alpha$ .

Let  $X$  be a separable, complete metric space with metric  $d$ , and  $\alpha > 0$  a “scale”. Define an associated (generally infinite) simplicial complex  $C_{X,\alpha}$  to  $(X, d, \alpha)$ . Let  $X^{\ell+1}$ , for  $\ell > 0$ , be the  $(\ell+1)$ -fold Cartesian product, with metric denoted by  $d, d : X^{\ell+1} \times X^{\ell+1} \rightarrow \mathbb{R}$  where  $d(x; y) = \max_{i=0, \dots, \ell} d(x_i; y_i)$ . Let

$$U_{\alpha}^{\ell+1}(X) = U_{\alpha}^{\ell+1} = \{x \in X^{\ell+1} : d(x; D^{\ell+1}) \leq \alpha\}$$

where  $D^{\ell+1} \subset X^{\ell+1}$  is the diagonal, so  $D^{\ell+1} = \{t \in X : (t, \dots, t), \ell+1 \text{ times}\}$ . Let  $C_{X,\alpha} = \cup_{\ell=0}^{\infty} U_{\alpha}^{\ell+1}$ . This has the structure of a simplicial complex whose  $\ell$  simplices consist of points of  $U_{\alpha}^{\ell+1}$ .

The  $\alpha$ -scale homology is that homology generated by the simplicial complex above.

The original exploration of example compact metric spaces resulted in low dimensional  $\alpha$ -scale homology groups. Missing from the results were any examples with infinite dimensional homology groups. In addition examination of the first  $\alpha$ -scale homology group was less promising for infinite dimensional results; the examination resulted in the proof that the first homology group is always finite, as shown in Section 9.

The infinite dimensional example in this paper was derived through several failed attempts to prove that the  $\alpha$ -scale homology was finite. The difficulty that presented itself was the inability to slightly perturb vertices and still have the perturbed object remain a simplex. This sensitivity is derived from the “equality” in the definition of  $U_{\alpha}^{\ell+1}$ . It is interesting to note the contrast between the first homology group and higher homology groups. In the case of first homology group all 1-cycles can be represented by relatively short simplices; there is no equivalent statement for higher homology groups:

**Lemma 5.** *A 1-cycle in  $\alpha$ -scale homology can be represented by simplices with length less than or equal to  $\alpha$ .*

*Proof.* For any  $[x_i, x_j]$  simplex with length greater than  $\alpha$  there exists a point  $p$  such that  $d(x_i, p) \leq \alpha$  and  $d(x_j, p) \leq \alpha$ . This indicates  $[x_i, p]$ ,  $[p, x_j]$ , and  $[x_i, p, x_j]$  are simplices. Since  $[x_i, p, x_j]$  is a simplex we can substitute  $[x_i, p] + [p, x_j]$  for  $[x_i, x_j]$  and remain in the original equivalence class.  $\square$

---

<sup>1</sup>anthony.w.baker@boeing.com

In the section that follows we present an example that relies on the rigid nature of the definition to produce an infinite dimensional homology group. The example is a countable set of points in  $\mathbb{R}^3$ . One noteworthy point is that from this example it is easy to construct a 1-manifold embedded in  $\mathbb{R}^3$  with infinite  $\alpha$ -scale homology. In addition to showing that for a fixed  $\alpha$  the homology is infinite, we consider the sensitivity of the result around that fixed  $\alpha$ .

The existence of an infinite dimensional example leads to the following question for future consideration: are there necessary and sufficient conditions on  $(X, d)$  for the  $\alpha$ -scale homology to be finite.


### A.1 An Infinite Dimensional Example

The following example illustrates a space such that the second homology group is infinite. For the discussion below fix  $\alpha = 1$ .

Consider the set of point  $\{A, B, C, D\}$  in the diagram below such that

$$d(A, B) = d(B, C) = d(C, D) = d(A, D) = 1$$

$$d(A, C) = d(B, D) = \sqrt{2}$$



The lines in the diagram suggest the correct structure of the  $\alpha$ -simplices for  $\alpha = 1$ . The 1-simplices are  $\{\{A, B\}, \{B, C\}, \{C, D\}, \{A, D\}, \{A, C\}, \{B, D\}\}$ . The 2-simplices are the faces  $\{\{A, B, C\}, \{A, B, D\}, \{A, C, D\}, \{B, C, D\}\}$ . There are no (non-degenerate) 3-simplices. A 3-simplex would imply a point such that all of the points are within  $\alpha = 1$  — no such point exists. The chain  $[ABC] - [ABD] + [ACD] - [BCD]$  is a cycle with no boundary.

Define  $R$  as  $R = \{r \in [0, 1, 1/2, 1/3, \dots]\}$ . Note in this example  $R$  acts as an index set and the dimension of the homology is shown to be at least that of  $R$ .

Let  $X = \{A, B, C, D\} \times R$ . Define  $A_r = (A, r)$ ,  $B_r = (B, r)$ ,  $C_r = (C, r)$ , and  $D_r = (D, r)$ .

We can again enumerate the 1-simplices for  $X$ . Let  $r, s, t, u \in R$ . The 1-simplices are

$$\begin{aligned} & \{\{A_r, B_s\}, \{B_r, C_s\}, \{C_r, D_s\}, \{A_r, D_s\}, \\ & \{A_r, A_s\}, \{B_r, B_s\}, \{C_r, C_s\}, \{D_r, D_s\}, \\ & \{\mathbf{B}_r, \mathbf{D}_r\}, \{\mathbf{A}_r, \mathbf{C}_r\}\}. \end{aligned}$$

The last two 1-simplices (highlighted) must have the same index in  $R$  due to the distance constraint.

The 2-simplices are

$$\begin{aligned} & \{\{A_r, B_s, C_r\}, \{A_s, B_r, D_r\}, \{A_r, C_r, D_s\}, \{B_r, C_s, D_r\}, \\ & \{A_r, B_s, B_r\}, \{B_r, C_s, C_r\}, \{C_r, D_s, D_r\}, \{A_r, D_s, D_r\}, \\ & \{A_s, A_r, B_s\}, \{B_s, B_r, C_s\}, \{C_s, C_r, D_s\}, \{A_r, A_r, D_s\}, \\ & \{A_r, A_s, A_t\}, \{B_r, B_s, B_t\}, \{C_r, C_s, C_t\}, \{D_r, D_s, D_t\}\}. \end{aligned}$$

The 3-simplices are

$$\begin{aligned} & \{\{A_r, B_s, B_t, C_r\}, \{A_s, A_t, B_r, D_r\}, \{A_r, C_r, D_s, D_t\}, \{B_r, C_s, C_t, D_r\}, \\ & \{A_r, B_t, B_s, B_r\}, \{B_r, C_t, C_s, C_r\}, \{C_r, D_t, D_s, D_r\}, \{A_r, D_t, D_s, D_r\}, \\ & \{A_t, A_s, A_r, B_s\}, \{B_t, B_s, B_r, C_s\}, \{C_t, C_s, C_r, D_s\}, \{A_t, A_r, A_r, D_s\}, \\ & \{A_r, A_s, A_t, A_u\}, \{B_r, B_s, B_t, B_u\}, \{C_r, C_s, C_t, C_u\}, \{D_r, D_s, D_t, D_u\}\}. \end{aligned}$$

Define  $\sigma_r := [A_r B_r C_r] - [A_r B_r D_r] + [A_r C_r D_r] - [B_r C_r D_r]$ . By calculation,  $\sigma_r$  is shown to be a cycle. Suppose that there existed a chain of 3-simplices such that the  $\sigma_r$  is the boundary then  $\gamma = [A_r A_s B_r D_r]$  must be included in the chain to eliminate  $[A_r B_r D_r]$ . Since the boundary of  $\gamma$  contains  $[A_s B_r D_r]$  there must be a term to eliminate this term. The only term with such a boundary is of the form  $[A_s A_t B_r D_r]$ . Again, a new simplex to eliminate the extra boundary term is in the same form. Either this goes on *ad infinitum*, impossible since the chain is finite, or it returns to  $A_r$  in which case the boundary of the original chain is 0 (contradicting that the  $[A_r B_r D_r]$  term is eliminated). For all  $r \in R$ ,  $\sigma_r$  is a generator for homology.

If  $s \neq t$  then  $\sigma_s$  and  $\sigma_t$  are not in the same equivalence class. Suppose they are. The same argument above shows that any term with the face  $[A_t B_t D_t]$  will necessarily have a face  $[A_u B_t D_t]$  for some  $u \in R$ . Such a term needs to be eliminated since it cannot be in the final image but such an elimination would cause another such term or cancel out the  $[A_t B_t D_t]$ . In either case the chain would not satisfy the boundary condition necessary to equivalence  $\sigma_s$  and  $\sigma_t$  together.

Each  $\sigma_s$  is a generator of homology and, therefore, the dimension of the homology is at least the cardinality of  $R$  which in this case is infinite.

**Theorem 12.** *For  $\alpha = 1$ , the second  $\alpha$ -scale homology group for*

$$X = \{A, B, C, D\} \times R$$

*is infinite dimensional.*

## A.2 Consideration for $\alpha < 1$

The example above is tailored for scale  $\alpha = 1$ . In this metric space the nature of the second  $\alpha$ -scale homology group changes significantly as  $\alpha$  changes.

Consider when  $\alpha$  falls below one. In this case the structure of the simplices collapses to simplices restricted to a line (with simplices of the form  $\{\{A_r, A_s, A_t\}, \{B_r, B_s, B_t\}, \{C_r, C_s, C_t\}, \{D_r, D_s, D_t\}\}$ ). These are clearly degenerate simplices resulting in a trivial second homology group.

In this example the homology was significantly reduced as  $\alpha$  decreased. This is not necessarily always the case. The above example could be further enhanced by replicating smaller versions of itself in a fractal-like manor so that as  $\alpha$  decreases the  $\alpha$ -scale homology encounters many values with infinite dimensional homology.

## A.3 Consideration for $\alpha > 1$

There are two cases to consider when  $\alpha > 1$ . The first is the behavior for very large  $\alpha$  values. In this case the problem becomes simple as illustrated by the lemma below.

Define  $\alpha$  **large** with respect to  $d$  if there exists an  $\rho \in X$  such that  $d(\rho, y) \leq \alpha$  for all  $y \in X$ .

**Lemma 6.** *Let  $X$  be a separable, compact metric space with metric  $d$ . If  $\alpha$  is large with respect to  $d$  then the  $\alpha$ -scale homology of  $X$  is trivial.*

*Proof.* Let  $\rho \in X$  satisfy the attribute above. Then  $U_\alpha^{\ell+1} = X^{\ell+1}$  since

$$d((\rho, \dots, \rho), (x_0, x_1, \dots, x_\ell)) \leq \alpha$$

for all values of  $x_i$ .

Let  $\sigma = \sum_{j=1}^k c_j(a_0^j, a_1^j, \dots, a_n^j)$  be an  $n$ -cycle. Define

$$\sigma_\rho = \sum_{j=1, k} c_j(a_0^j, a_1^j, \dots, a_n^j, \rho).$$

The  $n+1$ -cycle,  $\sigma_\rho$ , acts as a cone with base  $\sigma$ . Since  $\sigma$  is a cycle the terms in the boundary of  $\sigma_\rho$  containing  $\rho$  cancel each other out to produce zero. The terms without  $\rho$  are exactly the original  $\sigma$ . Therefore there exists no cycles without boundaries. This proves that for  $\alpha$  large and  $X$  infinite the homology of  $X$  is trivial.  $\square$

In the case that  $\alpha > 1$  but still close to 1, the second homology group changes significantly but does not completely disappear. In the example, simplices that existed only by the equality in the definition of  $\alpha$ -scale homology when  $\alpha = 1$  now find neighboring 2-simplices joined by higher dimensional 3-simplices. The result is larger equivalence classes of 2-cycles. This reduces the infinite dimensional homology for  $\alpha = 1$  to a finite dimension for  $\alpha$  slightly larger than 1. As  $\alpha$  gets closer to 1 from above the dimension of the homology increases without bound.

It is interesting to note that the infinite characteristics for  $\alpha = 1$  are tied heavily to the fact that the simplices that determined the structure lived on the bounds of being simplices. As  $\alpha$  changes from 1, the rigid restrictions on the simplices is no longer present in this example. The result is a significant reduction in the dimension of the homology.