THE INTEGRAL NOVIKOV CONJECTURE FOR ASYMPTOTICALLY FINITE DIMENSIONAL GROUPS

A.N. DRANISHNIKOV, S. FERRY, AND S, WEINBERGER

ABSTRACT. We show that the mod p Higson conjecture (for all p) for the universal cover of a geometrically finite group Γ implies the integral Novikov conjecture for Γ . Then we prove the mod p Higson conjecture for such groups with asdim $\Gamma < \infty$.

§1 INTRODUCTION

Coarse approach to the Novikov Higher Signatures Conjecture proved to be very productive. There are many statements on coarse geometry of a discrete group Γ that imply the Novikov Conjecture. One of such statements called the Higson Conjecture has a very short history because it turns out that it is false even in the cases where the Novikov conjecture is proven. One of the purposes of this paper is to revive the Higson Conjecture and to apply it for proving the integral Novikov conjecture for groups with finite asymptotic dimension.

We assume here that our discrete group Γ has a finite Eilenberg-MacLane complex $K(\Gamma, 1)$ also denoted by $B\Gamma$. Our method works in the case when the *n*-skeleton $B\Gamma^{(n)}$ is finite for every *n* but we do not address it in this paper. The universal covering space $X = E\Gamma$ of $B\Gamma$ admits an equivariant metric lifted from a metric on $B\Gamma$. Higson defined a certain compactification \bar{X} of any metric space X (called the Higson compactification) and proposed the following [Ro1]:

CONJECTURE. The Higson compactification \bar{X} of the above universal covering space X is rationally acyclic: $\tilde{H}^*(\bar{X}; \mathbf{Q}) = 0$.

1

Typeset by $\mathcal{AMS-TEX}$

¹⁹⁹¹ Mathematics Subject Classification. Primary 20H15.

 $Key\ words\ and\ phrases.$ dimension, asymptotic dimension, absolute extensor, Higson corona, Novikov conjecture..

The authors were partially supported by NSF grants DMS-0305152, DMS-, and DMS -.

We recall the definition of the Higson compactification. Let C_h be the set of all bounded continuous functions $f: X \to \mathbf{R}$ having the following property:

(*)
$$\lim_{d(x,x_0)\to\infty} diam(f(B_r(x))) = 0$$

for every r > 0 where $B_r(x)$ denotes the closed r-ball centered in $x \in X$ and $x_0 \in X$ is a fixed point. Clearly, this condition does not depend on the choice of x_0 . Higson defined his compactification as the maxiamal ideal space of the algebra C_h [Ro1]. It admits the following more topological decription. The Higson compactification of X is the closure \bar{X} in \mathbf{R}^{C_h} of the image of X under the imbedding $F: X \to \mathbf{R}^{C_h}$ defined by the formula $F(x) = \{f(x)\}_{f \in C_h}$. Since every function $f \in C_h$ is bounded, the image F(X) lies in a compact space

$$\prod_{f \in C_h} [\inf(f), sup(f)] \subset \mathbf{R}^{C_h}$$

and hence its closure is compact. The reminder $\overline{X} \setminus X$ is denoted by νX and it is called the Higson corona.

Different description of the Higson compactification are given in §2 and §3. We note that the Higson compactification plays a central role in connecting the bounded topology with the continuously controlled topology.

It was shown in [Ro1] that the Higson Conjecture implies the classical Novikov Conjecture. Soon after Keesling noticed that the Higson Conjecture does not hold for $\Gamma = \mathbf{Z}$ [Kee]. An example of a nontrivial (integral) cohomology element of the Higson compactification $\mathbf{\bar{R}}$ of $E\Gamma = \mathbf{R}$ can be given by means of a function $f : \mathbf{R} \to S^1$, $f(t) = e^{i\sqrt{t}}$. Since this function satisfies the condition (*), there is a continuous extension $\bar{f} : \mathbf{\bar{R}} \to S^1$. Clearly, this cohomology element has infinite order.

An important step in correction of the Higson conjecture is the following remark We note that Keesling's element is *p*-divisible for all *p*. Indeed, if $\phi : S^1 \to S^1$ is a *p*-fold covering, $z \xrightarrow{\phi} z^p$, the function \overline{f} admits a lift defined as an extension of the function $g: \mathbf{R} \to S^1, g(t) = e^{(i/p)\sqrt{t}}$, to the Higson corona. Thus, the following form of the Higson conjecture cannot be eliminated by Keesling's example.

Mod p Higson conjecture. If X is the universal covering of a finite aspherical complex supplied with a lifted metric and \bar{X} is its Higson compactification, then $\tilde{H}^*(\bar{X}; \mathbf{Z}_p) = 0$ for the prime p.

Our first result is the following

Theorem A. Suppose that a group Γ admits a finite complex for a classifying space $B\Gamma$ and the Mod p Higson conjecture holds for $E\Gamma$ for all primes p. Assume that dim $\nu\Gamma < \infty$. Then the integral Novikov Conjecture holds for Γ .

Here under the integral Novikov Conjecture for a group Γ we understand the statement that the assembly map

$$H_*(B\Gamma; \mathbf{L}) \xrightarrow{A} L_*(\mathbf{Z}\Gamma)$$

is a monomorphism. We recall that the classical Novikov Conjecture for Γ is equivalent to the statement that the rational assembly map

$$A \otimes 1_{\mathbf{Q}} : H_*(B\Gamma; \mathbf{L}) \otimes \mathbf{Q} \to L_*(\mathbf{Z}\Gamma) \otimes \mathbf{Q}$$

is a monomorphism [FRR].

The proof of Theorem A is based on the mod p version of a Carlsson-Pedersen theorem from [CP1].

Our next result states that groups with finite asymptotic dimension satisfy the Mod p Higson conjecture. The asymptotic dimension of a metric space was defined by Gromov [G1]. Here discrete finitely generated groups are considered as metric spaces supplied with the word metric. Since asdim $\mathbf{R}^n = n$, this results implies that all nontrivial *n*-cohomology of the Higson compactification of \mathbf{R}^n constructed in [DF] are *p*-divisible for all p.

Thus, together with Theorem A and the inequality $\dim \nu \Gamma \leq \operatorname{asdim} \Gamma$ [DKU] this result leads to the following.

Theorem B. Suppose that a group Γ admits a finite complex for a classifying space $B\Gamma$ and asdim $\Gamma < \infty$. Then the integral Novikov conjecture holds for Γ .

We note that the rational Novikov Conjecture for such groups Γ was proven first by Guoliang Yu [Yu1]. The integral K-theoretical Novikov Conjecture for asymptotically finite dimensional groups was proved recently by Carlsson and Goldfarb [CG2]. The class of discrete groups Γ with asdim $\Gamma < \infty$ is very large. It contains hyperbolic groups, polycyclic groups, Coxeter groups, arithmetic groups, etc. (see [G1],[Ro2], [BD1], [DJ], [Ji], [CG1]). In fact it is not easy to construct a finitely presented group with infinite asymptotic dimension. It is especially difficult in the class of groups with finite cohomological dimension. Only recently using expanders Gromov constructed asymptotically infinite dimensional groups with finite $B\Gamma$ [G2].

Theorem B shows that the Mod p Higson conjecture has a great potential. In addition to that we prove the following.

Theorem C. Suppose that a group Γ admits a finite complex for a classifying space $B\Gamma$ and the Mod 2 Higson conjecture holds for $E\Gamma$. Then the Novikov Higher Signatures conjecture holds for Γ .

$\S2$ L-theory and assembly map with \mathbf{Z}_p coefficients

For every ring with involution R (actually any additive category with involution) Ranicki defines a 4-periodic spectrum $\mathbf{L}_*(R)$ with $\pi_i(\mathbf{L}_*(R)) = L_i(R)$ [Ran1]. We use the notation $\mathbf{L} = \mathbf{L}_*(\mathbf{Z})$.

Below R is a PID with involution.

For a metric space (X, d) by $C_X(R)$ we denote the boundedly controlled Pedersen-Weibel category whose objects are locally finite direct sums $A = \bigoplus_{x \in X} A(x)$ of finite dimensional free *R*-modules and morphisms are given by matrices with bounded propagations. For a subset $V \subset X$ we denote by A(V) the sum $\bigoplus_{x \in V} A(x)$. Ranicki [Ran2] defined *X*-bounded quadratic *L*-groups $L_*(C_X(R))$ and the corresponding spectrum $\mathbf{L}_*(C_X(R))$. We will use a notation $\mathbf{L}^{bdd}(X) = \mathbf{L}_*(C_X(\mathbf{Z}))$.

Suppose that \bar{X} is a compactification of X with the corona $Y = \bar{X} \setminus X$. Then one can define continuously controlled category $B_{X,Y}(R)$ by taking the same objects as above and with morphisms $f : A \to B$ to be homomorphisms such that for every $y \in Y$ and every neighborhood $U \subset \bar{X}$ there is a smaller neighborhood $V \subset U$ such that $f(A(V)) \subset A(U)$. This category is additive and hence also admits the L-theory. The corresponding spectrum for $R = \mathbb{Z}$ we denote by $\mathbf{L}^{cc}(X) = \mathbf{L}_*(B_{X,Y}(\mathbb{Z}))$.

For a spectrum $\mathbf{E} = \{E_k \mid k \in \mathbf{Z}\}$ by $H_i(X; \mathbf{E})$ we denote the **E**-homology of X. If X is a complex, then $H_i(X; \mathbf{E}) = \pi_i(X_+ \wedge \mathbf{E}) = \lim_{k \to \infty} \pi_{i+k}(X_+ \wedge E_k)$. If X is a compact metric space, by $H_i(X; \mathbf{E})$ we denote the Steenrod **E**-homology of X, i.e. $H_i(X; \mathbf{E}) = \pi_i(\operatorname{holim}\{N^i_+ \wedge \mathbf{E}\})$ where $X = \lim_{\leftarrow} N^i$ is the inverse limit of polyhedra [CP2], [EH].

The following theorem was proven in [P], [CP1], [CP2].

Theorem 2.1. $L_i(B_{X,Y}(R)) = H_{i-1}(Y; \mathbf{L}(R))$ for all *i* where the homology is the Steenrod $\mathbf{L}_*(R)$ -homology.

Subsets $A, B \subset X$ of a metric space (X, d) are called *diverging* if

$$\lim_{r \to \infty} d(A \setminus B_r(x_0), B \setminus B_r(x_0)) = \infty$$

where $x_0 \in X$ is any point, $B_r(x_0)$ is the *r*-ball centered at x_0 , and the distance d(A', B')between sets is the infimum of distances between d(a, b), $a \in A'$, $b \in B'$. We note that the minimal compactification of X with respect to the property that the closures of every diverging subsets in X have empty intersection in the corona is the Higson compactification. A compactification \tilde{X} of X is called Higson dominated if the identity map $1_X : X \to X$ admits a continuous extension $\bar{X} \to \tilde{X}$. We note that for every Higson dominated compactification of X with the corona Y there is a forgetful functor $C_X(R) \to B_{X,Y}(R)$. This functor defines a forgetful map of spectra $\phi : \mathbf{L}^{bdd}(X) \to \mathbf{L}^{cc}(X)$.

If X is a universal covering of a finite complex with the fundamental group Γ , then Γ acts on $C_X(R)$ and hence on $\mathbf{L}_*(C_X(R))$ with the fixed set $\mathbf{L}_*(C_X(R))^{\Gamma} = L_*(R\Gamma)$ [CP]. We note that the base points of $\mathbf{L}_*(C_X(R))$ are fixed under Γ -action. If a compactification of X with the corona Y is equivariant, then Γ acts on $B_{X,Y}(R)$ and hence on $\mathbf{L}_*(B_{X,Y}(R))$.

We recall that the homotopy fixed set $X^{h\Gamma}$ of a pointed space X with a Γ action on it is defined as the space of equivariant maps $Map_{\Gamma}(E\Gamma_{+}, X)$.

For a general spectrum **E** and a torsion free group Γ it was proven [C], [CP1] that

$$\mathbf{H}_*(B\Gamma; \mathbf{E}) \cong \mathbf{H}^{lf}_*(E\Gamma; \mathbf{E})^{\Gamma} \cong \mathbf{H}^{lf}_*(E\Gamma; \mathbf{E})^{h\Gamma}.$$

Here we use notation $\mathbf{H}_*(X; \mathbf{E})$ for the spectrum \mathbf{S} with $\pi_i(\mathbf{S}) = H_i(X; \mathbf{E})$ and $\mathbf{H}_*^{lf}(X; \mathbf{E})$ for the spectrum \mathbf{S}' with $\pi_i(\mathbf{S}) = H_i^{lf}(X; \mathbf{E})$. Formally, $H_i^{lf}(X; \mathbf{E})$ is defined for a locally compact space X as the Steenrod \mathbf{E} -homology of the one point compactification αX . We note that for a complex N, $\mathbf{H}_*(N; \mathbf{E}) = N_+ \wedge \mathbf{E}$ and for a compact space $Y, \mathbf{H}_*(Y; \mathbf{E}) = \text{holim}\{N_+^{\alpha} \wedge \mathbf{E}\}$ where N^{α} runs over nerves of all finite open covers of Y. If Γ acts on a compact space Y, then it acts on the set of all finite open covers of Y and hence on the spectrum $\mathbf{H}_*(Y; \mathbf{E})$.

The following theorem is a combination of results of Carlsson-Pedersen [CP1],[CP2] and Ranicki [Ran3]. The first part of it in a more general setting is discussed in [Ros], Theorem 3.3.

Theorem 2.2.

For every group Γ with finite classifying complex $B\Gamma$ there are morphism of spectra called the bounded control assembly map

$$A^{bdd}: \mathbf{H}^{lf}_{*}(E\Gamma; \mathbf{L}) \to \mathbf{L}^{bdd}(E\Gamma)$$

which close a homotopy commutative diagram

$$\begin{array}{cccc}
\mathbf{H}_{*}(B\Gamma; \mathbf{L}) & \xrightarrow{A} & \mathbf{L}_{*}(\mathbf{Z}\Gamma) \\
\simeq & & trf \\
\mathbf{H}_{*}^{lf}(E\Gamma; \mathbf{L})^{h\Gamma} & \xrightarrow{A^{bdd,h\Gamma}} & \mathbf{L}^{bdd}(E\Gamma)^{h\Gamma}
\end{array}$$

where A is the standard assembly map, the vertical arrows are the natural maps from fixed sets to homotopy fixed sets.

If the universal covering space $E\Gamma$ admits a Higson dominated equivariant compactification X with the corona Y then the diagram can be extended

where $A^{cc,h\Gamma} = \phi^{h\Gamma} \circ A^{bdd,h\Gamma}$ is the composition with the forgetful map. Moreover, A^{cc} coincides with the boundary map from the Steenrod L-homology exact sequence of pair (X,Y):

$$A_p^{cc} = \partial : \mathbf{H}_*^{lf}(E\Gamma; \mathbf{L}) \to \mathbf{H}_{*-1}(Y; \mathbf{L}).$$

Let M(p) denote the Moore spectrum for the group \mathbf{Z}_p .

Lemma 2.1. Let X be a finite-dimensional compact metric space. Suppose that X is \mathbf{Z}_p -acyclic, i.e., $\tilde{H}^*(X; \mathbf{Z}_p) = 0$ for the reduced Čech cohomology. Then X is acyclic for the reduced Steenrod $\mathbf{L} \wedge M(p^k)$ -homology for all k.

Proof. First we note $\tilde{H}_*(X; \mathbf{Z}_p) = 0$ for the reduced Steenrod homology. Using coefficient exact sequence we can show that $\tilde{H}_*(X; \mathbf{Z}_{p^k}) = 0$ for all k. For every spectrum **S** there is the Steenrod homology Atiyah-Hirzebruch spectral sequence with

$$E_{i,j}^2 = \tilde{H}_i(X; H_j(S^0, \mathbf{S}))$$

which converges to $H_*(X; \mathbf{S})$ provided X is finite dimensional compact metric space [EH],[KS]. Since all groups $H_q(S^0, \mathbf{L} \wedge M(p^k))$ are \mathbf{Z}_{p^s} , we have $E_{i,j}^2 = 0$ for all i and j. \Box

REMARK 1. The finite dimensionality condition is essential here. Tere are acyclic compacta that have nontrivial mod p complex K-theory [T]. Namely, by results of Adams and Toda there is a map $f : \Sigma^k M(\mathbf{Z}_p, m) \to M(\mathbf{Z}_p, m)$ to a Moore space of the suspension of the Moore space which induces an isomorphism in K-theory. Then the inverse limit

THE INTEGRAL NOVIKOV CONJECTURE FOR ASYMPTOTICALLY FINITE DIMENSIONAL GROUPS7

of the suspensions of f is a an acyclic compactum, since all bonding maps are trivial in cohomology, and it has nontrivial K-theory and nontrivial mod p K-theory.

REMARK 2. The spectrum \mathbf{L} localaized at 2 is equivalent to the Eilenberg-MacLane 4-periodic spectrum generated by the space [MM]

$$\prod_{i=1}^{\infty} K(\mathbf{Z}_{(2)}, 4i) \times K(\mathbf{Z}_2, 4i-2).$$

Thus, $\mathbf{L} \wedge M(2) = \mathbf{L}_{(2)} \wedge M(2)$ is the Eilenberg-MacLane spectrum generated by the space

$$\prod_{i=1}^{\infty} K(\mathbf{Z}_2, 4i) \times K(\mathbf{Z}_2, 4i-1) \times K(\mathbf{Z}_2, 4i-2).$$

Now if compactum X is \mathbb{Z}_2 -acyclic it is $L \wedge M(2)$ -acyclic without assumption of finite dimensionality of X.

DEFINITION. The mod p assembly map A_p is by the definition

$$A \wedge 1_{M(p)} : \mathbf{H}_*(B\Gamma; \mathbf{L}) \wedge M(p) \to \mathbf{L}_*(\mathbf{Z}\Gamma) \wedge M(p).$$

Theorem 2.3. Suppose that the universal cover X of a finite aspherical complex B satisfies the mod p Higson conjecture and let $\dim \nu X < \infty$. Then the mod p^k assembly map for $\Gamma = \pi_1(B)$ is a split monomorphism for every k.

Proof. Let let $m = \dim \nu X$. Using Schepin's spectral theorem [Dr1] one can obtain a metrizable \mathbb{Z}_p -acyclic Γ -equivariant compactification \overline{X} of X with the corona Y of dimension dim Y = m (see the proof of Lemma 8.3. in [Dr1]). We introduce coefficients to the second diagram of Theorem 2.2 by the smash product with $M(p^k)$.

If Γ acts on a based space W such that the base point is free, there is the formula $(W \wedge N)^{h\Gamma} = W^{h\Gamma} \wedge N$ for the trivial action on N. Therefore, we have the commutative diagram

We show that

$$A^{cc} \wedge 1 : \mathbf{H}^{lf}_{*}(X; \mathbf{L}) \wedge M(p^{k}) \to \mathbf{L}^{cc}(X) \wedge M(p^{k})$$

is an isomorphism. This would imply that $(A^{cc} \wedge 1)^{h\Gamma}$ is an isomorphism and hence A_p is a split monomorphism.

In view of Theorems 2.1 and 2.2 we need to show that

$$\partial \wedge 1 : \mathbf{H}^{lf}_{*}(X; \mathbf{L}) \wedge M(p^{k}) \to \mathbf{H}_{*-1}(Y; \mathbf{L}) \wedge M(p^{k})$$

is isomorphism. We note that $\partial \wedge 1_{M(p^k)}$ is equivalent to the boundary homomorphism for the pair (\bar{X}, Y) in $\mathbf{L} \wedge M(p^k)$ -homology. By Lemma 2.1 we obtain $\mathbf{H}_*(\bar{X}; \mathbf{L} \wedge M(p^k)) =$ $\mathbf{H}_*(pt; \mathbf{L} \wedge M(p^k))$ and hence $\partial \wedge 1_{M(p^k)}$ is an isomorphism. \Box

It is known that the universal coefficient formula with \mathbf{Z}_p coefficients (UCF) holds true for every generalized homology theory and it is natural with respect to morphisms of spectra.

Proposition 2.1. For every morphism of spectra $A : \mathbf{E}_1 \to \mathbf{E}_2$ and every p and i there is a commutative diagram

Proof. We apply the smash product with \mathbf{E}_i , i = 1, 2 to the cofibration of spectra $\mathbf{S} \to \mathbf{S} \to M(p)$. Then the result follows from the homotopy exact sequence of the resulting cofibrations of spectra and the induced morphism between them



_

Theorem 2.4. Suppose that the mod p Higson conjecture holds for the universal cover of finite complex $B\Gamma$ for all p and let dim $\nu\Gamma < \infty$. Then the integral assembly map A is a monomorphism.

Proof. In view of compactness of $B\Gamma$ we have $\nu\Gamma = \nu E\Gamma$. By Theorem 2.3 $A \otimes 1_G$ is a split monomorphism for every finite abelian group G. Since $B\Gamma$ is a finite complex,

the standard induction argument on the number of cells show that the group $H_i(M; \mathbf{L})$ is finitely generated for every i. Hence for every $\alpha \in H_i(M; \mathbf{L})$ there is p such that $\alpha \otimes 1 \in H_i(M; \mathbf{L}) \otimes \mathbf{Z}_p$ is not zero. By the UCF there is a monomorphism $H_i(M; \mathbf{L}) \otimes \mathbf{Z}_p \to$ $H_i(M; \mathbf{L}(\mathbf{Z}_p))$ which together with the assembly map produces a commutative diagram

$$H_{i}(M; \mathbf{L}) \otimes \mathbf{Z}_{p} \longrightarrow H_{i}(M; \mathbf{L} \wedge M(p))$$

$$A \otimes 1 \downarrow \qquad A_{p} \downarrow$$

$$L_{i}(\mathbf{Z}\Gamma) \otimes \mathbf{Z}_{p} \longrightarrow \pi_{i}(\mathbf{L}(\mathbf{Z}\Gamma) \wedge M(p)).$$

This diagram implies that $A(\alpha) \neq 0$. \Box

Theorem 2.5. Suppose that a group Γ admits a finite complex for a classifying space $B\Gamma$ and the Mod 2 Higson conjecture holds for $E\Gamma$. Then the Novikov Higher Signatures conjecture holds for Γ .

Proof. In view of Remark 2 the argument of Theorem 2.3 for p = 2 works without the assumption dim $E\gamma < \infty$. Hence the mod 2^k assembly map

$$A \wedge 1_{M(2^k)} : \mathbf{H}_*(B\Gamma; \mathbf{L}) \wedge M(2^k) \to \mathbf{L}_*(\mathbf{Z}\Gamma) \wedge M(2^k)$$

is a split monomorphism. On the group level by the Universal Coefficient Formula we have the diagram

Since the group $H_i(B\Gamma; \mathbf{L})$ is finitely generated, it can be presented as $\bigoplus_{F_i} \mathbf{Z} \oplus Tor_i$. Since $A_* \otimes \mathbf{1}_{\mathbf{Z}_{2^k}}$ is a monomorphism, the kernel $ker(A_*)$ consists of 2^k divisible elements. Since k is arbitrary, $ker(A_*|_{\oplus \mathbf{Z}}) = 0$. Therefore, $A_* \otimes \mathbf{1}_{\mathbf{Q}}$ is a monomorphism. \Box

$\S3 \mod p$ Higson conjecture for asymptotically finite dimensional sapces

We recall that a map $f: X \to Y$ between metric spaces is λ -Lipschitz if $d_Y(f(x), f(x')) \leq \lambda d_X(x, x')$ for all $x, x' \in X$. Denote by

$$L(f) = \sup_{x \neq x'} \{ \frac{d_Y(f(x), f(x'))}{d_X(x, x')} \}$$

the minimal Lipschitz constant of f.

Every simplicial complex K carries a metric where all simplexes are the standard euclidean of size one. We will call the maximal such metric *uniform* and usually we will denote corresponding metric space as K_U . Note that the metric space K_U is geodesic If it is not specified, we will assume that a finite complex always supplied with the uniform metrics.

The following Lemma is a partial case of Theorem A from [SW].

Lemma 3.1. Let Y be a finite simplicial complex with $\pi_n(Y)$ finite. Then for every λ there is μ such that every map $f: B^n \to Y$ with $L(f|_{S^{n-1}}) \leq \lambda$ can be deformed to a μ -Lipschitz map $g: B^n \to Y$ by means of a homotopy $h_t: B^n \to Y$ with $h_t|_{S^{n-1}} = f|_{S^{n-1}}$.

Lemma 3.2. Let L be a finite dimensional complex and let K be a finite complex with finite homotopy groups $\pi_i(K)$ for $i \leq \dim L+1$. Let $f, g: L \to K$ be homotopic Lipschitz maps. Then every homotopy between f and g can be deformed to a Lipschitz homotopy $H: L \times [0, 1] \to K$.

Proof. Let $F: L \times I \to K$ be a homotopy between f and g. By induction on n, and using Lemma 3.1 we construct a μ_n -Lipschitz map $H_n: L^{(n)} \times I \cup L \times \{0, 1\} \to K$ which is a deformation of F restricted to the n-skeleton $L^{(n)}$ such that H_n extends H_{n-1} . Here the fact that L and hence $L \times I$ are geodesic is essential for the argument because this condition will guarantee that the union of λ -Lipschitz maps on $\Delta^n \times I$ is λ' -Lipschitz on $L^{(n)} \times I$ (see [Dr2] or [Dr3] for details). Then $H = H_m$ for $m = \dim L$. \Box

Let $x_0 \in X$ be a base point in a metric space X. For $x \in X$ and $A \subset X$ we denote by $||x|| = dist(x, x_0)$ and $||A|| = max\{||z \in A\}$. By $B_r(x)$ we denote a closed r-ball in X centered in x. We use notation $B_r = B_r(x_0)$.

Let σ be an *n*-dimensional simplex spanned in a Euclidean space. By $s_{\sigma} : \sigma \to \Delta^n$ we denote a simplicial homeomorphism onto the standard *n*-simples $\Delta^n = \{(x_i) \in \mathbf{R}^{n+1} \mid \sum x_i = 1, x_i \geq 0\}$. A locally finite simplicial complex *L* with a geodesic metric on it is called an *asymptotic polyhedron* if every simplex in *L* is isometric to a simplex σ spanned in a Euclidean space and $\lim_{\|\sigma\|\to\infty} L(s_{\sigma}) = 0$.

Proposition 3.3. Let (K, d) be an asymptotic polyhedron and let d_U denote the uniform geodesic metric on K. Then the identity map $u : (K, d) \to (K, d_U)$ satisfies the condition (*).

Proof. Let R and $\epsilon > 0$ be given. There is t such that $L(s_{\sigma}) < \epsilon/(2R)$ for σ with $\|sigma\| \ge t$. Let x be with $\|x\| \ge t + 2R$. For every two points $z, z' \in B_R(x)$ a geodesic segment J = [z, z'] joining them does not intersect B_t . There is a partition of J: $z = z_0 < z_1 \cdots < z_m = z'$ such that every segment $J_i = [z_i, z_{i+1}]$ lies in a

simplex σ_i . Hence $L(\sigma_i) < \epsilon/(2R)$. Therefore $d_U(z_i, z_{i+1}) \le \epsilon/(2R)d(z_i, z_{i+1})$ and hence $d_U(z, z') \le \sum \epsilon/(2R)d(z_i, z_{i+1}) = \epsilon/(2R)d(z, z') \le \epsilon$. \Box

Gromov defined the asymptotic dimension [G1] asdim X of a metric space X as follows: asdim $X \leq n$ if for every d there are n + 1 d-disjoint uniformly bounded families \mathcal{U}_i , $i = 0, \ldots, n$ of subsets of X such that the union $\mathcal{U} = \cup \mathcal{U}_i$ is a cover of X. A family of subsets \mathcal{A} is said to be d-disjoint if for every sets A and A' we have d(x, x') > d for all $x \in A$ and $x' \in A'$.

Gromov's definition can be equivalently reformulated as follows: asdim $X \leq n$ if for any arbitrary large number d there is a uniformly bounded open cover \mathcal{U} of X with multiplicity $\leq n + 1$ and with the Lebesgue number $\geq d$ (see Assertion 1 in [BD2] for a proof).

We note that for every *n*-dimensional asymptotic polyhedron L, asdim L = n.

Let \mathcal{U} be an open cover of a metric space X. The canonical projection to the nerve $p: X \to Nerve(\mathcal{U})$ is defined by the partition of unity $\{\phi_U : X \to \mathbf{R}\}_{U \in \mathcal{U}}$, where $\phi_U(x) = d(x, X \setminus U) / \sum_{V \in \mathcal{U}} d(x, X \setminus V)$. The family $\{\phi_U : X \to \mathbf{R}\}_{U \in \mathcal{U}}$ defines a map p to the Hilbert space $l_2(\mathcal{U})$ of square summable functions on \mathcal{U} with the Dirac functions $\delta_U, U \in \mathcal{U}$ as the basis. The nerve $N(\mathcal{U})$ of the cover \mathcal{U} is realized in $l_2(\mathcal{U})$ by taking every vertex U to δ_U . Clearly, the image of p lies in the nerve. Given a family of positive numbers $\overline{\lambda} = \{\lambda_U\}_{U \in \mathcal{U}}$ we can change the above imbedding of the nerve $N(\mathcal{U})$ into $l_2(\mathcal{U})$ by taking each vertex U to $\lambda_U \delta_U$. Then the projection $p_{\mathcal{U}}^{\overline{\lambda}} : X \to N(\mathcal{U})$ to modified realization is given by the formula $p_{\mathcal{U}}^{\overline{\lambda}}(x) = \{\lambda_U \phi_U(x)\}_{U \in \mathcal{U}}$.

For a subset $A \subset X$ we denote by $L(\mathcal{U}|_A)$ the Lebesgue number of \mathcal{U} restricted to A. Thus, $L(\mathcal{U}|_A) = \inf_{y \in A} \max_{V \in \mathcal{U}} d(y, X \setminus V)$.

Proposition 3.4. Let \mathcal{U} be a locally finite cover of a geodesic metric space X with the multiplicity $\leq m$. Then the above projection to the nerve $p_{\mathcal{U}}^{\bar{\lambda}} : X \to N = N(\mathcal{U})$ for $\bar{\lambda} = \{\lambda_U\}$ with $\lambda_U = L(\mathcal{U}|_U)/(2m+1)^2$ is 1-Lipschitz where the nerve is taken with the intrinsic metric induced from $l_2(\mathcal{U})$.

Proof. We show that the map $\bar{p} = p_{\mathcal{U}}^{\lambda}$ is 1 - Lipschitz as a map to $l_2(\mathcal{U})$. The partition of N into simplices defines a locally finite partition on X such that \bar{p} is 1-Lipschitz on every piece considered as the map to N with the intrinsic metric. Since X is geodesic metric space, this would imply that \bar{p} is 1-Lipschitz. Let $x, y \in X$ and $U \in \mathcal{U}$. The triangle inequality implies

$$|d(x, X \setminus U) - d(y, X \setminus U)| \le d(x, y).$$

It is easy to see that $\sum_{V \in \mathcal{U}} d(x, X \setminus V) \geq L(\mathcal{U}|_U) = (2m+1)^2 \lambda_U$ for $x \in U$. Then

$$\begin{aligned} |\phi_U(x) - \phi_U(y)| &\leq \\ \frac{1}{\sum_{V \in \mathcal{U}} d(x, X \setminus V)} d(x, y) + d(y, X \setminus U) |\frac{1}{\sum_{V \in \mathcal{U}} d(x, X \setminus V)} - \frac{1}{\sum_{V \in \mathcal{U}} d(y, X \setminus V)}| \\ &\leq \frac{1}{(2m+1)^2 \lambda_U} d(x, y) + \frac{1}{(2m+1)^2 \lambda_U} (\sum_{V \in \mathcal{U}} |d(x, X \setminus V) - d(y, X \setminus V)|) \leq \frac{1}{\sqrt{2m+1} \lambda_U} d(x, y) \end{aligned}$$

Then ||p(x) - p(y)|| =

$$= \left(\sum_{U \in \mathcal{U}} \lambda_U^2 (\phi_U(x) - \phi_U(y))^2\right)^{\frac{1}{2}} \le \left((2m)\frac{1}{2m+1}d(x,y)^2\right)^{\frac{1}{2}} \le d(x,y)$$

Lemma 3.5. Suppose that X is a geodesic metric space with $\operatorname{asdim} X \leq n$ and let $f: X \to \mathbf{R}_+$ be a given proper function. Then there are a compact set $C \subset X$ and a map $\phi: X \to N$ to a 2n + 1-dimensional asymptotic polyhedron with $\operatorname{diam}(\phi^{-1}(\Delta)) \leq f(z)$ for all $z \in \phi^{-1}(\Delta) \setminus C$.

Proof. We construct ϕ as the projection $p_{\mathcal{U}}^{\bar{\lambda}}$ from Proposition 3.4 to the nerve of a cover \mathcal{U} of X.

Fix a monotone tending to infinity sequence of l_i . Since $\operatorname{asdim} X \leq n$, for every i there is a uniformly bounded cover \mathcal{U}_i of multiplicity n + 1 with the Lebesgue number $L(\mathcal{U}^i) > l_i$. Let m_i be an upper bound for the diameter of elements of \mathcal{U}_i . We define a sequence of r_i such that $f(X \setminus B_{r_i-m_i}) \geq 2m_{i+1}$. We take $\mathcal{U} = \bigcup_i \mathcal{U}'_i$ where

$$\mathcal{U}_i' = \{ U \setminus B_{r_{i-1}} \mid U \in \mathcal{U}_i, U \cap B_{r_i} \neq \emptyset \}$$

We take $C = B_{r_1}$ and check the condition of Lemma. By the definition the preimage $\phi^{-1}(\Delta)$ lies in the union $\cup U'_J$ of sets from \mathcal{U} with nonempty intersection. Let $x \in \cap U'_J$. Each U'_j belongs to the cover \mathcal{U}_i for some *i*. Let *k* be maximal among those *is*. Then $x \notin B_{r_{k-1}}$. Therefore $U'_j \subset X \setminus B_{r_{k-1}-m_{k-1}}$ for all *j*. Hence $f(z) \geq 2m_k \geq diam \cup_j U'_j$ for $z \in X \setminus B_{r_{k-1}-m_{k-1}}$ and k > 1.

Since $l_i \to \infty$ the nerve N realized in $l_2(\mathcal{U})$ as above with the intrinsic metric is an asymptotic polyhedron. Since the covers \mathcal{U}'_i and \mathcal{U}'_j do not intersect for |i - j| > 1, the multiplicity of \mathcal{U} is $\leq 2n + 2$. \Box

REMARK. By a standard dimension theoretic trick the polyhedron N can be chosen to be n-dimensional.

A metric space X is called *uniformly contractible* if there is a function $S : \mathbf{R}_+ \to \mathbf{R}_+$ such that every ball $B_r(x)$ is contractible to a point in the ball $B_{S(r)}(x)$.

Let $x_0 \in X$ be a base point. We denote $||x|| = d_X(x, x_0)$.

THE INTEGRAL NOVIKOV CONJECTURE FOR ASYMPTOTICALLY FINITE DIMENSIONAL GROUP\$3

Lemma 3.6. Let X be a uniformly contractible proper metric space with asdim X = n. Then given a proper function $g: X \to \mathbf{R}_+$ there are an n-dimensional asymptotic polyhedron N, a proper 1-Lipschitz map $\phi: X \to N$, and a proper homotopy inverse map $\gamma: N \to X$ with $d(x, \gamma \phi(x)) < g(x)$ for all $x \in X$. Moreover, there is a compact set $C \subset X$ such that $diam(f^{-1}(\Delta) \leq g(x)$ for all $x \in \Delta \setminus C$.

Proof. We define by induction on i a lift γ on the i-skeleton $N^{(i)}$ of the nerve of a cover of X given by Lemma 3.5 for an appropriate choice of f. We take $\gamma(v) \in \phi^{-1}(v)$ for every vertex v. Then using the uniform contractibility of X we can extend γ with control over the 1-skeleton $N^{(1)}$ and so on. Without loss of generality we may assume that X is a polyhedron of the dimension n supplied with a triangulation of mesh ≤ 1 . By induction on i we define a homotopy $H : X^{(i)} \times I \to X$ joining the identity map with $\gamma \circ \phi$. We consider a function $\psi(x) = ||x|| - \max\{d(x,y) \mid y \in H(x \times I)\}$. If ψ tends to infinity, then the map H is proper. Therefore it suffices to show that ψ tends to infinity for an appropriate choice of f. Let S be a contractibility function. We define $\rho(t) = S^{-1}(t/2)$ where S^{-1} is the inverse function for S. Then we take $f = \rho^{2n+1} \circ g$, the composition of g and 2n + 1 times iteration of ρ . Clearly $g \leq f$. We assume here that $g(x) \leq ||x||/2$. Then it is easy to verify that $\psi(x) \geq ||x||/2$. \Box

A metric space X is called *proper* if every ball $B_r(x)$ in X is compact. We recall that the Higson compactification \overline{X} of a proper metric space X can be defined as the maximal ideal space of the completion of the ring of bounded functions with the gradient tending to zero at infinity [Ro1]. The defining property of the Higson corona is the following:

(**) A continuous map $f : X \to Z$ to a compact metric space is extendable to the Higson corona νX if and only if it satisfies the condition: For arbitrary large R

$$\lim_{\|x\|\to\infty} diam(f(B_R(x))) = 0.$$

Note that a proper Lipschitz map $f: X \to Y$ induces a continuous mapping between the Higson coronas $\overline{f}: \nu X \to \nu Y$.

Theorem 3.7. Let X be a uniformly contractible geodesic proper metric space with a finite asymptotic dimension and let \bar{X} be the Higson compactification. Then $\check{H}^n(\bar{X}; \mathbf{Z}_p) = 0$ for all n and all p.

Proof. We show that every map $\alpha : \overline{X} \to K(\mathbf{Z}_p, n)$ is null homotopic. Since \overline{X} is compact, the image $\alpha(\overline{X})$ is contained in the k-skeleton $K = K(\mathbf{Z}_p, n)^{(k)}$, which is a finite complex. We fix a geodesic metric on K. Let ϵ_K be an injectivity radius in K, i.e. every two points within a distance ϵ_K can be joined by the unique geodesic. Since the map $\alpha|_X \to K$ is extendable over the Higson corona the function $R_{\alpha}(t) = L(\alpha|_{X \setminus B_t(x_0)})$ tends to zero

at infinity. We apply Lemma 3.6 with $g(x) \leq \min\{\epsilon_K/R_\alpha(||x||/2), ||x||/4\}$ to obtain an asymptotic polyhedron N and maps $\phi: X \to N$ and $\gamma: N \to X$. We may assume that ϕ is surjective. Let [u, v] be an edge in N, then $d_K(\alpha\gamma(u), \alpha\gamma(v)) \leq R_\alpha(t_0)d_X(\gamma(u), \gamma(v))$ where $t_0 = \min\{||\gamma(u)||, ||\gamma(v)||\}$. We may assume that there are $x, y \in X$ such that $\phi(x) = u$ and $\phi(y) = v$. Then

$$d_X(\gamma(u), \gamma(v)) \le d_X(x, y) + \epsilon_K / R_\alpha(\frac{1}{2} ||x||) + \epsilon_K / R_\alpha(\frac{1}{2} ||y||) \le diam\phi^{-1}[u, v] + 2\epsilon_K / R_\alpha(\frac{1}{2} ||x||)$$

provided $||x|| \leq ||y||$. Because of the inequality $g(x) \leq ||x||/4$ we have that $R_{\alpha}(t_0) \leq R_{\alpha}(||x||/2)$. We may assume that f < g and then $diam(\phi^{-1}([u, v])) < g(x)$. Summarizing all this, we obtain the inequality $d_K(\alpha\gamma(u), \alpha\gamma(v)) \leq 3\epsilon_K$. This means that the map $\alpha \circ \gamma$ is $3\epsilon_K$ -Lipschitz for N taken with the uniform metric.

Since X is contractible, the map $\alpha \circ \gamma$ is null homotopic. Note that the homotopy groups $\pi_i(K)$ are finite for $i \leq \dim N + 1$. We apply Lemma 3.2 to obtain a λ -Lipschitz homotopy $H : N_U \times I \to K$ of $\alpha \circ \gamma$ to a constant map. This homotopy defines a Lipschitz map $\tilde{H} : N_U \to K^I_{\lambda}$ to the space of λ -Lipschitz mappings of the unit interval Ito K. We note that the space K^I_{λ} is compact. Then by Proposition 3.3 $\tilde{H} \circ u : N \to K^I_{\lambda}$ satisfies the Higson extendibility condition (**). Let $\bar{h} : \bar{N} \to K^I_{\lambda}$ be the extension over the Higson corona. This extension defines a map $\bar{H} : \bar{N} \times I \to K$. The map \bar{H} is a homotopy between the extension $\xi = \overline{\alpha \circ \gamma}$ and a constant map. To complete the proof, we show that α is homotopic to $\xi \circ \bar{\phi}$ where $\bar{\phi}$ is the extension of the Lipschitz map ϕ to the Higson compactifications. Note that

$$d_K(\alpha(x), \alpha\gamma\phi(x)) \le R_\alpha(t_0)d(x, \gamma\phi(x)) \le R_\alpha(t_0)\epsilon_K/R_\alpha(\|x\|/2) \le \epsilon_K$$

where $t_0 = \min\{\|x\|, \|\gamma\phi(x)\|\} \ge \|x\|/2$. Then for every $x \in X$ we join the points $\alpha(x)$ and $\alpha\gamma\phi(x)$ by the unique geodesic $\psi_x: I \to K$. This defines a map $\tilde{\psi}: X \to K^I_{\mu}$. Since both α and $\alpha \circ \gamma \circ \phi$ satisfy the condition (*), the map $\tilde{\psi}$ has the property (*). Let $\bar{\psi}: \bar{X} \to K^I_{\mu}$ be the extension of $\tilde{\psi}$ to the Higson corona. The map $\bar{\psi}$ defines a homotopy $\Psi: \bar{X} \times I \to K$ between α an $\xi \circ \bar{\phi}$. \Box

Corollary 3.8. Suppose that a group Γ has a finite asymptotic dimension and $B\Gamma$ is a finite complex. Then the integral assembly map $A : H_*(B\Gamma; \mathbf{L}) \to L_*(\mathbf{Z}\Gamma)$ is a monomorphism.

Proof. In view of the inequality dim $\nu \Gamma \leq \operatorname{asdim} \Gamma$ [DKU] we can apply Theorem 2.4. \Box

THE INTEGRAL NOVIKOV CONJECTURE FOR ASYMPTOTICALLY FINITE DIMENSIONAL GROUP\$5

References

- [BD1] G. Bell and A. Dranishnikov, On asymptotic dimension of groups, Algebr. Geom. Topol. 1 (2001), 57-71.
- [BD2] G. Bell and A. Dranishnikov, On asymptotic dimension of groups acting on trees, Geom. Dedicata 103 (2004), 89-101.
- [C] G. Carlsson, Novikov conjectures, index theorems and rigidity, Vol. 2 (Oberwolfach, 1993), 5-127, London Math. Soc. Lecture Note Ser., 226, Cambridge Univ. Press, Cambridge, 1995.
- [CG1] G. Carlsson and B. Goldfarb, On homological coherence of discrete groups, Journal of Algebra 276 No 2 (2004), 77-97.
- [CG2] G. Carlsson and B. Goldfarb, The integral K-theoretic Novikov conjecture for groups with finite asymptotic dimension, Inventiones Math. (2004).
- [CP1] G. Carlsson and E.K. Pedersen, Controlled algebra and the Novikov conjectures for K- and L-theory, Topology 34 (1993), 731-758.
- [CP2] G. Carlsson and E.K. Pedersen, Čech homology and the Novikov conjectures for K- and L-theory, Math Scand (1997).
- [Dr1] A. Dranishnikov, Asymptotic topology, Russian Math. Surveys 55:6 (2000), 71-116.
- [Dr2] A. Dranishnikov, On hypersphericity of manifolds with finite asymptotic dimension, Trans. Amer. Math. Soc. **355 no 1** (2003), 155-167.
- [Dr3] A. Dranishnikov, On Hypereuclidean manifolds, Preprint (2000).
- [DF] A. Dranishnikov and S. Ferry, On the Higson-Roe corona, Russian Math. Surveys **52 no 5** (1997), 1017-1028.
- [DJ] A. Dranishnikov and T. Januszkiewicz, Every Coxeter group acts amenably on a compact space, Topology Proceedings **24** (1999), 135-141.
- [DKU] A. Dranishnikov, J. Keesling, V. V. Uspenskij, On the Higson corona of uniformly contractible spaces, Topology 37 (1999), no. 4, 791-803.
- [EH] D. A. Edwards and H. M. Hastings, Čech and Steenrod Homology Theories with Applications to Geometric Topology, Lecture Notes in Mathematics Proceedings 542 (1976), Springer.
- [FRR] S. Ferry, A. Ranicki and J. Rosenberg; Editors, Novikov conjectures, index theorems and rigidity, Vol. 1, 2. London Math. Soc. Lecture Note Ser., 226,, Cambridge Univ. Press, Cambridge, 1995.
- [FW] S. Ferry and S. Weinberger, A coarse approach to the Novikov conjecture, Novikov conjectures, index theorems and rigidity, Vol. 1 (Oberwolfach, 1993), 147–163, London Math. Soc. Lecture Note Ser., 226, (1995), Cambridge Univ. Press, Cambridge.
- [G1] M. Gromov, Asymptotic invariants of infinite groups, Geometric Group Theory, vol 2, Cambridge University Press, 1993.
- [G2] M. Gromov, Spaces and questions, GAFA Special Volume, Part I (2000), 118-161.
- [HR] N. Higson and J. Roe, Analytic K-homology, Oxford University Press, Oxford, 2000.
- [Ji] L. Ji, Asymptotic dimension of arithmetic groups, preprint (2003).
- [KS] J. Kaminker and C. Schochet, *K*-theory and Steenrod homology: applications to the Brown-Douglas-Fillmore theory of operator algebras, Trans. Amer. Math. Soc. **227** (1977), 63-107.
- [Kee] J. Keesling, The one-dimensional Čech cohomology of the Higson compactification and its corona, Topology Proc. **19** (1994), 129–148.
- [MM] Madsen I. and Milgram R.J., *The universal smooth surgery class*, Comm. Math. Helv. **50** (1975), 281-310.

- [P] E.K. Pedersen, Bounded and continuous control, Novikov conjectures, index theorems and rigidity, Vol. 2 (Oberwolfach, 1993), 277-284, London Math. Soc. Lecture Note Ser., 226, (1995), Cambridge Univ. Press, Cambridge.
- [Ran1] A.A. Ranicki, Algebraic L-theory and topological manifolds, Cambridge Univ. Press, Cambridge, 1992.
- [Ran2] A.A. Ranicki, Additive L-theory, K-theory 3 (1989), 163-195.
- [Ran3] A.A. Ranicki, On the Novikov conjecture, in the book "Novikov conjectures, index theorems and rigidity", Vol. 1 (Oberwolfach, 1993), 272-337, London Math. Soc. Lecture Note Ser., 226, (1995), Cambridge Univ. Press, Cambridge.
- [Ro1] J. Roe, Coarse cohomology and index theory for complete Riemannian manifolds, Memoirs Amer. Math. Soc. No. 497, 1993.
- [Ro2] J. Roe, Lectures on coarse geometry, University Lecture series, Volume 31 (2003), AMS.
- [Ros] J. Rosenberg, Novikov conjectures, index theorems and rigidity, Vol. 1 (Oberwolfach, 1993),
- 338-372, London Math. Soc. Lecture Note Ser., 226,, Cambridge Univ. Press, Cambridge, 1995.
 [SW] J. Siegel and F. Williams, Uniform bounds for isoperimetric problems, Proc. Amer. Math. Soc. 107 (1989), no 2, 459-464.
- [T] J. L. Taylor, A counter-Example in Shape Theory, Bull. Amer. Math. Soc. 81 (1975), 629-632.
- [Yu1] G. Yu, The Novikov conjecture for groups with finite asymptotic dimension, Ann. of Math 147 (1998), no. 2, 325-355.

UNIVERSITY OF FLORIDA, DEPARTMENT OF MATHEMATICS, P.O. BOX 118105, 358 LITTLE HALL, GAINESVILLE, FL 32611-8105, USA

Department of Mathematics, Rutgers University, 110 Frelinghuysen Road, Piscataway, NJ08854-8019

DEPARTMENT OF MATHEMATICS, THE UNIVERSITY OF CHICAGO, CHICAGO, IL 60637 *E-mail address:* dranish@math.ufl.edu