

Configuration-Spaces and Iterated Loop-Spaces.

Segal, Graeme

pp. 213 - 222



Terms and Conditions

The Göttingen State and University Library provides access to digitized documents strictly for noncommercial educational, research and private purposes and makes no warranty with regard to their use for other purposes.

Some of our collections are protected by copyright. Publication and/or broadcast in any form (including electronic) requires prior written permission from the Goettingen State- and University Library.

Each copy of any part of this document must contain there Terms and Conditions. With the usage of the library's online system to access or download a digitized document you accept there Terms and Conditions.

Reproductions of material on the web site may not be made for or donated to other repositories, nor may be further reproduced without written permission from the Goettingen State- and University Library

For reproduction requests and permissions, please contact us. If citing materials, please give proper attribution of the source.

Contact:

Niedersächsische Staats- und Universitätsbibliothek

Digitalisierungszentrum

37070 Goettingen

Germany

Email: gdz@www.sub.uni-goettingen.de

Purchase a CD-ROM

The Goettingen State and University Library offers CD-ROMs containing whole volumes / monographs in PDF for Adobe Acrobat. The PDF-version contains the table of contents as bookmarks, which allows easy navigation in the document. For availability and pricing, please contact:

Niedersaechische Staats- und Universitaetsbibliothek Goettingen - Digitalisierungszentrum

37070 Goettingen, Germany, Email: gdz@www.sub.uni-goettingen.de

Configuration-Spaces and Iterated Loop-Spaces

Graeme Segal (Oxford)

§ 1. Introduction

The object of this paper is to prove a theorem relating “configuration-spaces” to iterated loop-spaces. The idea of the connection between them seems to be due to Boardman and Vogt [2]. Part of the theorem has been proved by May [6]; the general case has been announced by Giffen [4], whose method is to deduce it from the work of Milgram [7].

Let C_n be the space of finite subsets of \mathbb{R}^n . It is topologized as the disjoint union $\coprod_{k \geq 0} C_{n,k}$, where $C_{n,k}$ is the space of subsets of cardinal k , regarded as the orbit-space of the action of the symmetric group Σ_k on the space $\tilde{C}_{n,k}$ of ordered subsets of cardinal k , which is an open subset of \mathbb{R}^{nk} .

There is a map from C_n to $\Omega^n S^n$, the space of base-point preserving maps $S^n \rightarrow S^n$, where S^n is the n -sphere. One description of it (at least when $n > 1$) is as follows. Think of a finite subset c of \mathbb{R}^n as a set of electrically charged particles, each of charge $+1$, and associate to it the electric field E_c it generates. This is a map $E_c: \mathbb{R}^n - c \rightarrow \mathbb{R}^n$ which can be extended to a continuous map $E_c: \mathbb{R}^n \cup \infty \rightarrow \mathbb{R}^n \cup \infty$ by defining $E_c(\xi) = \infty$ if $\xi \in c$, and $E_c(\infty) = 0$. Then E_c can be regarded as a base-point-preserving map $S^n \rightarrow S^n$, where the base-point is ∞ on the left and 0 on the right. Notice that the map $c \mapsto E_c$ takes $C_{n,k}$ into $\Omega^n S^n_{(k)}$, the space of maps of degree k .

Our object is to prove that C_n is an approximation to $\Omega^n S^n$, in the sense that the two spaces have composition-laws which are respected by the map $C_n \rightarrow \Omega^n S^n$, and the induced map of classifying-spaces is a homotopy-equivalence. In view of the “group-completion” theorem of Barratt-Priddy-Quillen [1, 8] one can say equivalently that $C_{n,k} \rightarrow \Omega^n S^n_{(k)}$ induces an isomorphism of integral homology up to a dimension tending to ∞ with k . But to make precise statements it is convenient to introduce a modification of the space C_n .

If $u \leq v$ in \mathbb{R} , let $\mathbb{R}^n_{u,v}$ denote the open set $]u, v[\times \mathbb{R}^{n-1}$ in \mathbb{R}^n . Then C_n is homotopy-equivalent to the space

$$C'_n = \{(c, t) \in C_n \times \mathbb{R} : t \geq 0, c \subset \mathbb{R}^n_{0,t}\},$$

which has an associative composition-law given by juxtaposition, i.e. $(c, t) \cdot (c', t') = (c \cup T_t c', t + t')$, where $T_t: \mathbb{R}_{0,t}^n \rightarrow \mathbb{R}_{t,t+t'}^n$ is translation. As a topological monoid C'_n has a classifying-space BC'_n .

Theorem 1. $BC'_n \simeq \Omega^{n-1} S^n$, the $(n-1)$ -fold loop-space of S^n .

More generally, let X be a space with a good base-point denoted by 0. (That means that there exists a homotopy $h_t: X \rightarrow X$ ($0 \leq t \leq 1$) such that $h_0 = \text{identity}$, $h_1(0) = 0$, and $h_1^{-1}(0)$ is a neighbourhood of 0.) Let $C_n(X)$ be the space of finite subsets of \mathbb{R}^n "labelled by X " in the following sense. A point of $C_n(X)$ is a pair (c, x) , where c is a finite subset of \mathbb{R}^n , and $x: c \rightarrow X$ is a map. But (c, x) is identified with (c', x') if $c \subset c'$, $x'|_c = x$, and $x'(\xi) = 0$ when $\xi \notin c$.

$C_n(X)$ is topologized as a quotient of the disjoint union

$$\coprod_{k \geq 0} (\tilde{C}_{n,k} \times X^k) / \Sigma_k.$$

As before, $C_n(X)$ is homotopy-equivalent to a topological monoid $C'_n(X) \subset C_n(X) \times \mathbb{R}_+$, consisting of triples (c, x, t) such that $t \geq 0$ and $c \subset \mathbb{R}_{0,t}^n$.

Theorem 2. $BC'_n(X) \simeq \Omega^{n-1} S^n X$.

Here $S^n X$ is the n -fold reduced suspension of X . Theorem 1 is Theorem 2 for the case $X = S^0$. If X is path-connected so is $C'_n(X)$, and then $\Omega BC'_n(X) \simeq C'_n(X) \simeq C_n(X)$, and one has

Theorem 3. If X is path-connected $C_n(X) \simeq \Omega^n S^n X$.

This has been proved by May [6].

Some other special cases are

(a) If $n=1$, $C'_n(X)$ is equivalent to the free monoid MX on X , in the sense that there is a homomorphism $C'_1(X) \rightarrow MX$ which is a homotopy-equivalence. Thus one obtains the theorem of James [5] that $BMX \simeq SX$.

(b) If $n=2$, $C_{2,k}$ is the classifying-space for the braid group B_k on k strings. Thus one has $B(\coprod_{k \geq 0} B(B_k)) \simeq \Omega S^2$.

(c) Because $\tilde{C}_{n,k}$ is $(n-2)$ -connected, being the complement of some linear subspaces of codimension n in \mathbb{R}^{n^k} , one has $C_{n,k} \rightarrow B\Sigma_k$ as $n \rightarrow \infty$. This gives the theorem of Barratt-Priddy-Quillen that $B(\coprod_{k \geq 0} B\Sigma_k) \simeq \Omega^{\infty-1} S^\infty$.

The theorems above do not mention a specific map between the configuration-spaces and the loop-spaces. I shall return to this question in § 3.

§ 2. Proofs

Theorem 2 is obtained by induction from

Proposition (2.1). $BC'_n(X) \cong C_{n-1}(SX)$.

For $C_{n-1}(SX)$ is connected, so $C_{n-1}(SX) \cong C'_{n-1}(SX) \cong \Omega BC'_{n-1}(SX) \cong \Omega C_{n-2}(S^2 X) \cong \dots \cong \Omega^{n-1} C_0(S^n X) = \Omega^{n-1} S^n X$.

The proof of (2.1) is based on the idea of a “partial monoid”.

Definition (2.2). A *partial monoid* is a space M with a subspace $M_2 \subset M \times M$ and a map $M_2 \rightarrow M$, written $(m, m') \mapsto m \cdot m'$, such that

(a) there is an element 1 in M such that $m \cdot 1$ and $1 \cdot m$ are defined for all m in M , and $1 \cdot m = m \cdot 1 = m$.

(b) $m \cdot (m' m'') = (m \cdot m') \cdot m''$ for all m, m', m'' in M , in the sense that if one side is defined then the other is too, and they are equal.

A partial monoid M has a classifying-space BM , defined as follows. Let $M_k \subset M \times \dots \times M$ be the space of composable k -tuples. The M_k form a (semi)simplicial space, in which $d_i: M_k \rightarrow M_{k-1}$ and $s_i: M_k \rightarrow M_{k+1}$ are defined by

$$\begin{aligned} d_i(m_1, \dots, m_k) &= (m_2, \dots, m_k) & \text{if } i=0 \\ &= (m_1, \dots, m_i \cdot m_{i+1}, \dots, m_k) & \text{if } 0 < i < k \\ &= (m_1, \dots, m_{k-1}) & \text{if } i=k \\ s_i(m_1, \dots, m_k) &= (m_1, \dots, m_i, 1, m_{i+1}, \dots, m_k) & \text{if } 0 \leq i \leq k. \end{aligned}$$

BM is defined as the realization of this simplicial space [9]. If M is actually a monoid (i.e. if $M_2 = M \times M$) then BM is the usual classifying-space. On the other hand if M has trivial composition (i.e. $M_2 = M \vee M$) then $BM = SM$, the reduced suspension of M .

The space $C_{n-1}(X)$ can be regarded as a partial monoid, in which (c, x) and (c', x') are composable if and only if c and c' are disjoint, and then $(c, x) \cdot (c', x') = (c \cup c', x \cup x')$.

Proposition (2.3). $BC_{n-1}(X) \cong C_{n-1}(SX)$.

Proof. Write $M = C_{n-1}(X)$. By definition BM is a quotient of the disjoint union of the spaces $M_k \times \Delta^k$ for $k \geq 0$. Regard the standard simplex Δ^k as $\{(t_1, \dots, t_k) \in \mathbb{R}^k: 0 \leq t_1 \leq \dots \leq t_k \leq 1\}$. Define $M_k \times \Delta^k \rightarrow C_{n-1}(SX)$ by $((c_1, x_1), \dots, (c_k, x_k); t_1, \dots, t_k) \mapsto (c, \tilde{x})$, where $c = \bigcup c_i$ and $\tilde{x}: c \rightarrow SX$ takes $P \in c_i$ to $(t_i, x_i(P)) \in SX$.

These maps induce a map $BM \rightarrow C_{n-1}(SX)$ which is obviously surjective. It is injective because each point of BM is representable in the above form with $0 < t_1 < \dots < t_k < 1$, all c_i non-empty, and $x_i(c_i) \subset X - \{0\}$. It is a homeomorphism because one can define a continuous inverse-

map, observing that $C_{n-1}(SX)$ is a quotient of the disjoint union for $k \geq 0$ of the spaces $\tilde{C}_{n-1,k} \times X^k \times [0, 1]^k$, which map to $M_k \times [0, 1]^k$.

The partial monoid $C_{n-1}(X)$ is related to the monoid $C'_n(X)$ by means of a sub-partial-monoid of the latter. Call an element (c, x, t) of $C'_n(X)$ *projectable* if c is mapped injectively by the projection

$$pr_2: \mathbb{R}_{0,t}^n \rightarrow \mathbb{R}^{n-1}.$$

The projectable elements form a subspace which the composition-law of $C'_n(X)$ makes into a partial monoid. Projection defines a homomorphism $C''_n(X) \rightarrow C_{n-1}(X)$, and elements of $C''_n(X)$ are composable if and only if their images in $C_{n-1}(X)$ are. Furthermore $C''_n(X) \rightarrow C_{n-1}(X)$ is a homotopy-equivalence (with inverse $(c, x) \mapsto (s(c), x, 1)$, where s is any cross-section of $pr_2: \mathbb{R}_{0,t}^n \rightarrow \mathbb{R}^{n-1}$); and so is the map $C''_n(X)_k \rightarrow C_{n-1}(X)_k$ of spaces of composable k -tuples. Because the simplicial spaces in question are good (see Appendix 2), this implies that $BC''_n(X) \cong BC_{n-1}(X)$. So to prove (2.1) it is enough to prove

Proposition (2.4). *The inclusion $C''_n(X) \rightarrow C'_n(X)$ induces a homotopy-equivalence of classifying-spaces.*

To prove (2.4) I shall use an alternative description of the classifying-space of a partial monoid M . Let $\mathcal{C}(M)$ be the topological category [9] whose objects are the elements of M , and whose morphisms from m to m' are pairs of elements $m_1, m_2 \in M$ such that $m_1 \cdot m \cdot m_2 = m'$. Thus $\text{ob } \mathcal{C}(M) = M$, and $\text{mor } \mathcal{C}(M) = M_3$. Let $|\mathcal{C}(M)|$ be the realization or "classifying space" of $\mathcal{C}(M)$ in the sense of [9].

Proposition (2.5). $|\mathcal{C}(M)| \cong BM$.

This is a particular case of a subdivision theorem for arbitrary simplicial spaces, proved in Appendix 1. For the proof of (2.4) I shall use a modification of the category $\mathcal{C}(C'_n(X))$. Let Q be the ordered space whose points are 4-tuples $(u, v; c, x)$, with $u, v \in \mathbb{R}$, $u \leq 0 \leq v$, c a finite subset of $\mathbb{R}_{u,v}^n$, and $x: c \rightarrow X$, subject to the usual equivalence-relation. It is ordered by defining $(u, v; c, x) \leq (u', v'; c', x')$ if $[u, v] \subset [u', v']$, $c = c' \cap ([u, v] \times \mathbb{R}^{n-1})$, and $x'|_c = x$. Thinking of the topological ordered set Q as a topological category, define a functor $\pi: Q \rightarrow \mathcal{C}(C'_n(X))$ by $(u, v; c, x) \mapsto (T_{-u}c, x, v - u)$.

Lemma (2.6). $|\pi|: |Q| \rightarrow |\mathcal{C}(C'_n(X))|$ is shrinkable [3] (i.e. it has a section s such that $s|\pi| \simeq \text{identity}$ by a homotopy h_t for which $|\pi|h_t = |\pi|$).

Proof. Using the homomorphism of monoids $C'_n(X) \rightarrow \mathbb{R}_+$ which takes (c, x, t) to t one can regard Q as the fibre-product (of categories) $\mathcal{C}(C'_n(X)) \times_{\mathcal{C}(\mathbb{R}_+)} \mathcal{J}$, where \mathcal{J} is the space of intervals $[u, v]$ with $u \leq 0 \leq v$ ordered by inclusion. Forming the nerve commutes with fibre-products. But $|\mathcal{J}| \rightarrow |\mathcal{C}(\mathbb{R}_+)|$ is easily seen to be shrinkable; so $|\pi|$ is shrinkable.

Continuing the proof of (2.4), if P is the sub-ordered-space of Q consisting of all $(u, v; c, x)$ with c projectable then $\pi(P) = \mathcal{C}(C''_n(X))$; so it will be enough to show that $|P| \rightarrow |Q|$ is a homotopy-equivalence. Heuristically this is so because P is co-initial in Q , i.e. for each $q \in Q$ there is a $p \in P$ such that $p \leq q$, and if $p_1 \leq q$ and $p_2 \leq q$ there is a $p_3 \in P$ such that $p_3 \leq p_1$ and $p_3 \leq p_2$. But some further conditions are needed, and I shall use the following ad hoc lemma.

Proposition (2.7). *Let Q be a good ordered space such that*

(a) $q_1 \cap q_2 = \inf(q_1, q_2)$ *is defined whenever there exists $q \in Q$ such that $q_1 \leq q$ and $q_2 \leq q$, and*

(b) $q_1 \cap q_2$ *depends continuously on (q_1, q_2) where defined.*

Let Q_0 be an open subspace of Q such that if $q \in Q$ and $p \in Q_0$ and $q \leq p$ then $q \in Q_0$. Suppose there is a numerable covering [3] $U = \{U_\alpha\}$ of Q , and maps $f_\alpha: U_\alpha \rightarrow Q_0$ such that $f_\alpha(q) \leq q$ for all $q \in U_\alpha$.

Then $|Q_0| \rightarrow |Q|$ is a homotopy-equivalence.

In the application of (2.7) Q is as above. Using the fact that X has a good base-point, choose a homotopy $h_t: X \rightarrow X$ such that h_0 is the identity and $h_1^{-1}(0)$ is a neighbourhood of 0. This induces $h_t: Q \rightarrow Q$. Let $Q_0 = h_1^{-1}(P)$, a neighbourhood of P . One might call Q_0 the "almost projectable" elements of Q . Obviously $|P| \rightarrow |Q_0|$ is a homotopy-equivalence, with inverse induced by h_1 . But Q_0 and Q satisfy the hypotheses of (2.7) — it is proved in Appendix 2 that Q is good. Thus (2.4) is proved modulo (2.7).

The proof of (2.7) would be almost trivial if one could choose the maps f_α compatibly so as to get a continuous map $Q \rightarrow Q_0$. In general it depends on the fact that one does not change the homotopy-type of a topological category by breaking apart the space of objects into the sets of a numerable covering and introducing isomorphism between the reduplicated objects. To be precise, if C is a topological category, and $U = \{U_\alpha\}_{\alpha \in A}$ is a numerable covering of $\text{ob}(C)$, then the *disintegration* of C by U is the topological category \tilde{C} whose objects are pairs (x, α) with $\alpha \in A$ and $x \in U_\alpha$, and whose morphisms $(x, \alpha) \rightarrow (x', \alpha')$ are the morphisms from x to x' in C . Thus $\text{ob}(\tilde{C}) = \coprod_{\alpha \in A} U_\alpha$ and $\text{mor}(\tilde{C}) = \coprod_{\alpha, \beta \in A} V_{\alpha\beta}$, where $V_{\alpha\beta}$ consists of the elements of $\text{mor}(C)$ with source in U_α and target in U_β .

Proposition (2.8). *If C is a topological category, U is a numerable covering of $\text{ob}(C)$, and \tilde{C} is the disintegration of C by U , then the projection $|\tilde{C}| \rightarrow |C|$ is shrinkable.*

Proof. Let $\{C_k\}_{k \geq 0}$ be the simplicial space associated to C (i.e. $C_0 = \text{ob}(C)$, $C_1 = \text{mor}(C)$, etc.), and let $\{\tilde{C}_k\}_{k \geq 0}$ be that associated to \tilde{C} . Then $\tilde{C}_k = (\tilde{C}_0)^{k+1} \times_{(C_0)k+1} C_k$. Now for any space Y there is a simplicial space $\{Y^{k+1}\}_{k \geq 0}$ whose realization is a contractible space called EY [9].

Because realization commutes with fibre-products $|\tilde{C}| = E\tilde{C}_0 \times_{EC_0} |C|$, and it is enough to show that $E\tilde{C}_0 \rightarrow EC_0$ is shrinkable. But $\{EU_a\}_{a \in A}$ is a numerable covering of EC_0 over which $E\tilde{C}_0 \rightarrow EC_0$ is shrinkable, so the result follows from [3].

Proof of (2.7). Let $p: \tilde{Q} \rightarrow Q$ be the disintegration associated to U . \tilde{Q} is a preordered space. The f_a define a map $f: \tilde{Q} \rightarrow Q_0$ such that $f(\xi) \leq p(\xi)$ for all $\xi \in \tilde{Q}$. Let $\text{chn}(\tilde{Q})$ be the space of finite chains of \tilde{Q} ordered by inclusion (cf. Appendix 2). Define $F: \text{chn}(\tilde{Q}) \rightarrow Q_0$ by $(\xi_0 \leq \dots \leq \xi_k) \mapsto \inf(f(\xi_0), \dots, f(\xi_k))$. This is order-preserving, and $F(\sigma) \leq r \text{chn}(p)(\sigma)$ for $\sigma \in \text{chn}(Q)$, where $r: \text{chn}(\tilde{Q}) \rightarrow \tilde{Q}$ is $(\xi_0 \leq \dots \leq \xi_k) \mapsto \xi_0$. As $|r|$ is a homotopy-equivalence by Appendix 2, and $|p|$, and so $|\text{chn}(p)|$, is one by (2.8), and $|F| \simeq |r| |\text{chn}(p)|$ by [9] (), it follows that the composite $|\text{chn}(\tilde{Q})| \rightarrow |Q_0| \rightarrow |Q|$ is a homotopy-equivalence. Similarly the composite $|\text{chn}(\tilde{Q}_0)| \rightarrow |\text{chn}(\tilde{Q})| \rightarrow |Q_0|$ is a homotopy-equivalence; and so $|Q_0| \rightarrow |Q|$ is one, as desired.

§ 3. The Map $C_n(X) \rightarrow \Omega^n S^n X$

Despite its picturesqueness the electrostatic map described in §1 is not very convenient in practice. It is homotopic, however, to the following map. Let $D_n(X)$ be the space of finite sets of disjoint open unit disks in \mathbb{R}^n , labelled by X . (This is a closed subset of $C_n(X)$, and obviously a deformation retract of it.) Choose a fixed map f of degree 1 from a standard disk D to S^n , taking the boundary to the base-point. Then associate to a point $(\{i_a: D \rightarrow \mathbb{R}^n\}_{a \in c}, x: c \rightarrow X)$ of $D_n(X)$ the map $\phi: \mathbb{R}^n \cup \infty \rightarrow S^n X$ defined by

$$\begin{aligned} \phi(\xi) &= (f i_a^{-1}(\xi), x(a)) \quad \text{if } \xi \in i_a(D), \text{ and} \\ &= \text{base-point otherwise.} \end{aligned}$$

Regard ϕ as a point of $\Omega^n S^n X$.

$D_n(X)$ can be regarded as a partial monoid in n different ways, the composition in each case being superimposition, but two sets of disks being composable for the i -th law if and only if they are separated by a hyperplane perpendicular to the i -th coordinate direction. These composition-laws are compatible in the sense that each is a homomorphism for the others, so one can use each of them in turn and thus define an n -fold classifying-space $B^n D_n(X)$. There is a map $X \rightarrow D_n(X)$ taking x to the unit disk at the origin labelled by x . Giving X n trivial composition-laws (so that $B^n X = S^n X$), $X \rightarrow D_n(X)$ is a homomorphism for all n laws, and induces $S^n X \rightarrow B^n D_n(X)$. Evidently what was proved in §2 was that $S^n X \rightarrow B^n D_n(X)$.

On the other hand the space of maps $\phi: \mathbb{R}^n \cup \infty \rightarrow S^n X$ also has n partial composition-laws. Define support $(\phi) = \phi^{-1}(S^n X - \{0\})$. Then ϕ_1

and ϕ_2 are composable for the i -th law if their supports are separated by a hyperplane perpendicular to the i -th axis, and the composite is in any case given by "union". The map $X \rightarrow \Omega^n S^n X$ is an n -fold homomorphism, so induces $S^n X \rightarrow B^n \Omega^n S^n X$, which one knows classically to be a homotopy-equivalence. But the map $D_n(X) \rightarrow \Omega^n S^n X$ defined above is an n -fold homomorphism, and the diagram


$$\begin{array}{ccc} & X & \\ & \swarrow \quad \searrow & \\ D_n(X) & \rightarrow & \Omega^n S^n X \end{array}$$

commutes. So $D_n(X) \rightarrow \Omega^n S^n X$ induces an isomorphism of classifying-spaces, as desired.

Appendix 1. The Edgewise Subdivision of a Simplicial Space

This is more or less due to Quillen.

The standard n -simplex $\Delta^n = \{(t_1, \dots, t_n) \in \mathbb{R}^n : 0 \leq t_1 \leq \dots \leq t_n \leq 1\}$ can be subdivided into 2^n n -simplexes corresponding to the 2^n possible orders in which the 2^n numbers $(t_1, \dots, t_n; 1-t_n, \dots, 1-t_1)$ can occur in $[0, 1]$. When $n=2$ the diagram is



In general one puts a new vertex P_{ij} at the mid-point of each edge $P_{ii}P_{jj}$ of Δ^n , the original vertices being denoted $P_{ii} (0 \leq i \leq n)$. And $P_{i_0 j_0}, \dots, P_{i_k j_k}$ span a simplex of the subdivision if $i_0 \geq \dots \geq i_k$ and $j_0 \leq \dots \leq j_k$.

This subdivision of a simplex, being functorial for simplicial maps, induces a subdivision of $|A|$ for any simplicial space A . Thereby $|A|$ is expressed as the realization of a simplicial space B such that $B_n = A_{2n+1}$. To be precise, let Δ be the category of finite ordered sets, so that A is a contravariant functor from Δ to spaces. There is a functor $T: \Delta \rightarrow \Delta$ which takes the set $\{\alpha_0, \dots, \alpha_n\}$ with $n+1$ elements to the set $\{\alpha_0, \dots, \alpha_n, \alpha'_n, \dots, \alpha'_0\}$ with $2n+2$ elements, ordered as written. Then B is $A \cdot T$.

Proposition (A.1). For any simplicial space A , $|A| \cong |A \cdot T|$.

Proof. To write down maps $|A| \hookrightarrow |A \cdot T|$, observe that the 2^n simplexes into which Δ^n is subdivided can be indexed Δ^n_θ , where θ runs through a set of 2^n maps $[2n+1] \rightarrow [n]$; and Δ^n_θ is the image of the composition $\theta_* i$, where $i: \Delta^n \rightarrow \Delta^{2n+1}$ is $(t_1, \dots, t_n) \mapsto (\frac{1}{2}t_1, \dots, \frac{1}{2}t_n, \frac{1}{2}, 1 - \frac{1}{2}t_n, \dots, 1 - \frac{1}{2}t_1)$.

Then the maps $A_{2n+1} \times \Delta^n \rightarrow A_{2n+1} \times \Delta^{2n+1}$ given by $(a, \xi) \mapsto (a, i\xi)$ induce $|A \cdot T| \rightarrow |A|$; and the maps $A_n \times \Delta^n \rightarrow A_{2n+1} \times \Delta^n$ given by $(a, \theta_* i\xi) \mapsto (\theta^* a, \xi)$ induce its inverse.

Appendix 2. Good Simplicial Spaces

Goodness is a condition on a simplicial space which ensures that its realization has convenient properties. (May [6] uses “strictly proper” for a similar idea.) I have discussed the condition at greater length in [10].

First observe that for any simplicial space $A = \{A_n\}$ there are n distinguished subsets $A_{n,i}$ ($1 \leq i \leq n$) in A_n , homeomorphic images of A_{n-1} by the degeneracy maps $s_i: A_{n-1} \rightarrow A_n$.

Definition. A simplicial space A is *good* if for each n there exists a homotopy $f_t: A_n \rightarrow A_n$ ($0 \leq t \leq 1$) such that

- (i) $f_0 = \text{identity}$
- (ii) $f_t(A_{n,i}) \subset A_{n,i}$ for $1 \leq i \leq n$
- (iii) $f_1^{-1}(A_{n,i})$ is a neighbourhood of $A_{n,i}$ in A_n for $1 \leq i \leq n$.

Two results about good simplicial spaces are used in this paper. They are proved in [10].

Proposition (A.1). *If $\phi: A \rightarrow B$ is a map of good simplicial spaces, and $\phi_n: A_n \rightarrow B_n$ is a homotopy-equivalence for each n then $|f|: |A| \rightarrow |B|$ is a homotopy-equivalence.*

To state the second result one first associates to a simplicial space A a topological category $\text{simp}(A)$. Its objects are pairs (n, a) , where $n \geq 0$ and $a \in A_n$, and its morphisms $(n, a) \rightarrow (m, b)$ are morphisms $\theta: [n] \rightarrow [m]$ in Δ such that $\theta^* b = a$. Thus

$$\text{ob}(\text{simp}(A)) = \coprod_{n \geq 0} A_n, \quad \text{and} \quad \text{mor}(\text{simp}(A)) = \coprod_{\theta: [n] \rightarrow [m]} A_m.$$

There is a natural map $|\text{simp}(A)| \rightarrow |A|$.

Proposition (A.2). *If A is a good simplicial space, $|\text{simp}(A)| \rightarrow |A|$ is a homotopy-equivalence.*

In the case occurring in this paper A is the simplicial space arising from an ordered space Q . Then $\text{simp}(A)$ is precisely $\text{chn}(Q)$, the space of finite chains $q_0 \leq \dots \leq q_n$ in Q , ordered by inclusion; and $|\text{simp}(A)| \rightarrow |A|$ is induced by the order-reversing map $(q_0 \leq \dots \leq q_n) \mapsto q_0$.

I call a monoid, partial monoid, category, ordered space, etc., *good* if it gives rise to a good simplicial space. One needs to know that the condition holds for all the examples arising in the paper. Notice first:

1. A monoid is good if and only if it is locally contractible at 1.
2. A neighbourhood of the identity in a good monoid is a good partial monoid.
3. The edgewise subdivision of a good simplicial space is good.

Now we must consider in turn $C_{n-1}(X)$, $C'_n(X)$, $C''_n(X)$, Q . Because X has a good base-point there is a homotopy $h_t: X \rightarrow X$ such that $h_0 = id$, $h_1^{-1}(0) = U$, a neighbourhood of 0. The map $h_t: C_{n-1}(X) \rightarrow C_{n-1}(X)$ contracts a neighbourhood of the identity through partial-monoid-homomorphisms, proving $C_{n-1}(X)$ is good. On the other hand $C'_n(U)$ is a contractible neighbourhood of the identity in $C'_n(X)$, so the latter is good. For the same reason $h_1^{-1} C''_n(X)$ is good. But this is homotopy-equivalent to $C''_n(X)$ as *partial monoid*, so $C''_n(X)$ is good. Finally Q is good because $Q \rightarrow \mathcal{C}(C'_n(X))$ is shrinkable.

References

1. Barratt, M.G., Priddy, S.: On the homology of non-connected monoids and their associated groups. *Comment. Math. Helvet.* **47**, 1–14 (1972).
2. Boardman, J.M., Vogt, R.M.: Homotopy-everything H -spaces. *Bull. Amer. Math. Soc.* **74**, 1117–1122 (1968).
3. Dold, A.: Partitions of unity in the theory of fibrations. *Ann. of Math.* **78**, 223–255 (1963).
4. Giffen, C.: To appear.
5. James, I. M.: Reduced product spaces. *Ann. of Math.* **62**, 170–197 (1955).
6. May, J.P.: The geometry of iterated loop spaces. *Springer Lecture Notes in Mathematics* **271**. Berlin-Heidelberg-New York 1972.
7. Milgram, R.J.: Iterated loop spaces. *Ann. of Math.* **84**, 386–403 (1966).
8. Quillen, D.G.: The group completion of a simplicial monoid. To appear.
9. Segal, G.B.: Classifying spaces and spectral sequences. *Publ. Math. I.H.E.S. Paris* **34**, 105–112 (1968).
10. Segal, G.B.: Categories and cohomology theories. *Topology*.

Graeme Segal
Mathematical Institute
University of Oxford
24–29. St. Giles
Oxford OX1 3LB, U.K.

(Received March 15, 1973)

