

# CUP- $i$ PRODUCTS AND THE ADAMS SPECTRAL SEQUENCE†‡

DANIEL S. KAHN

(Received 28 October 1968)

## §1. INTRODUCTION

THE COHOMOLOGY  $H(A)$  of the mod 2 Steenrod algebra  $A$  is isomorphic to the  $E_2$  term of the mod 2 Adams spectral sequence  $\{E_r(S^0)\}$  [2]. In this paper we extend the results of [5] in relating cup- $i$  products in  $H(A)$  to the structure of the Adams spectral sequence.

The main result of the paper is contained in

**THEOREM (1.1).** *Suppose that  $\alpha \in E_2^{s,s+t}(S^0)$  is a permanent cycle and that both*

- (i)  $(k+1) \leq \rho(t+k+1)$  and
  - (ii)  $(t+k) \equiv 2^{i+1} - 1 \pmod{2^{i+1}}$ ,  $i = 0, 1, 2$  or 3.
- Then  $Sq_k \alpha$  is a permanent cycle and  $Sq_{k+2^i} \alpha$  survives to  $E_{2^i+1}(S^0)$  where*

$$\delta_{2^i+1}(Sq_{k+2^i} \alpha) = h_i(Sq_k \alpha).$$

(1.1) occurs in this paper as Theorems (5.1) and (5.4). In the statement of (1.1),  $\rho((2a+1)2^{c+4d}) = 2^c + 8d$  where  $0 \leq c < 4$ . Also,  $Sq_k \alpha = \alpha \cup_k \alpha$ . ( $Sq_k \alpha$  should not be confused with  $Sq^k \alpha$ .)

For  $k = 0$ , the differentials given in (1.1) correspond to identities in  $\pi_*^S(S^0) = G_*$ . For example, if  $\alpha \in G_t$ , then  $2\alpha^2 = 0$  if  $t$  is even and  $\eta\alpha^2 = 0$  if  $t \equiv 3(4)$ . Identities such as these are proved using the quadratic construction which is also the main tool in the present study. The quadratic construction is a functor from pointed spaces to filtered spaces which has been studied by J. F. Adams, M. G. Barratt and M. Mahowald (unpublished). Theorem (4.4), our main technical result, was an early conjecture in the study of the quadratic construction. The author is indebted to J. F. Adams, M. G. Barratt and M. Mahowald for conversations and correspondence which were helpful in the present work.

This paper is organized as follows:

§2 recalls the construction of the cup- $i$  products in the cohomology of the Steenrod algebra. §3 outlines the quadratic construction. The main technical results are proved in §4, and the applications of Theorem (4.4) to the Adams spectral sequence are carried out in §5. The quadratic construction can also be used to define cup- $i$  products in  $G_*$ . §6 discusses the relation of some of these homotopy operations to the cup- $i$  products in  $H(A)$ .

† This research was supported in part by NSF GP-5591.

‡ A preliminary announcement of this research appeared in: D. S. KAHN, Squaring operations in the Adams spectral sequence, *Bull. Am. math. Soc.*, to appear.

§2. CUP- $i$  PRODUCTS IN  $H(A)$ 

In this section, we recall the definition of squaring operations in the cohomology of  $A$ , the mod 2 Steenrod algebra [6; Chapt. II, §5]. The construction itself applies to any connected co-commutative Hopf algebra over  $Z_2$ , but our only concern is with the Steenrod algebra.

The cohomology of  $A$  may be calculated from the bar construction  $B(A)$  [1, p. 32]. Thus,  $H^{s,t}(A)$  is a subquotient of  $\text{Hom}_A(B_s(A), Z_2)$ .  $B(A) \otimes B(A)$  is made into an  $A$ -module via the diagonal map  $\psi : A \rightarrow A \otimes A$ . Since  $B(A) \otimes B(A)$  is acyclic,  $\exists$  an  $A$ -map  $D_0 : B(A) \rightarrow B(A) \otimes B(A)$  commuting with the augmentation.  $D_0$  may be used to compute cup products in  $H(A)$ .

Since  $A$  is co-commutative, the twisting map  $\rho : B(A) \otimes B(A) \rightarrow B(A) \otimes B(A)$  given by  $x \otimes y \rightarrow y \otimes x$  is an  $A$  homomorphism commuting with augmentation. Again using the acyclicity of  $B(A) \otimes B(A)$ , there exists a chain homotopy

$$D_1 : B(A) \rightarrow B(A) \otimes B(A)$$

such that  $\partial D_1 + D_1 \partial = \rho D_0 + D_0$ . (No signs are needed since we are working mod 2.) Continuing as in the definition of the Steenrod operations in ordinary topology [8; V, §9], we obtain a sequence of  $A$  homomorphisms

$$D_j : B_n(A) \rightarrow (B(A) \otimes B(A))_{n+j}$$

such that

(2.1)  $D_0$  commutes with augmentation, and

(2.2) for  $j > 0$ ,  $\partial D_j + D_j \partial + D_{j-1} + \rho D_{j-1} = 0$ .

If  $\{D_j\}$  and  $\{D'_j\}$  are any two such sequences, there exists a sequence of  $A$  maps

$$E_j : B_n(A) \rightarrow (B(A) \otimes B(A))_{n+j}, j \geq 0$$

such that

(2.3)  $E_0 = 0$ , and

(2.4) for  $j \geq 0$ ,  $\partial E_{j+1} + E_{j+1} \partial + E_j + \rho E_j + D_j + D'_j = 0$

We now define, for  $u \in \text{Hom}_A(B_n(A), Z_2)$ , an element  $u \cup_i u \in \text{Hom}_A(B_{2n-i}(A), Z_2)$  by

$$(u \cup_i u)(\sigma) = (u \times u)(D_i \sigma),$$

where  $\sigma \in B_{2n-i}(A)$ . Using (2.1) and (2.2), we see that if  $u$  is cocycle, so also is  $u \cup_i u$  and that the cohomology class of  $u \cup_i u$  depends only on the cohomology class of  $u$ . Using (2.3) and (2.4) we see that the cohomology class of  $u \cup_i u$  does not depend on the choice of the sequence  $\{D_j\}$  satisfying (2.1) and (2.2). Thus we have defined  $\alpha \cup_i \alpha \in H^{2s-i, 2t}(A)$  for  $\alpha \in H^{s,t}(A)$ . We will also write

$$\text{Sq}_i \alpha = \text{Sq}^{2s-i} \alpha = \alpha \cup_i \alpha.$$

*Remark (2.5).* In order to define  $\alpha \cup_i \alpha$ , it suffices to define  $D_j$  satisfying (2.1) and (2.2) only for  $j \leq i$ .

## §3. THE QUADRATIC CONSTRUCTION

We describe here the quadratic construction, a functor from pointed spaces to filtered spaces which has been studied by Adams, Barratt and Mahowald (unpublished). The results of this section are well known to these authors. For convenience, we shall only discuss the construction for finite *CW* complexes with base point.

Let  $X$  be a finite *CW* complex with base point. Denote by  $Q^n(X)$  the space  $S^n \times (X \wedge X)$ . (If  $B$  is a space with base point  $b_0$ ,  $A \times B$  shall mean  $(A \times B)/(A \times b_0)$ .) Define the involution  $T: Q^n(X) \rightarrow Q^n(X)$  by  $T(x, y, z) = (-x, z, y)$ , where  $-x$  is the point of  $S^n$  antipodal to  $x$ . This defines an action of  $Z_2$  on  $Q^n(X)$  and we define  $Q^n(X) = Q^n(X)/Z_2$ .  $Q(X)$  shall mean  $Q^\infty(X)$  and is called the *quadratic construction* on  $X$ .  $Q(X)$  is naturally filtered by the subspaces  $\{Q^n(X)\}$ .

If  $f: X \rightarrow Y$  is a map of pointed complexes, then  $Q'(f): Q'(X) \rightarrow Q'(Y)$  defined by  $(x, y, z) \rightarrow (x, f(y), f(z))$  induces a map  $Q(f): Q(X) \rightarrow Q(Y)$  and  $Q(f)|Q^n(X) = Q^n(f): Q^n(X) \rightarrow Q^n(Y)$ . Thus each  $Q^n$  is a functor and if  $n < m$ ,  $Q^n$  is a subfunctor of  $Q^m$ .

*Example* (3.1).  $Q^n(S^m)$  is homeomorphic with  $S^m \wedge P_m^{m+n} = \Sigma^m P_m^{m+n}$ . (By  $P_a^b$  we mean the stunted projective space  $RP^b/RP^{a-1}$ .)

*Proof.*  $Q^n(S^m)$  may also be described as the one point compactification of  $(S^n \times R^m \times R^m)/Z_2$  where the action of  $Z_2$  on  $S^n \times R^m \times R^m$  is given by  $T'(x, y, z) = (-x, z, y)$ . Now  $S^m \wedge P_m^{m+n}$  can be described as the one point compactification of  $(S^n \times R^m \times R^m)/Z_2$  where the action of  $Z_2$  on  $S^n \times R^m \times R^m$  is given by  $T''(x, y, z) = (-x, y, -z)$  [4, p. 205]. Since, as is easily seen,  $(S^n \times R^m \times R^m, T')$  and  $(S^n \times R^m \times R^m, T'')$  are equivariantly homeomorphic, the result follows.

We leave it to the reader to verify the following three elementary properties of the quadratic construction.

**PROPOSITION (3.2).** *If  $f, g: X \rightarrow Y$  are homotopic relative base points, then there exists a homotopy  $H: Q(X) \times I \rightarrow Q(Y)$  of  $Q(f)$  to  $Q(g)$  such that  $H|Q^n(X) \times I$  gives a homotopy of  $Q^n(f)$  to  $Q^n(g)$ .*

**COROLLARY (3.3).** *If  $X$  and  $Y$  have the same homotopy type as pointed spaces, then  $Q(X)$  and  $Q(Y)$  have the same homotopy type, preserving filtration.*

**PROPOSITION (3.4).**  $Q^n(X)/Q^{n-1}(X) \approx S^n \wedge X \wedge X$  and  $Q^n(f)/Q^{n-1}(f) \approx S^n \wedge f \wedge f$ .

## §4. THE MAIN THEOREM

Our discussion of the Adams spectral sequence will follow the exposition given in [2], with the exception that we will use the smash product rather than the join in treating products. Finite *CW* approximations through the stable range respecting base points will be used. We will omit specific mention of the skeletons on which various stable constructions may be carried out.

Let  $W_0 \supset W_1 \supset \dots$  be a realization for  $B(A)$  with  $W_0$  having the homotopy type of  $S^{2n}$ . Let  $Y_0 \supset Y_1 \supset \dots$  be a realization for  $B(A)$  with  $Y_0$  having the homotopy type of  $S^n$ .

Then  $Z_0 = Y_0 \wedge Y_0$  with the product filtration is a realization for  $B(A) \otimes B(A)$ . The switching map  $\tau: Y \wedge Y \rightarrow Y \wedge Y$  is a realization of  $\rho: B(A) \otimes B(A) \rightarrow B(A) \otimes B(A)$ , the switching map for  $B(A) \otimes B(A)$ .

LEMMA (4.1): If  $n = 2^q$  where  $q \geq \phi(r)$  [3],  $S^n = P_n^n$  is a retract of  $P_n^{n+r}$ .

Proof. [3, 4].

Let  $\phi_r: \Sigma^n P_n^{n+r} = Q^r(S^n) \rightarrow S^{2n}$  be the  $n$ -fold suspension of such a retraction. Since  $Q^r(Y_0) \simeq Q^r(S^n)$ , there exists a map  $\Phi_r: Q^r(Y_0) \rightarrow W_0$  equivalent to  $\phi_r$ .

Let  $\theta_t: S^t | \times (Y_0 \wedge Y_0) \rightarrow Q^t(Y_0)$  be the identification map. Denote by  $E_+^t (E_-^t)$  the upper (lower) hemisphere of  $S^t$ . Let

$$\psi_t^\pm: (E_\pm^t | \times (Y_0 \wedge Y_0) \rightarrow S^t | \times (Y_0 \wedge Y_0)$$

be the inclusion map.

PROPOSITION (4.2). Let  $n = 2^q$ ,  $q \geq \phi(r)$ . Then there exists a map  $\Theta_r: S^r | \times (Y_0 \wedge Y_0) \rightarrow W_0$  such that

- A)  $\Theta_r \circ T = \Theta_r$ ,
- B) there is an equivariant homotopy  $H^r$  of  $\Theta_r$  to  $\Phi \circ \theta_r$  (yielding  $\Theta_r/Z_2 \simeq \Phi_r: Q^r(Y_0) \rightarrow W_0$ ), and
- C)  $\Theta_r(S^t | \times Z_s) \subset W_{s-t}$  for  $t \leq r$ .

Proof. We proceed by induction, denoting by  $A_t, B_t, C_t$  conditions A, B, C of (4.2) for  $r = t$ .

Since  $\Phi: Q^0(Y_0) = Y_0 \wedge Y_0 \rightarrow W_0$  is equivalent to the identity map  $S^{2n} \rightarrow S^{2n}$ , we may use Lemma 3.4 of [2] to obtain a map

$$\Theta_0^+: \psi_0^+(E_+^0 | \times (Y_0 \wedge Y_0)) \approx Y_0 \wedge Y_0 \rightarrow W_0$$

such that  $\Theta_0^+ \psi_0^+(E_+^0 | \times Z_s) \subset W_s$  and such that there is a homotopy  $H_+^0$  of  $\Theta_0^+$  with  $\Phi_0$ . Define  $\Theta_0^-: \psi_0^-(E_-^0 | \times (Y_0 \wedge Y_0)) \rightarrow W_0$  by  $\Theta_0^- = \Theta_0^+ \circ T \circ \psi_0^-$ . This defines  $\Theta_0$ . We define  $H_-^0$  by the composite

$$\psi_0^-(E_-^0 | \times (Y_0 \wedge Y_0)) \times |I \xrightarrow{T \times 1} \psi_0^+(E_+^0 | \times (Y_0 \wedge Y_0)) \xrightarrow{H_+^0} W_0.$$

This defines  $H^0$ . Conditions  $A_0, B_0$  and  $C_0$  are readily verified.

Assume now that  $\Theta_t$  is defined satisfying  $A_t, B_t$  and  $C_t$  where  $t < r$ . Now since  $t < r$ ,  $\Theta_t/Z_2: Q^t(Y_0) \rightarrow W_0$  extends to a map  $\alpha: Q^{t+1}(Y_0) \rightarrow W_0$  with  $H^t/Z_2$  extending to a homotopy  $K$  of  $\alpha$  with  $\Phi_{t+1}$ . Consider the composite

$$\alpha \circ \theta_{t+1} \circ \psi_{t+1}^+: E_+^{t+1} | \times (Y_0 \wedge Y_0) \rightarrow W_0.$$

Using an argument similar to one given in Lemma 3.5 of [2], we see that  $\alpha \circ \theta_{t+1} \circ \psi_{t+1}^+$  is homotopic relative to  $S^t | \times (Y_0 \wedge Y_0)$  to a map  $\bar{\Theta}_{t+1}^+$  such that  $\bar{\Theta}_{t+1}^+(E_+^{t+1} | \times Z_s) \subset W_{s-(t+1)}$ . Define  $\Theta_{t+1}^+$  by the equation  $\Theta_{t+1}^+ = \bar{\Theta}_{t+1}^+ \circ \psi_{t+1}^+$ . This also yields a homotopy  $H_{t+1}^+$  extending  $H^t$  to  $\Theta_{t+1}^+$ . Now set  $\Theta_{t+1}^- = \Theta_{t+1}^+ \circ T$  and  $H_{t+1}^- = H_{t+1}^+ \circ (T \times 1)$ . One checks that conditions  $A_{t+1}, B_{t+1}$  and  $C_{t+1}$  are satisfied. This completes the induction.

Id  
homon  
 $S^r | \times ($   
an ison

Conside

$(E_\pm^j,$   
Then we

LEM

Proc

The resul

In th

THEC  
an integer  
A)

B) 1  
 $(E^{2(n+t)+k}$   
cycle of  $E_1$   
istic map c

Reman  
an integer

Proof.  
vant portic  
which satis

Now, s  
We define t

By (3.  
(4.4A). Not  
from (3.4) a

In this s  
(4.4) to relat  
tions on perr

Identifying  $H^*(W_k, W_{k+1})$  with  $B_k(A)$  and  $H^*(Z_m, Z_{m+1})$  with  $(B(A) \otimes B(A))_m$ , we define homomorphisms  $D_j: B_n(A) \rightarrow (B(A) \otimes B(A))_{n+j}$ ,  $0 \leq j \leq r$  as follows: Let  $\Theta = \Theta_r: S^r \times (Y_0 \wedge Y_0) \rightarrow W_0$  be a map given by Proposition (4.2). The Künneth Theorem yields an isomorphism

$$\chi_{\pm}: H^*((E_{\pm}^j, S^{j-1}) \times (Z_m, Z_{m+1})) \approx H^{*-j}(Z_m, Z_{m+1}) \approx (B(A) \otimes B(A))_m$$

Consider the composites

$$(E_{\pm}^j, S^{j-1}) \times (Z_m, Z_{m+1}) \xrightarrow{\psi_j^{\pm}} (S^j, S^{j-1}) \times (Z_m, Z_{m+1}) \xrightarrow{\Theta} (W_{m-j}, W_{m-j+1}) \approx B_{m-j}(A).$$

Then we define  $D_j: B_{m-j}(A) \rightarrow (B(A) \otimes B(A))_m$  by

$$D_j = \chi_{+} \circ (\psi_j^{+})^{*} \circ \Theta^{*}.$$

LEMMA (4.3). The homomorphisms  $D_j$  satisfy (2.1) and (2.2) for  $j \leq r$ .

Proof. Since we are working mod 2, it follows by (4.2A) that

$$\rho D_j = \chi \circ (\psi_j^{-})^{*} \circ \Theta^{*}.$$

The result now follows from the cell structure of  $S^r$ .

In the following theorem, we retain the notation of this section.

THEOREM (4.4). Let  $\alpha \in E_2^{s,s+t}(S^0)$  be a permanent cycle and  $r$  an integer. Then there is an integer  $n$  and a mapping  $\bar{\Theta}: \Sigma^{n+t} P_{n+t}^{n+t+r} \rightarrow W_0$  such that

A)  $\bar{\Theta}(\Sigma^{n+t} P_{n+t}^{n+t+k}) \subset W_{2s-k}$ ,  $0 \leq k \leq r$ , and

B) the composite  $(E_1^{2(n+t)+k}, S^{2(n+t)+k-1}) \xrightarrow{\Theta} (\Sigma^{n+t} P_{n+t}^{n+t+k}, \Sigma^{n+t} P_{n+t}^{n+t+k-1}) \xrightarrow{\bar{\Theta}} (W_{2s-k}, W_{2s-k+1})$  represents a cycle of  $E_1^{2s-k, 2(s+t)}(S^0)$  whose image in  $E_2^{2s-k, (s+t)}(S^0)$  is  $\alpha_k \cup \alpha$ . ( $\bar{\Theta}$  denotes the characteristic map of the top cell of  $\Sigma^{n+t} P_{n+t}^{n+t+k}$ .)

Remark (4.5). The integer  $n$  may be chosen to be as large as one likes. In fact, there is an integer  $M$  so that we may choose  $n = 2^q$ ,  $q \geq M$ .

Proof. Choose  $n = 2^q$  where  $q \geq \phi(r)$ . We also choose  $q$  large enough so that the relevant portion of  $B(A)$  is realizable. Then we may find a map  $\Theta: S^r \times (Y_0 \wedge Y_0) \rightarrow W_0$  which satisfies conditions A, B and C of (4.2).

Now, since  $\alpha$  is a permanent cycle, it may be represented by a mapping  $f: S^{n+t} \rightarrow Y_s$ . We define the map  $\bar{\Theta}$  to be the composite  $(\Theta/Z_2) \circ Q^r(f): Q^r(S^{n+t}) \rightarrow W_0$ .

By (3.1),  $Q^r(S^{n+t})$  is homeomorphic with  $\Sigma^{n+t} P_{n+t}^{n+t+r}$ . Proposition (4.2C) implies (4.4A). Note that  $f \wedge f: S^{n+t} \wedge S^{n+t} \rightarrow (Z_{2s}, Z_{2s-1})$  represents  $\alpha \times \alpha$ . (4.4B) now follows from (3.4) and (4.3).

## §5. SOME DIFFERENTIALS IN THE ADAMS SPECTRAL SEQUENCE

In this section, we use information concerning the cell structure of  $P_n^{n+k}$  and Theorem (4.4) to relate the differential in the Adams spectral sequence to some of the squaring operations on permanent cycles. The results for  $Sq_1$  were obtained earlier in [5].

**THEOREM (5.1).** *Let  $\alpha \in E_2^{s+t}(S^0)$  be a permanent cycle. Then if  $k+1 \leq \rho(t+k+1)$ ,  $\text{Sq}_k \alpha$  is also a permanent cycle.*

*Proof.* Recall that if  $m = (2a+1)2^{c+4d}$ ,  $0 \leq c \leq 3$ , then  $\rho(m) = 2^c + 8d$ . Observe that if  $t+k+1 = 2^r(2b+1)$  and  $q > r$ , then  $\rho(2^q+t+k+1) = \rho(t+k+1)$ .

It follows from (4.4) and (4.5) that we need only show  $P_{2^q+t+k}^{2^q+t+k}$  is stably reducible if  $q$  is large enough. But it is known that this is true if and only if  $(k+1) \leq \rho(2^q+t+k+1) = \rho(t+k+1)$  [3].

We shall use the symbols  $\bar{h}_0, \bar{h}_1, \bar{h}_2$  and  $\bar{h}_3$  to denote any elements in  $G_*$  which correspond to the elements  $h_0, h_1, h_2$  and  $h_3$  in  $E_2(S^0)$ , respectively. This means that  $\bar{h}_i$  is a generator of the 2-component of  $G_{2^i-1}$ .

Now let  $f_i : S_{n+t}^{n+t+2^i-1} \rightarrow P_{n+t+2^i-1}^{n+t+2^i-1}$  denote the attaching map of the top cell of  $P_{n+t+2^i}^{n+t+2^i}$ . Let  $g : S_{n+t}^{n+t} \rightarrow P_{n+t}^{n+t+2^i-1}$  denote the inclusion map.

**LEMMA (5.2).** *Let  $n = 2^q$ ,  $q$  large enough, and  $i = 0, 1, 2$  or  $3$ . Then  $[f_i] = g_* \bar{h}_i$  if  $t \equiv 2^{i+1} - 1 \pmod{2^{i+1}}$ .*

*Proof.* This follows from the fact that  $\text{Sq}^{2^i}$  is non-zero in  $P_{n+t+2^i}^{n+t+2^i}$  and that  $P_{n+t+2^i}^{n+t+2^i}$  is  $S$ -reducible under the hypothesis of (5.2).

*Notation.*  $\delta_r$  denotes the differential in  $E_r(S^0)$ .

**THEOREM (5.3).** *Let  $i = 0, 1, 2$  or  $3$ . If  $\alpha \in E_2^{s,t}(S^0)$  is a permanent cycle and  $t \equiv 2^{i+1} - 1 \pmod{2^{i+1}}$ , then  $\text{Sq}_{2^i} \alpha$  survives to  $E_{2^i+1}(S^0)$  and*

$$\delta_{2^i+1}(\text{Sq}_{2^i} \alpha) = h_i \alpha^2.$$

*Proof.* Let  $\bar{\Theta} : \Sigma_{n+t}^{n+t} P_{n+t}^{n+t+2^i} \rightarrow W_0$  be a map given by (4.4) with  $n = 2^q$ ,  $q$  large. Then by (5.2) it follows that  $\text{Sq}_{2^i} \alpha$  is represented by a map  $(E^{2(n+t)+2^i}, S^{2(n+t)+2^i-1}) \xrightarrow{f} (W_{2s-2^i}, W_{2s})$  such that  $f|_{S^{2(n+t)+2^i-1}} = f'$  is homotopic in  $W_{2s}$  to the composite of  $\bar{h}_i$  with a map representing  $\alpha^2$ . It follows that  $f'$  is homotopic through  $W_{2s}$  to a map  $f'' : S^{2(n+t)+2^i-1} \rightarrow W_{2s+1}$  which represents  $h_i \alpha^2$ . (For  $i = 0$ , this is Lemma (2.2) of [5]. The proof of this statement for  $i = 1, 2$  and  $3$  is similar.)

A specific homotopy through  $W_{2s}$  of  $f'$  to  $f''$  can be used to alter  $f$  so as to obtain a mapping

$$g : (E^{2(n+t)+2^i}, S^{2(n+t)+2^i-1}) \rightarrow (W_{2s-2^i}, W_{2s+1})$$

so that  $g$  represents  $\text{Sq}_{2^i} \alpha$  and  $g|_{S^{2(n+t)+2^i-1}}$  represents  $h_i \alpha^2$ . This completes the proof of (5.3).

**THEOREM (5.4).** *Suppose that  $\alpha \in E_2^{s,t}(S^0)$  is a permanent cycle and that both*

(i)  $k+1 \leq \rho(t+k+1)$  and

(ii)  $t+k \equiv 2^{i+1} - 1 \pmod{2^{i+1}}$ ,  $i = 0, 1, 2$  or  $3$ .

*Then  $\text{Sq}_{k+2^i} \alpha$  survives to  $E_{2^i+1}(S^0)$  and*

$$\delta_{2^i+1}(\text{Sq}_{k+2^i} \alpha) = h_i(\text{Sq}_k \alpha).$$

*Proof.* Let  $n = 2^q$ ,  $q$  large enough. Condition (i) implies that (stably)  $P_{n+t+k}^{n+t+k}$  has the same homotopy type as  $P_{n+t+k-1}^{n+t+k-1} \vee S^{n+t+k}$ . Let

$$f : S_{n+t+k+2^i-1}^{n+t+k+2^i-1} \rightarrow P_{n+t+k+2^i-1}^{n+t+k+2^i-1}$$

denote  
topic t

Using t

We

(4.4) w

+  $\bar{\Theta}_*$ [ $\Sigma$

$\pi_*(W_{2s-}$

$\bar{\Theta}_0(\Sigma^{n+1}$

follows

$S^{2(n+t)+}$

Age

We

More pr

Defi

$\beta'$  the im

represent

homomo

The

to Massey

(5.3) are l

We s

Defin

is called  $v_i$

and  $\beta$  is re

The n

LEMMA

which maps

Proof.

The qu  
to define cu

denote the attaching map of the top cell of  $P_{n+t}^{n+t+k+2^i}$ . Lemma (5.2) implies that  $f$  is homotopic to a map  $g: S^{n+t+k+2^i-1} \rightarrow P_{n+t}^{n+t+k}$ . Now

$$\pi_{n+t+k+2^i-1}(P_{n+t}^{n+t+k}) \approx \pi_{n+t+k+2^i-1}(S^{n+t+k}) \oplus \pi_{n+t+k+2^i-1}(P_{n+t}^{n+t+k-1}).$$

Using this decomposition write  $[g] = ([g_1], [g_2])$ . Condition (ii) implies that  $[g_1] = \bar{h}_i$ .

We proceed as in the proof of (5.3). Let  $\bar{\Theta}: \Sigma^{n+t} P_{n+t}^{n+t+k+2^i} \rightarrow W_0$  be a map given by (4.4) with  $n = 2^q$ ,  $q$  large. Then, as in (5.3),  $\delta_{2^i}(\text{Sq}_{k+2^i} \alpha)$  is represented by  $\bar{\Theta}_*[\Sigma^{n+t} g_1] + \bar{\Theta}_*[\Sigma^{n+t} g_2] \in \pi_*(W_{2s-k})$ . Now  $\bar{\Theta}_*[\Sigma^{n+t} g_1]$  is homotopic in  $W_{2s-k}$  to an element of  $\pi_*(W_{2s-k+1})$  which represents  $h_i(\text{Sq}_k \alpha)$ . Because  $\bar{\Theta}_0(\Sigma^{n+t} P_{n+t}^{n+t+k-1}) \subset W_{2s-k+1}$ ,  $\bar{\Theta}_0(\Sigma^{n+t} g_2)(S^{2(n+t)+2^i-1})$  is contained in  $W_{2s-k+1}$ , and since  $\bar{H}^*(g_2; Z_2) = 0$ , it follows by Lemma (3.3) of [2] that  $\bar{\Theta}_0 \Sigma^{n+t} g_2$  is homotopic in  $W_{2s-k+1}$  to a map carrying  $S^{2(n+t)+k+2^i-1}$  into  $W_{2s-k+2}$ .

Again as in the proof of (5.3), it follows that

$$\delta_{2^i+1}(\text{Sq}_{k+2^i} \alpha) = h_i(\text{Sq}_k \alpha).$$

We conclude this section by observing that the differentials given in (5.3) are "honest". More precisely:

**Definition (5.5).** Let  $\beta \in \pi_*(W_m)$  such that the image of  $\beta$  in  $\pi_*(W_0)$  is zero. Denote by  $\beta'$  the image of  $\beta$  in  $E_r^{m,*}(S^0)$ . Then a relation  $\delta_r \gamma' = \beta'$  is called *honest rel.  $\beta$*  if  $\gamma'$  can be represented by an element  $\gamma \in \pi_*(W_{m-r}, W_m)$  such that  $\partial_* \gamma = \beta$ ,  $\partial_*$  being the boundary homomorphism of the pair  $(W_{m-r}, W_m)$ .

The condition of honesty of relations occurs in the problem of relating Toda brackets to Massey products. See, for example, the work of Moss [7]. It is clear that the relations of (5.3) are honest.

We shall need another criterion of honesty in the next section.

**Definition (5.6).** A relation

$$\delta_r^{s-r, t+1+s-r} \gamma' = \beta' \in E_r^{s, s+t}$$

is called *visibly honest rel.  $\beta$*  if

$$\delta_{r+k+l}^{s-r-k, t+1+s-r-k} = 0 \text{ for } k \geq 0, l \geq 1$$

and  $\beta$  is represented by an element  $\tilde{\beta} \in \pi_*(W_r)$  whose image in  $\pi_*(W_0)$  is zero.

The nomenclature of (5.6) is justified by the following:

**LEMMA (5.7).** If  $\delta_r \gamma' = \beta'$  is visibly honest rel.  $\beta'$ , it is honest rel.  $\beta$ , for any  $\beta \in \pi_*(W_s)$  which maps to zero in  $\pi_*(W_0)$  and to  $\pi'$  in  $\beta_r$ .

*Proof.* [7].

## §6. CUP- $i$ PRODUCTS IN HOMOTOPY

The quadratic construction has been used by Adams, Barratt, B. Gray and Mahowald to define cup- $i$  products in  $G_*$ . We do not attempt a systematic discussion of these opera-

tions. Rather, we just treat some illustrative cases and their relation to the Adams spectral sequence.

Let  $f: S^{n+t} \rightarrow S^n$  represent  $\alpha \in G_t$ , where  $n = 2^q$ ,  $q$  large. Then from (4.1) we obtain a mapping

$$\bar{\Theta}_r: Q'(S^{n+t}) = \Sigma^{n+t} P_{n+t}^{n+t+r} \rightarrow S^{2n}$$

which is the composition of a retraction  $\Psi: \Sigma^n P_n^{n+r} \rightarrow S^{2n}$  with  $Q'(f)$ . Denote by  $g: S^{n+t+r-1} \rightarrow P_{n+t}^{n+t+r-1}$  the attaching map of the top cell. Since  $\Psi \circ Q'^{-1}(f)$  extends to  $\Psi \circ Q'(f)$ , it follows that  $\bar{\Theta}_{r-1} \circ \Sigma^{n+t} g$  has a null-homotopy  $H$ . Now if for "different" reasons, there exists a null-homotopy  $K$  of  $\bar{\Theta}_{r-1} \circ \Sigma^{n+t} g$ , we define  $\alpha \cup_r \alpha$  to contain the difference element  $d(H, K)$ . Thus  $\alpha \cup_r \alpha \in G_{2r+r}(\alpha \cup_0 \alpha = \alpha^2)$ . We consider a few examples.

*Examples (6.1).* A.) If  $r+1 \leq \rho(t+r+1)$ , then  $g$  itself is null-homotopic. In this case  $\alpha \cup_r \alpha$  is defined by demanding that the null-homotopy  $K$  be the composite of  $\Psi$  with a null-homotopy  $K'$  of  $\Sigma^{n+t} g$ . The indeterminacy of  $\alpha \cup_r \alpha$  comes from varying the null-homotopy  $K'$  and the retraction  $\Psi$ .

B.) Let  $t \equiv 2^{i+1} - 1 \pmod{2^{i+1}}$  and  $r = 2^i$  for  $i = 0, 1, 2$  or  $3$ . Then we may suppose that  $g: S^{n+t+2^i-1} \rightarrow S^{n+t}$ . Assume that  $\alpha^2 = 0$ . Let  $K'$  be a null-homotopy of  $f \wedge f$ . In this case, we define  $\alpha \cup_r \alpha$  by demanding that  $K$  be the composite of  $K'$  with the cone on  $g$ . The indeterminacy comes from varying the compression of  $g$  into  $S^{n+t}$ , varying  $K'$  and again varying the retraction  $\Psi$ .

C.) We retain the hypothesis of (6.1B) with the exception that we replace the condition  $\alpha^2 = 0$  by the condition that  $f \circ g$  is null-homotopic for some choice of compression  $g: S^{n+t+2^i-1} \rightarrow S^{n+t}$  of the attaching map. Let  $K'$  be such a null-homotopy. We define  $\alpha \cup_r \alpha$  in this case by demanding that  $K$  be of the form  $K' \wedge f$ .

**THEOREM (6.2).** Let  $\bar{\alpha} \in E_2^{s,s+t}(S^0)$  and suppose that  $r+1 \leq \rho(t+r+1)$ . Then there is an element of  $\bar{\alpha} \cup_r \bar{\alpha}$  as defined in (6.1A) which corresponds to  $\alpha \cup_r \alpha \in E_2^{2s-r, 2s+2t}(S^0)$ .

*Proof.* Defining  $d(H, K) \in \bar{\alpha} \cup_r \bar{\alpha}$ , by (4.4)  $H$  may be chosen so as to represent  $\alpha \cup_r \alpha$  and  $K$  may be compressed into  $W_{2s-r+1}$ . Thus  $d(H, K)$  represents  $\alpha \cup_r \alpha$  in  $E_2^{2s-r, 2s+2t}(S^0)$ .

**THEOREM (6.3).** Suppose that  $\bar{\alpha} \in G_t$  corresponds to  $\alpha \in E_2^{s,s+t}(S^0)$  and that  $t = 2^{i+1} - 1 \pmod{2^{i+1}}$ . Suppose also that  $\bar{\alpha}^2 = 0$  and that there is a visibly honest differential  $\delta_m \beta = \alpha^2 \text{ rel. } \alpha^2$ . Then  $\text{Sq}_{2^i} \bar{\alpha}$  contains an element which corresponds to  $\text{Sq}_{2^i} \alpha \frown h_i \beta$ .

*Notation.* If  $\gamma \in E_2^{s,s+k}$  and  $\gamma' \in E_2^{s',s'+k}$ , then  $\gamma \frown \gamma'$  denotes  $\gamma, \gamma + \gamma'$  or  $\gamma'$  as  $s < s'$ ,  $s = s'$  or  $s > s'$ , respectively.

*Proof.* Let  $r = 2^i$ . Using (4.4) and the proof of (5.3), we compress  $\bar{\Theta}_{r-1} \circ \Sigma^{n+t} g$  into  $W_{2s+1}$  with a null-homotopy into  $W_{2s-r}$  representing  $\alpha \cup_r \alpha$ . By (5.7),  $\delta_m \beta = \alpha^2$  is honest. It follows that  $\delta_m h_i \beta = h_i \alpha^2$  is honest rel.  $h_i \alpha^2$ . Use a representative  $K$  of  $h_i \beta$  which gives null-homotopy of  $\bar{\Theta}_{r-1} \circ \Sigma^{n+t} g$  into  $W_{2s+1-m}$ .  $K$  exists because of the honesty of the differential. Now  $d(H, K)$  represents  $\bar{\alpha} \cup_r \bar{\alpha}$  in  $G_*$  and  $\alpha \cup_r \alpha \frown h_i \beta$  in  $E_2(S^0)$ .

**THEOREM (6.4).** Suppose that  $\bar{\alpha} \in G^t$  corresponds to  $\alpha \in E_2^{s,s+t}(S^0)$  and that  $t \equiv 2^{i+1} - 1 \pmod{2^{i+1}}$ . Suppose also that  $\text{Sq}_{2^i} \bar{\alpha}$  is defined as in (6.1C) and that there is a visibly honest

differ.  
 $\text{Sq}_{2^i} \alpha$

T

1. J. F.
2. J. F.  
(195
3. J. F.
4. D. I
5. D. S
6. A. I  
Am.
7. M. I
8. E. F

A

E



differential  $\delta_m \beta = h_i \alpha$  rel.  $h_i \alpha$ . Then  $Sq_{2^i} \bar{\alpha}$  contains an element which corresponds to  $Sq_{2^i} \alpha \neq \beta \alpha$ .

The proof is similar to that of (6.3).

## REFERENCES

1. J. F. ADAMS: On the non-existence of elements of Hopf invariant one, *Ann. Math.* **72**(1960), 20-104.
2. J. F. ADAMS: On the structure and applications of the Steenrod Algebra, *Comment. math. helvet.* **32**(1958), 180-214.
3. J. F. ADAMS: Vector fields on spheres, *Ann. Math.* **75** (1962), 603-632.
4. D. HUSEMOLLER: *Fibre Bundles*, McGraw-Hill, 1966.
5. D. S. KAHN: A differential in the Adams spectral sequence, *Proc. Am. math. Soc.*, to appear.
6. A. LIULEVICIUS: The factorization of cyclic reduced powers by secondary cohomology operations, *Mem. Am. math. Soc.* **42** (1962).
7. M. MOSS: Secondary compositions and the Adams spectral sequence, to appear.
8. E. H. SPANIER: *Algebraic Topology*, McGraw-Hill, 1966.

Northwestern University  
Evanston, Illinois, U.S.A.