

ON THE HANDLEBODY DECOMPOSITION ASSOCIATED TO A LEFSCHETZ FIBRATION

A. KAS

The classical Lefschetz hyperplane theorem in algebraic geometry describes the homology of a projective algebraic manifold M in terms of "simpler" data, namely the homology of a hyperplane section X of M and the vanishing cycles of a Lefschetz pencil containing X . This paper is a first step in proving a diffeomorphism version of the Lefschetz hyperplane theorem, namely a description of the diffeomorphism type of M in terms of "simpler" data.

Let $\tilde{M}$ be the manifold obtained from M by blowing up the axis of a Lefschetz pencil. There is a holomorphic mapping $f: \tilde{M} \rightarrow \mathbb{CP}^1$ which is a Lefschetz fibration, i.e., f has only nondegenerate critical points (in the complex sense). Using the Morse function $z \rightarrow |f(z)|^2$ on $\tilde{M} - f^{-1}(\infty)$, one obtains a handlebody decomposition of $\tilde{M} - f^{-1}(\infty)$ which may be described as follows: Let $X = f^{-1}(a)$ be a regular fiber of f . Choose a system of smooth arcs $\gamma_1, \dots, \gamma_\mu$ starting at a and ending at the critical values of f such that the γ 's are pairwise disjoint except for their common initial point. The γ 's are ordered such that the tangent vectors $\gamma'_1(0), \dots, \gamma'_\mu(0)$ rotate in a counter clockwise manner. To each γ_j one may associate a "vanishing cycle", i.e., an imbedding $\phi_j: S^n \rightarrow X$ ($\dim X = 2n$) defined up to isotopy, together with a bundle isomorphism $\phi'_j: \tau \rightarrow \nu$ where τ is the tangent bundle to S^n and ν is the normal bundle of S^n in X corresponding to the imbedding ϕ_j . ϕ'_j together with the well known bundle isomorphism $\tau \oplus \varepsilon \simeq \varepsilon^{n+1}$ determines a trivialization of the normal bundle of $e^{2\pi i j/\mu} \times \phi_j(S^n)$ in $S^1 \times X$. This trivialization allows one to attach a n -handle to $D^2 \times X$ with the core $e^{2\pi i j/\mu} \times \phi_j(S^n)$. If this is done for each $j, j=1, \dots, \mu$, the resulting manifold is diffeomorphic to $\tilde{M}$ -(tubular neighborhood of $f^{-1}(\infty)$).

Using the bundle isomorphism ϕ'_j and the tubular neighborhood theorem one may identify a closed tubular neighborhood T of $\phi_j(S^n)$ in X with the tangent unit disk bundle to S^n . One may then define a diffeomorphism, up to isotopy, $\delta_j: X \rightarrow X$ with support in T . δ_j is a generalization of the classical Dehn-Lickorish twist. δ_j is the geometric monodromy corresponding to the j th critical value of f . It follows that the composition $\delta_\mu \circ \dots \circ \delta_1$ is smoothly isotopic to the identity $1_X: X \rightarrow X$. A smooth isotopy is given by a smooth arc λ in $\text{Diff}(X)$ joining the identity to $\delta_\mu \circ \dots \circ \delta_1$. The choice of λ , up to homotopy, determines the way in which one closes off $\tilde{M}$ -(tubular neighborhood of $f^{-1}(\infty)$) to obtain $\tilde{M}$.

Thus the diffeomorphism type of $\tilde{M}$ is determined by the invariants $\phi_1, \phi'_1, \dots, \phi_\mu, \phi'_\mu$ and $\{\lambda\}$, the homotopy class of λ . Conversely, given a compact oriented $2n$ dimensional manifold X , imbeddings $\phi_j: S^n \rightarrow X$, $j=1, \dots, \mu$, and bundle isomorphisms $\phi'_j: \tau \sim \nu_j$ such that $\delta_\mu \circ \dots \circ \delta_1$ is smoothly isotopic to 1_X , and a homotopy class $\{\lambda\}$ of arcs in $\text{Diff}(X)$ with initial point 1_X and end point $\delta_\mu \circ \dots \circ \delta_1$, one may construct a $2n+2$ dimensional manifold $\tilde{M}$ and a Lefschetz fibration $f: \tilde{M} \rightarrow \mathbb{CP}^1$. It is shown that in the case $n=1$, apart from certain exceptions, $\tilde{M}$ is uniquely determined by $\phi_1, \dots, \phi_\mu$, i.e., the bundle isomorphisms $\phi'_1, \dots, \phi'_\mu$ and the smooth isotopy class $\{\lambda\}$ are superfluous.

O. Introduction. The classical Lefschetz hyperplane theorem in algebraic geometry describes the homology groups of an algebraic manifold M in terms of those of a hyperplane section X and in terms of the "vanishing cycles" of X . This paper was inspired by the Morse theoretic proof of the Lefschetz hyperplane theorem due to Andreotti and Frankel [1]. Their approach is to blow up the base locus of a generic Lefschetz pencil so as to obtain a manifold $\tilde{M}$ and a "Lefschetz fibration" $f: \tilde{M} \rightarrow P^1(C)$. They then use the Morse function $|f|^2$ to describe $\tilde{M}$, at least up to homotopy type, and finally they show how to relate the homology groups of M to those of $\tilde{M}$. Now according to Smale's handlebody theory [11], it should be possible to use the Morse function $|f|^2$ to determine the full diffeomorphism class of $\tilde{M}$, not just its homotopy type. In order to do this we must describe the framings of the imbedded spheres (vanishing cycles) corresponding to the critical points of $|f|^2$. In general, the framing associated to a critical point of index $n+1$, has an ambiguity measured by $\pi_n(SO(n+1))$. In our situation, we can improve this to $\pi_n(SO(n))$. This completely determines the framing in certain cases, most notably if M is a complex surface.

In this paper, we describe a set of invariants associated to a Lefschetz fibration $f: M \rightarrow P^1$, which allows one, in principle, to give a handlebody decompositions of certain complex surfaces.

There is a certain amount of overlap between some of the ideas of this paper and certain papers of B. Moishezon and R. Mandelbaum (cf. e.g., [6], [7], [10]).

1. The framings associated with a Lefschetz fibration. Let M be a smooth manifold of dimension ≥ 2 and let $f: M \rightarrow S^2$ be a smooth mapping. A point $p \in M$ will be called a critical point of f if the differential $df_p: T(M)_p \rightarrow T(S^2)_{f(p)}$ is not surjective. Now let M be a closed compact oriented smooth manifold of even dimension,

say $\dim M = 2n + 2$, $n \geq 0$, and assume that $f: M \rightarrow S^2$ is a surjective mapping with a finite number of critical points. We will identify S^2 with the extended complex plane $C \cup \infty$. If $z \in S^2$ is a regular value of f , then $f^{-1}(z)$ is called a regular fiber of f . It is clear that up to diffeomorphism, the regular fibers of f are independent of the regular value z .

DEFINITION 1.1. The smooth mapping $f: M \rightarrow S^2$ will be called a Lefschetz fibration if each critical point p of f admits a coordinate neighborhood with complex valued coordinates $(w_1, \dots, w_{n+1})$, consistent with the given orientation of M , and if $f(p)$ has a coordinate neighborhood with a complex coordinate z , consistent with the orientation of S^2 , such that locally, f has the form:

$$f(w) = z_0 + w_1^2 + \dots + w_{n+1}^2.$$

DEFINITION 1.2. Two Lefschetz fibrations

$$f_1: M \longrightarrow S^2, \quad \text{and} \quad f_2: M \longrightarrow S^2$$

are said to be equivalent, if $f_2 = g \circ f_1$ where g is an orientation preserving diffeomorphism of S^2 .

Let X denote any regular fiber of the Lefschetz fibration $f: M \rightarrow S^2$. Notice that up to diffeomorphism, X only depends on the equivalence class of the Lefschetz fibration $f: M \rightarrow S^2$. Notice also, that X has a unique orientation consistent with the orientations of M and S^2 .

We wish to describe a handlebody decomposition of M associated to the Lefschetz fibration $f: M \rightarrow S^2$. We will assume that a handlebody decomposition of X is already known.

We first recall some standard facts about handles and handlebodies. Let N be a manifold with boundary and let $n = \dim N$. Let

$$\Phi: S^{k-1} \times D^{n-k} \longrightarrow \partial N$$

be a smooth imbedding. Form the union $N_1 = N \cup_{\Phi} D^k \times D^{n-k}$ where we identify each point of $S^{k-1} \times D^{n-k} \subset \partial(D^k \times D^{n-k})$ with its image under Φ . N_1 is a manifold with boundary and corner points. Let V denote the unique manifold (possibly with boundary) obtained from N_1 by "straightening the corners" of N_1 (cf. [2]).

DEFINITION 1.3. V is called: the manifold obtained from N by attaching a k -handle along Φ .

It is easy to see that, up to diffeomorphism, V depends only on the smooth isotopy class of Φ . Let Φ_0 denote the restriction of

Φ to $S^{k-1} \times 0$. It follows easily from the tubular neighborhood theorem [9], that Φ is determined, up to smooth isotopy, by a bundle isomorphism:

$$\Phi': \varepsilon^{n-k} \longrightarrow \nu$$

where ε^{n-k} is the trivial $n-k$ bundle on S^{k-1} , and ν is the normal bundle of S^{k-1} in ∂N under the imbedding Φ_0 . Thus the isotopy class of the imbedding Φ is determined by:

(i) A smooth isotopy class of imbeddings

$$\Phi_0: S^{k-1} \longrightarrow \partial N;$$

(ii) For each Φ_0 in (i), a smooth isotopy class of bundle isomorphisms:

$$\Phi': \varepsilon^{n-k} \longrightarrow \nu.$$

Notice that (i) is a "knot invariant". As for (ii), the distinct bundle isomorphisms, up to smooth isotopy, are classified by the group $\pi_{k-1}(SO(n-k))$. Φ' is called a framing of Φ_0 .

Let $F: M \rightarrow \mathbf{R}$ be a Morse function, and let $c \in \mathbf{R}$ be a critical value such that $F^{-1}(c)$ contains a single critical point p , where F has index λ at p . For each real number $a \in \mathbf{R}$, let $M_a = \{x \in M | F(x) \leq a\}$. Then for $\varepsilon > 0$ sufficiently small, $M_{c+\varepsilon}$ is diffeomorphic to the manifold obtained from $M_{c-\varepsilon}$ by attaching a λ -handle along $\Phi: S^{\lambda-1} \times D^{n-\lambda} \rightarrow \partial M_{c-\varepsilon}$. To construct Φ explicitly, one may choose (by Morse's lemma) a system of coordinates $x_1, \dots, x_n$ in M centered at p , such that $F(x_1, \dots, x_n) = c - x_1^2 - \dots - x_\lambda^2 + x_{\lambda+1}^2 + \dots + x_n^2$ (cf. e.g., [8]). Then if $\xi = (\xi_1, \dots, \xi_\lambda) \in S^{\lambda-1}$, $\eta = (\eta_{\lambda+1}, \dots, \eta_n) \in D^{n-\lambda}$, $\Phi: S^{\lambda-1} \times D^{n-\lambda} \rightarrow F^{-1}(c-\varepsilon)$ is defined by setting

$$\Phi(\xi, \eta) = (x_1, \dots, x_n)$$

where

$$\begin{aligned} (x_1, \dots, x_\lambda) &= \sqrt{\varepsilon + |\eta|^2} \xi \\ (x_{\lambda+1}, \dots, x_n) &= \eta. \end{aligned}$$

Similarly, if $F^{-1}(c)$ contains several nondegenerate critical points $p_1, \dots, p_\mu$ of indices $\lambda_1, \dots, \lambda_\mu$, then $M_{c+\varepsilon}$ is diffeomorphic to the manifold obtained from $M_{c-\varepsilon}$ by attaching μ handles, where the j th handle is a λ_j -handle attached along $\Phi_j: S^{\lambda_j-1} \times D^{n-\lambda_j} \rightarrow F^{-1}(c-\varepsilon)$, $j = 1, \dots, \mu$, and where the images of the Φ_j are disjoint.

DEFINITION 1.4. A handlebody decomposition of a manifold M is given by:

(i) a sequence $M = M_N \supset \dots \supset M_0$, where each M_j is a sub-

manifold of M ;

(ii) diffeomorphisms $\Phi_j: S^{\lambda_j-1} \times D^{n-\lambda_j} \rightarrow \partial M_k$, and

$$\begin{aligned}\Psi_{j+1}: M_{j+1} &\longrightarrow M_j \cup_{\Phi_j} D^{\lambda_j} \times D^{n-\lambda_j}, \\ \Psi_0: M_0 &\longrightarrow D^n.\end{aligned}$$

A Morse function $F: M \rightarrow \mathbb{R}$ together with a gradient-like vector field for F , determine a handlebody decomposition of M (cf. [6]).

Now let $f: M \rightarrow S^2$ be a Lefschetz fibration with regular fiber X , where $\dim X = 2n$, $\dim M = 2n + 2$. We will assume that 0 and ∞ are regular values of f . Define $F: M \rightarrow \mathbb{R} \cup \infty$ by $F(p) = |f(p)|^2$. Then it is easy to verify that outside of $f^{-1}(0) \cup f^{-1}(\infty)$, F has only nondegenerate critical points, each of index $n + 1$ (cf. [1]).

DEFINITION 1.5. The Lefschetz fibration $f: M \rightarrow S^2$ is said to be normalized if the critical values of f are precisely the μ roots of unity.

Every Lefschetz fibration is equivalent to a normalized Lefschetz fibration. Explicitly, if $a_1, \dots, a_\mu \in S^2$ are the critical values of f , and if $a \in S^2$ is a regular value of f , choose arcs γ_j from a to a_j , $j = 1, \dots, \mu$, which do not intersect except at a . Construct an orientation preserving diffeomorphism between a regular neighborhood of $\bigcup_{j=1}^\mu \gamma_j$ and the disk of radius $1 + \varepsilon$ such that a_j is mapped to $e^{2\pi i j / \mu}$. Let $g: S^2 \rightarrow S^2$ be an extension of this diffeomorphism. Then $g \circ f: M \rightarrow S^2$ is a normalized Lefschetz fibration.

Now assume that $f: M \rightarrow S^2$ is a normalized Lefschetz fibration, and let $D_t \subset S^2$ denote the disk of radius t centered at the origin. Then $f^{-1}(D_t) \cong X \times D^2$ for $t < 1$, while for $t > 1$, $f^{-1}(D_t)$ is diffeomorphic to the manifold obtained from $X \times D^2$ by attaching μ $(n + 1)$ -handles by means of imbedding $\Phi_j: S^n \times D^{n+1} \rightarrow \partial(X \times D^2) = X \times S^1$, $j = 1, \dots, \mu$, where the Φ_j have disjoint images.

LEMMA 1.6. *The imbeddings Φ_j may be chosen so that $\Phi_j(S^n \times 0) \subset f^{-1}(z_j)$ for some z_j , $j = 1, \dots, \mu$.*

The proof of Lemma 1.6 will be given together with the proof of Theorem 1.7.

Let $\phi_j: S^n \rightarrow X$ be the imbedding defined by Φ_j restricted to $S^n \times 0$, and the identification $f^{-1}(D_t) \cong X \times D^2$, for $t < 1$. Let ν_1 denote the normal bundle of S^n in X corresponding to the imbedding ϕ_j , and let ν denote the normal bundle of S^n in $F^{-1}(1 - \varepsilon)$. Clearly we have $\nu \cong \nu_1 \oplus \varepsilon$. Let τ denote the tangent bundle of S^n .

THEOREM 1.7. *There exists a bundle isomorphism $\phi'_j: \tau \rightarrow \nu_1$. The framing $\phi'_j: \varepsilon^{n+1} \rightarrow \nu$ coincides with the isomorphism.*

$$\varepsilon^{n+1} \xrightarrow{\sim} \tau \oplus \varepsilon \xrightarrow{\phi'_j \oplus 1} \nu_1 \otimes \varepsilon \xrightarrow{\sim} \nu.$$

Moreover ϕ'_j is determined, up to smooth isotopy, by the local behavior of the function f at the critical point p .

The proof of Lemma 1.6 and Theorem 1.7 are based on the following easy lemma.

LEMMA 1.8. *Let $f: \mathbf{R}^n \rightarrow \mathbf{R}$ be a smooth function, and let $a = (a_1, \dots, a_n)$ be a regular point of f . Let $f(x) = f(a) + L(x - a) + R(x - a)$ where $L(x)$ is a linear function and $|R(x - a)| \leq k|x - a|^2$ for $|x - a|$ sufficiently small. Then for $\varepsilon > 0$ sufficiently small, there exists a smooth function $F: \mathbf{R}^n \times I \rightarrow \mathbf{R}$ satisfying:*

- (i) $F(x, 0) = f(x)$;
- (ii) $F(x, t) = f(x)$ for $|x - a| > \varepsilon$;
- (iii) $F(x, 1) = f(a) + L(x - a)$ for $|x - a| < \varepsilon/2$;
- (iv) $x \rightarrow F(x, t)$ has no critical points in $\{|x - a| < \varepsilon\}$.

Proof. We may assume that $x = 0$, and $f(a) = 0$. Let $\rho_\varepsilon(s)$ be a smooth function such that

$$\rho_\varepsilon(s) = \begin{cases} 1, & s < \varepsilon/2 \\ 0, & s > \varepsilon \end{cases}.$$

Let $F(x, t) = (tL(x) + (1 - t)f(x))\rho_\varepsilon(|x|) + f(x)(1 - \rho_\varepsilon(|x|))$. It is obvious that F satisfies conditions (i), (ii), and (iii). To check condition (iv), we restrict our choice of $\rho_\varepsilon(s)$. Let $\rho(s)$ be a smooth function such that $\rho(s) = \begin{cases} 1, & s < 1/2 \\ 0, & s > 1 \end{cases}$. Then set $\rho_\varepsilon(s) = \rho(s/\varepsilon)$. Let M be a constant such that $|\rho'(s)| \leq M$. Then $|\rho'_\varepsilon(s)| \leq M/\varepsilon$. Letting $\vec{V}f = (\partial f/\partial x_1, \dots, \partial f/\partial x_n)$, and defining $F_t(x) = F(x, t)$ we have:

$$\vec{V}F_t(x) = -t\rho_\varepsilon(|x|)\vec{V}R(x) - tR(x)\rho'_\varepsilon(|x|)\frac{\vec{x}}{|x|} + \vec{V}f$$

where $\vec{x} = (x_1, \dots, x_n)$. The first term on the right is small for small values of ε since $\vec{V}R(x) \rightarrow 0$ as $x \rightarrow 0$. As for the second term, we have $|R(x)| < k\varepsilon^2$ and $|\rho'_\varepsilon(x)| \leq M/\varepsilon$ for $|x| < \varepsilon$. Finally, $\vec{V}f(0) \neq 0$ since 0 is a regular point. Therefore for ε sufficiently small $\vec{V}F_t(x) \neq 0$ for $|x| < \varepsilon$.

Proof of Lemma 1.6 and Theorem 1.7. Assume, for simplicity, that $j = 1$. Since the Morse function F is the composition of the Lefschetz fibration $w \rightarrow f(w)$, and the function $z \rightarrow |z|^2$, we can find, by Lemma 1.8, an arc of Morse functions $F_t: M \rightarrow \mathbf{R}$ such that

- (i) F_t coincides with F on $\{w \in M \mid |f(w) - 1| > \varepsilon\}$

$$(ii) \quad F_0 = F$$

$$(iii) \quad F_1(w) = 2\operatorname{Re} f(w) - 1 \text{ on } \{w \in M \mid |f(w) - 1| \leq \varepsilon/2\}.$$

Since F_1 can be connected to F_0 by an arc of Morse functions, the handlebody decomposition associated with F_1 differs from that associated with F_0 by a smooth isotopy. Now if we use local complex coordinates $(w_1, \dots, w_{n+1})$ in a neighborhood of the critical point p_1 such that: $f(w_1, \dots, w_{n+1}) = 1 + w_1^2 + \dots + w_{n+1}^2$, then $F_1(w) = 2\operatorname{Re} f(w) - 1 = 1 + 2|u|^2 - 2|v|^2$ where $w = u + iv$, $u, v \in \mathbb{R}^{n+1}$. Then $\Phi_1(S^n \times 0) = \{(u, v) \mid u = 0, 1 - 2|v|^2 = 1 - \varepsilon\} \subset f^{-1}(\sqrt{1 - \varepsilon})$, which proves Lemma 1.6.

To prove Theorem 1.7, consider the function: $g: S^n \times \mathbb{R}^{n+1} \rightarrow \mathbb{R}$ defined by $g(\xi, \eta) = \xi \cdot \eta$. $g^{-1}(0)$ is the total space of the tangent bundle to S^n ; thus if ν_1 is the normal bundle of $S^n \times 0$ in $g^{-1}(0)$, then ν_1 is naturally isomorphic to $\tau = \tau_{S^n}$. Since g is a fibration, the normal bundle to $g^{-1}(0)$ in $S^n \times \mathbb{R}^{n+1}$ is trivial. Restricting these bundles to $S^n \times 0$, and using the standard metric on $S^n \times \mathbb{R}^{n+1}$, we get the splitting of bundles on S^n , $\varepsilon^{n+1} \cong \tau \oplus \varepsilon$. An easy computation shows that this coincides with the splitting induced by the standard imbedding of S^n in $\mathbb{R}^{n+1}$. Now consider the diagram:

$$\begin{array}{ccc} S^n \times D^{n+1} & \xrightarrow{\phi} & F_1^{-1}(1 - \varepsilon) \\ g \downarrow & & \downarrow \operatorname{Im} f \\ \mathbb{R} & \xrightarrow{\sqrt{2\varepsilon}} & \mathbb{R} \end{array}$$

The diagram does not commute, but it does “commute to first order” on $S^n \times 0$. More precisely, this diagram induces a commutative diagram of bundle maps which proves Theorem 1.7.

DEFINITION 1.8. $\phi_j(S^n) \subset X$ is called a vanishing cycle for the Lefschetz fibration $f: M \rightarrow S^2$. The bundle isomorphism $\phi'_j: \tau \rightarrow \nu$ will be called a normalization of ϕ_j . The pair $\tilde{\phi}_j = (\phi_j, \phi'_j)$ will be called a normalized vanishing cycle. The sequence of normalized vanishing cycles, $(\tilde{\phi}_1, \dots, \tilde{\phi}_\mu)$ will be called an admissible sequence of normalized vanishing cycles.

Note that the bundle isomorphism $\phi': \tau \rightarrow \nu$ preserves or reverses orientations according to the factor $(-1)^{n(n-1)/2}$. Thus, e.g., if X is a real oriented surface, then a normalized vanishing cycle is completely determined by an imbedded circle. If X is an oriented 4 manifold, the “core” of a normalized vanishing cycle is an imbedded S^2 with self intersection -2 .

Consider the case where $\dim M = 4$, and therefore X is an oriented surface. A vanishing cycle is then an imbedded circle in X , and the normalization of each vanishing cycle is unique. If

$f: M \rightarrow S^2$ is a Lefschetz fibration, then for a sufficiently large disk $G \subset S^2$, $f^{-1}(G)$ is diffeomorphic to the manifold obtain from $X \times D^2$ by attaching a finite number of 2-handles using attaching maps $\phi_j: S^1 \times D^2 \rightarrow X \times S^1$, $j = 1, \dots, \mu$, where $\phi_j(S^1 \times 0)$ is an imbedded circle γ_j , in a fiber $X \times c_j$ of $X \times S^1$. Now there is a natural way to trivialize the normal bundle of γ_j in $X \times S^1$ corresponding to the fact that the normal bundle of γ_j in X is trivial, and the normal bundle of X in $X \times S^1$ is trivial. However, our Theorem 1.7 states that the framing $\phi'_j: \varepsilon^2 \rightarrow \nu$ is obtained by identifying the normal bundle of γ_j in X with the tangent bundle τ of γ_j and then using the isomorphism

$$\varepsilon^2 \cong \tau \oplus \varepsilon.$$

If one also pays attention to orientations, it is not hard to see that, relative to the “natural” framing, $\phi'_j: \varepsilon^2 \rightarrow \nu$ has framing -1 . In other words, if we identify ν with ε^2 by the “natural” framing, then $\phi'_j: \varepsilon^2 \rightarrow \nu$ is given by the mapping $S^1 \rightarrow SO(2) \cong S^1$ of degree -1 . This explains the well known fact (??) in algebraic geometry that relative vanishing cycles have self intersection number -1 .

2. Invariants of a Lefschetz fibration. Let $TS^n(1)$ denote the closed tangent unit disk bundle to S^n . We wish to describe a diffeomorphism

$$\delta: TS^n(1) \longrightarrow TS^n(1)$$

such that:

- (i) δ is the identity on the boundary of $TS^n(1)$; and
- (ii) δ is the antipodal map on $S^n \subset TS^n(1)$.

Let $(p, v) \in TS^n(1)$, where $p \in S^n$ and $v \in TS^n_p$ is a tangent vector of length ≤ 1 . Form the geodesic arc on S^n with initial point p , initial velocity v , and length $\pi|v|$, where $|v|$ denotes the length of v . Let q be the end point, and w the terminal velocity of this arc. Then we put $\delta(p, v) = (q^*, w^*)$ where (q^*, w^*) is the image of (q, w) under the antipodal map.


Now let X be a closed manifold of dimension $2n$; let $\phi: S^n \rightarrow X$ be an imbedding, and let $\phi': \tau \rightarrow \nu$ be a bundle isomorphism where τ is the tangent bundle to S^n , and ν is the normal bundle of S^n in X . By the tubular neighborhood theorem, ϕ' induces a diffeomorphism between $TS^n(1)$ and a tubular neighborhood of $\phi(S^n)$ in X . Thus we can apply δ to a tubular neighborhood of $\phi(S^n)$ in X , and after smoothing near the boundary, we can extend δ by the identity to a diffeomorphism of X which we denote

$$\delta_{\phi, \phi'}: X \longrightarrow X.$$


Note that up to smooth isotopy $\delta_{\phi, \phi'}$ only depends on the smooth isotopy class of the imbedding ϕ and the bundle isomorphism ϕ' .

DEFINITION 2.1. $\delta_{\phi, \phi'}$ is called the Dehn twist of X with center (ϕ, ϕ') .

Notice that if $\dim X = 2$, then $\delta_{\phi, \phi'}$ is the classical right handed Dehn twist, i.e., $\delta_{\phi, \phi'}$ maps the cylinder



to



Now let $f: M \rightarrow S^2$ be a Lefschetz fibration with regular fiber X , and let $p \in M$ be a critical point of f . The part of M which lies over the boundary of a small disk about $f(p)$ is then a fiber bundle over S^1 with typical fiber X . It is well known that such a fiber bundle is diffeomorphic to $X \times I / (x, 1) \sim (h(x), 0)$ where $h: X \rightarrow X$ is a diffeomorphism which is uniquely defined up to smooth isotopy. h is the geometric monodromy of the fibering $f: M \rightarrow S^2$, associated with the critical value $f(p)$. Assume now that p is the only critical point with critical value $f(p)$. Then as in §1, we may construct an imbedding $\phi: S^n \rightarrow X$, and normalization $\phi': \tau \rightarrow \nu$.

THEOREM 2.1 (cf. [3], page 148). *The Dehn twist $\delta_{\phi, \phi'}: X \rightarrow X$ is the geometric monodromy of M about $f(p)$.*

Proof. This theorem is proved in [3], but we will give another proof which is more consistent with our point of view. Let $I = [-1, 1]$ and fix an imbedding $\iota: I \rightarrow S^1$, e.g., $t \mapsto e^{i\pi t}$. Now given a Lefschetz fibration $f: M \rightarrow S^2$ with critical value $f(p)$, choose a small circle γ about $f(p)$; then $f^{-1}(\gamma) = Y$ is the manifold obtained from $X \times S^1$ by surgery. The surgery is performed as follows: look at $\phi' \times \iota: TS^n(1) \times I \rightarrow X \times S^1$. $TS^n(1) \times I$ may be identified with $S^n \times D^n$ in a standard way (this involves straightening corners). The surgery consists of removing $S^n \times D^n = \phi' \times \iota(TS^n(1) \times I)$ and sewing it back in by the automorphism of the boundary $S^n \times S^n$ given by $(\xi, \eta) \rightarrow (\eta, \xi)$. Y is given a differentiable structure by identifying Y with $(X \times S^1 - \phi(S^n) \times \{1\}) \cup D^{n+1} \times S^n$ where $(\xi', \eta') \in (D^{n+1} - 0) \times S^n$ is identified with a point of $X \times S^1 - \phi(S^n) \times \{1\}$ by the following map: first send (ξ', η') to $(\xi, \eta) = (\xi' / |\xi'|, |\xi'| |\eta'|) \in S^n \times D^{n+1}$; then send (ξ, η) to $(u, v) \times t \in TS^n(1) \times I$, and finally send this point to $\phi'(u, v) \times$

$\iota(t)$. We may write

$$Y = (X \times S^1 - \phi(S^n) \times \{1\}) \bigcup_h TS^n(1) \times I$$

where $h: TS^n(1) \times I - S^n \times \{0\} \rightarrow X \times S^1 - \phi(S^n) \times \{1\}$ is given as follows: if $(u, v) \times t \in TS^n(1) \times I - S^n \times \{0\}$, then $h((u, v), t) = \phi'(u', v') \times \iota(t)$ where (u', v') are given by:

$$\begin{aligned} u' &= \frac{t}{\sqrt{t^2 + |v|^2}} u + \frac{1}{\sqrt{t^2 + |v|^2}} v \\ v' &= \frac{|v|^2}{\sqrt{t^2 + |v|^2}} u - \frac{t}{\sqrt{t^2 + |v|^2}} v. \end{aligned}$$

Thus the vector $\begin{pmatrix} u' \\ v' \end{pmatrix}$ is the product of the vector $\begin{pmatrix} u \\ v \end{pmatrix}$ by the matrix:

$$\begin{pmatrix} \frac{t}{\sqrt{t^2 + |v|^2}} & \frac{1}{\sqrt{t^2 + |v|^2}} \\ \frac{|v|^2}{\sqrt{t^2 + |v|^2}} & -\frac{t}{\sqrt{t^2 + |v|^2}} \end{pmatrix} = \begin{pmatrix} 1 & 0 \\ 0 & |v| \end{pmatrix} \begin{pmatrix} \cos \theta & \sin \theta \\ -\sin \theta & \cos \theta \end{pmatrix} \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \\ \times \begin{pmatrix} 1 & 0 \\ 0 & |v|^{-1} \end{pmatrix}$$

where $\theta = \arcsin t/\sqrt{t^2 + |v|^2}$. These formulae show how the projection $X \times S^1 - \phi(S^n) \times \{1\} \rightarrow S^1$ extends to a fibration $f: Y \rightarrow S^1$ where f is defined on $TS^n(1) \times I$ by $f((u, v), t) = \iota(t)$.

The mapping $h: TS^n(1) \times I - S^n \times 0 \rightarrow TS^n(1) \times I - S^n \times 0$ given by $(u, v, t) \mapsto (u', v', t)$ has the following geometric description: Given $u \in S^n$ and $v \neq 0$, v tangent to S^n at u , define (u', v') so that u' lies on the great circle through u in the direction v and the angle between u and u' is $\theta + \pi/2$. v' is tangent to the great circle, has the same length as v and points backwards towards u . For fixed t , as v approaches 0, $\theta = \arcsin t/\sqrt{t^2 + |v|^2}$ approaches $\pi/2$ for $t > 0$ and $-\pi/2$ for $t < 0$. Thus $h|_{S^n \times \{t\}}$ is the identity for $t < 0$ and the antipodal map for $t > 0$. For $|v| = 1$, $\theta = \theta(t)$ goes from $-\pi/4$ to $\pi/4$ as t goes from -1 to 1 .

Now to calculate the geometric monodromy of $f: Y \rightarrow S^1$, we construct a flow on Y . This is accomplished by constructing a smooth map $g: X \times [\varepsilon, 2\pi + \varepsilon] \rightarrow Y$ such that $g(x, s) \in f^{-1}(e^{is})$ and $g_s: X \rightarrow f^{-1}(e^{is})$ is a diffeomorphism (the flow is $g_*(\partial/\partial s)$). The geometric monodromy is given by $g_{2\pi+\varepsilon} \circ g_\varepsilon^{-1}$.

Identifying Y with

$$X \times S^1 - \phi(S^n) \times \{1\} \bigcup_h TS^n(1) \times I$$

we first define g on $X \times [\varepsilon, 2\pi + \varepsilon] - \phi'(TS^n(1/2)) \times [2\pi - \varepsilon/2, 2\pi + \varepsilon]$ by setting $g(x, s) = (x, e^{is})$. We extend g by a mapping $\tilde{g}: TS^n(1) \times I \rightarrow TS^n(1) \times I$ such that on $TS^n(1) \times I - TS^n(1/2) \times [-1/2, 1]$ $h \circ g = \text{id}$. To construct $\tilde{g}$, choose a smooth function $\zeta(|v|, t)$ such that: (i) $\zeta(|v|, t) = \theta(|v|, t)$ on $\{t \leq -1/2\} \cup \{|v| \geq 1/2\}$ and (ii) for fixed t , $\zeta(|v|, t) \rightarrow -\pi/2$ as $|v| \rightarrow 0$. Now define $\tilde{g}$ just as we defined h , but use ζ instead of θ . Specifically,

$$\tilde{g}((u, v), t) = \begin{cases} ((u, v), t) & \text{if } v = 0 \\ ((u', v'), t) & \text{if } v \neq 0 \end{cases}$$

where u' lies on the great circle through u in the direction v , the angle between u and u' is $\pi/2 + \zeta$, and v' points backwards towards u along this geodesic and has the same length as v .

To calculate the monodromy $g_{2\pi+\varepsilon} \circ g_\varepsilon^{-1}$, it is clear that the monodromy is concentrated on $TS^n(1) \subset X$, i.e., it is the identity outside of $TS^n(1)$. Notice that g_ε is defined by the following diagram:

$$\begin{array}{ccc} X = X - S^n \bigcup_{\text{id}} TS^n(1) & & \\ g_\varepsilon \downarrow & \text{id} \downarrow & \downarrow h_1^{-1} \\ f^{-1}(e^{i\varepsilon}) = X - S^n \bigcup_{h_1} TS^n(1) & & \end{array}$$

where $h_t = h|_{TS^n(1) \times t} \cdot g_{2\pi+\varepsilon}$ is defined by:

$$\begin{array}{ccc} X = X - S^n \bigcup_{\text{id}} TS^n(1) & & \\ g_{2\pi+\varepsilon} \downarrow & \downarrow \text{id} & \downarrow \tilde{g}_1 \\ f^{-1}(e^{i\varepsilon}) = X - S^n \bigcup_{h_1} TS^n(1) . & & \end{array}$$

Thus the geometric monodromy $g_{2\pi+\varepsilon} \circ g_\varepsilon^{-1}$ is concentrated on $TS^n(1) \subset X$ and there it is equal to $\tilde{g}_1^{-1} \circ h_1$. This mapping sends a point $(u, v) \in TS^n(1)$ to (u', v') where u' lies on the great circle through u tangent to v , $|v'| = |v|$, and the angle between u and u' is $\phi(|v|, 1) - \zeta(|v|, 1)$. This angle is a decreasing function of $|v|$, equal to π for $|v| = 0$ and equal to 0 for $|v| = 1$. This map is therefore isotopic to the Dehn twist $\delta_{\phi, \phi'}$.

Let $f: M \rightarrow S^2$ be a normalized Lefschetz fibration with fiber X , and critical values $e^{2\pi i j / \mu}$, $j = 0, 1, \dots, \mu - 1$. Let E denote the subset of the disk of radius $1 + \varepsilon$ obtained by removing the sets E_j , $j = 1, \dots, \mu$ where E_j is described in polar coordinates (r, θ) by:

$$E_j = \{(r, \theta) | 1 - \varepsilon < r \leq 1 + \varepsilon, \frac{2\pi j}{\mu} - \varepsilon < \theta < \frac{2\pi j}{\mu} + \varepsilon\}.$$

Identifying $f^{-1}(E)$ with $X \times E$, one may construct a vector field σ on $f^{-1}\{|z| = 1 + \varepsilon\}$ such that $\sigma = (0, \partial/\partial\theta)$ outside of the E_j , while on E_j the flow along σ maps the fiber over $\theta = 2\pi j/\mu - \varepsilon$ to the fiber over $\theta = 2\pi j/\mu + \varepsilon$ by a Dehn twist δ_j , $j = 1, \dots, \mu$. In particular, if we let X_θ denote the fiber over θ , then the flow along σ defines diffeomorphisms,

$$h_\theta: X_0 \longrightarrow X_\theta \text{ such that ,}$$

$$h_{2\pi} = \delta_\mu \circ \dots \circ \delta_1. \text{ Since } f \text{ has no critical}$$

values outside the unit circle, the fibration defined by f may be trivialized over the complement of E . In particular each fiber X_θ is then identified with $X = X_0$, and $\theta \rightarrow h_\theta$ may be regarded as an arc in $\text{Diff}(X)$ joining the identity to $\delta_\mu \circ \dots \circ \delta_1$. Any two such trivializations of the fibering f are related by a diffeomorphism

$$D^2 \times X \longrightarrow D^2 \times X$$

of the form: $(z, x) \rightarrow (z, g_z(x))$. The function $D^2 \rightarrow \text{Diff}(X)$ given by $z \rightarrow g_z$ gives a homotopy between the arcs $\theta \rightarrow h_\theta$, corresponding to the two trivializations. Thus the homotopy class of the arc $\theta \rightarrow h_\theta$ is an invariant of the normalized Lefschetz fibration $f: M \rightarrow S^2$.

THEOREM 2.3. *The normalized Lefschetz fibration $f: M \rightarrow S^2$ is uniquely determined by:*

(1) *a sequence of normalized vanishing cycles $(\tilde{\phi}_1, \dots, \tilde{\phi}_\mu)$; and*

(2) *a homotopy class of arcs $\theta \rightarrow h_\theta$ in $\text{Diff}(X)$, where $h_0 = \text{id}_X$ and $h_{2\pi} = \delta_\mu \circ \dots \circ \delta_1$, where $\delta_j = \delta_{\tilde{\phi}_j}$ is the Dehn twist of X with center $\tilde{\phi}_j$.*

Proof. This theorem is essentially a direct consequence of Theorem 1.7, and our above discussion. More precisely, the manifold M and the fibration f are constructed as follows: starting with $D^2 \times X$, one attaches μ $(n+1)$ -handles along the centers $\Phi_{j_0}: S^n \rightarrow \varepsilon_j \times X$, $\varepsilon_j = e^{2\pi i j/\mu}$, where Φ_{j_0} coincides with $\phi_j: S^n \rightarrow X$, and the framing of Φ_{j_0} is given by the prescription of Theorem 1.7. The boundary of the resulting manifold fibers over a circle, with fiber X , and by Theorem 2.1, one may construct a vector field on this boundary whose flow gives rise to the diffeomorphism $\delta_\mu \circ \dots \circ \delta_1$. The smooth arc $\theta \rightarrow h_\theta$ in $\text{Diff}(X)$ joining id_X to $\delta_\mu \circ \dots \circ \delta_1$ is then used to identify the boundary with $S^1 \times X$, and we finally use this identification to close up the manifold by attaching a copy of $D^2 \times X$.

The normalized vanishing cycle $\tilde{\phi}$ was defined (Def. 1.8) as a

pair (ϕ, ϕ') where $\phi: S^n \rightarrow X$ is an imbedding, and $\phi': \tau_{S^n} \rightarrow \nu$ is a bundle isomorphism. ϕ and ϕ' are defined up to smooth isotopy. The distinct bundle isomorphisms $\tau_{S^n} \rightarrow \nu$ are classified, up to isotopy, by the group $\pi_n(SO(n))$. Thus for those values of n for which $\pi_n(SO(n)) = 0$, ϕ' is uniquely determined. This is true, in particular, for $n = 1$ and 2 .

Given an admissible sequence $(\tilde{\phi}_1, \dots, \tilde{\phi}_\mu)$, let M_1 be the manifold with boundary obtained by attaching μ $(n+1)$ -handles to $D^2 \times X$ as in Theorem 1.7. The distinct homotopy classes of arcs $\theta \rightarrow h_\theta$ in $\text{Diff}(X)$ joining id_X to $\delta_\mu \circ \dots \circ \delta_1$ are classified by $\pi_1(\text{Diff}(X))$. Now a closed loop in $\text{Diff}(X)$ gives rise to a fiber preserving diffeomorphism of $S^1 \times X$, and a homotopy class corresponds to an isotopy class of such fiber preserving diffeomorphisms. Let $\{\theta \rightarrow h_\theta\}, \{\theta \rightarrow h'_\theta\}$ be two homotopy classes of arcs in $\text{Diff}(X)$ joining id_X to $\delta_\mu \circ \dots \circ \delta_1$ and $g: S^1 \times X \rightarrow S^1 \times X$ be a diffeomorphism representing their difference. If g can be extended to a fiber preserving diffeomorphism of M_1 , then the Lefschetz fibering, $f: M \rightarrow S^2$ associated to h_θ is equivalent to the Lefschetz fibration $f': M' \rightarrow S^2$ associated to h'_θ .

THEOREM 2.4. *If X is an oriented surface of genus > 1 , then every Lefschetz fibration $f: M \rightarrow S^2$ with fiber X is uniquely determined, up to equivalence, by an admissible sequence of vanishing cycles. In case X has genus 1, the theorem is still true provided that $f: M \rightarrow S^2$ has at least one vanishing cycle $\phi: S^1 \rightarrow X$ which is not homologous to zero.*

Proof. It is clear that there is a unique normalization for any vanishing cycle in a surface (cf. the remarks on orientation, following Def. 1.8). The theorem in the case of genus $(X) > 1$ follows from the fact that $\text{Diff}(X)$ is contractible (cf. [4]). The case, genus $(X) = 1$, follows from results of Moishezon [10] (Lemma 8 on page 179 and Lemma 7 on page 164).

It follows from Theorem 2.4, that to each equation of the form:

$$\delta_\mu \circ \dots \circ \delta_1 = 1$$

in the mapping class group of the oriented surface X , where δ_j is the right-handed Dehn twist of X , centered at some circle γ_j in X , one may associate a 4-manifold M and a normalized Lefschetz fibration $f: M \rightarrow S^2$ whose critical values are the μ roots of unity and whose vanishing cycles are $\gamma_1, \gamma_2, \dots, \gamma_\mu$. Let $g: S^2 \rightarrow S^2$ be an orientation preserving diffeomorphism which leaves invariant the μ roots of unity. Then $f' = g \circ f: M \rightarrow S^2$ is a normalized Lefschetz fibration with vanishing cycles $\gamma'_1, \dots, \gamma'_\mu$. Following [10], page 180, we

will say that the admissible sequence $(\gamma'_1, \dots, \gamma'_\mu)$ is equivalent to $(\gamma_1, \dots, \gamma_\mu)$. This relation is generated by the elementary transformations:

$$T_j: (\gamma_1, \dots, \gamma_\mu) \longrightarrow (\gamma'_1, \dots, \gamma'_\mu)$$

where:

$$\begin{aligned} \gamma'_j &= \delta_j^{-1}(\gamma_{j+1}) \\ \gamma'_{j+1} &= \gamma_j \\ \gamma'_k &= \gamma_k \quad \text{if } k \neq j, j+1 \end{aligned}$$

(here the subscript j is computed modulo μ). The corresponding Dehn twists δ'_k are given by:

$$\begin{aligned} \delta'_j &= \delta_j \circ \delta_{j+1} \circ \delta_j^{-1} \\ \delta'_{j+1} &= \delta_j \\ \delta'_k &= \delta_k \quad \text{if } k \neq j \text{ or } j+1. \end{aligned}$$

EXAMPLE 2.5. Let $X = T^2$ be a surface of genus 1, i.e., a 2-dimensional torus. Let m denote a standard meridian circle, and let l denote a standard longitude on X . Let $(\gamma_1, \dots, \gamma_\mu)$ be an admissible sequence of imbedded circles on X , such that no γ_j is homologous to zero. Then Moishezon has proved (cf. [10]) that $\mu \equiv 0 \pmod{12}$ and $(\gamma_1, \dots, \gamma_\mu)$ is equivalent to the sequence $(m, l, \dots, m, l)$ (μ terms).

3. Some open problems. In this section we will discuss several problems on Lefschetz fibrations of 4-manifolds. Such fibrations arise naturally from Lefschetz pencils. Thus let M be an algebraic surface imbedded algebraically in a complex projective space P^N . Let H^{N-2} be a generic linear space of codimension 2 in P^N , and let $\tilde{M}$ be the manifold obtained from M by blowing up the points of $M \cap H^{N-2}$. Then there is a Lefschetz fibration $f: \tilde{M} \rightarrow P^1$ such that $f^{-1}(t) \cong M \cap L_t$, where $L_t \subset P^N$ is a hyperplane containing H^{N-2} (such hyperplanes are naturally parametrized by P^1). For the details of this construction cf [1]. By Theorem 2.4, the differentiable structure of $\tilde{M}$ together with the Lefschetz fibration is completely determined by an admissible sequence of vanishing cycles. M is obtained from $\tilde{M}$ by blowing down certain exceptional curves. These exceptional curves are holomorphic sections of the fibration $f: \tilde{M} \rightarrow P^1$ with self intersection -1 . Moreover if we assume that M is a minimal surface and that M is neither rational nor ruled, then M is obtained from $\tilde{M}$ by blowing down all holomorphic sections of $f: \tilde{M} \rightarrow P^1$ of self intersection -1 . We do not know whether every continuous section with self intersection -1 is homotopic to

a holomorphic section. This leads to the following problem:

Problem 3.1. Is the differentiable structure of an algebraic surface M (minimal, not rational or ruled) determined by an admissible sequence of vanishing cycles arising from a Lefschetz pencil?

Let $f_1: M_1 \rightarrow S^2$ and $f_2: M_2 \rightarrow S^2$ be two Lefschetz fibrations whose regular fibers are oriented surfaces of genus g . Choose regular fibers $f_1^{-1}(a) \in M_1$ and $f_2^{-1}(b) \in M_2$, and a diffeomorphism $\alpha: f_1^{-1}(a) \rightarrow f_2^{-1}(b)$. We may construct a new Lefschetz fibration $f_1 \#_\alpha f_2: M_1 \#_\alpha M_2 \rightarrow S^2$ as follows: Let $T_1 \cong f_1^{-1}(a) \times D^2$ be a tubular neighborhood of $f_1^{-1}(a)$ in M_1 , and let $T_2 \cong f_2^{-1}(b) \times D^2$ be a tubular neighborhood of $f_2^{-1}(b)$ in M_2 . Then $M_1 \#_\alpha M_2$ is the union of $M_1 - \text{int}(T_1)$ with $M_2 - \text{int}(T_2)$ where we identify the boundaries by the mapping $f_1^{-1}(a) \times S^1 \rightarrow f_2^{-1}(b) \times S^1$ which sends (x, z) to $(\alpha(x), \bar{z})$. The mapping $f_1 \# f_2$ is defined in an obvious way. $f_1 \# f_2: M_1 \#_\alpha M_2 \rightarrow S^2$ will be called a fiber connected sum of $f_1: M_1 \rightarrow S^2$ and $f_2: M_2 \rightarrow S^2$. In general $M_1 \#_\alpha M_2$ will depend on the diffeomorphism α . It is not hard to see that if $\alpha, \beta: f_1^{-1}(a) \rightarrow f_2^{-1}(b)$ are two diffeomorphisms, then $M_1 \#_\alpha M_2$ is equivalent (in the sense of Lefschetz fibrations) to $M_1 \#_\beta M_2$ if either $\alpha^{-1} \circ \beta: f_1^{-1}(a) \rightarrow f_1^{-1}(a)$ is in the image of the (geometric) monodromy group of M_1 , or if $\alpha \circ \beta^{-1}: f_2^{-1}(b) \rightarrow f_2^{-1}(b)$ is in the image of the (geometric) monodromy group of M_2 .

DEFINITION 3.2. We will say that a Lefschetz fibration $f: M \rightarrow S^2$ is irreducible if it is not equivalent to a fiber connected sum $f_1 \# f_2: M_1 \#_\alpha M_2 \rightarrow S^2$ where both f_1 and f_2 have at least one critical point. Algebraically, an irreducible Lefschetz fibration with regular fiber X is determined by an equation: $\delta_\mu \cdots \delta_1 = 1$ in the mapping class group of X such that $\delta_\mu \cdots \delta_1$ is not equivalent (in the sense given at the end of § 2) to $\delta'_\mu \cdots \delta'_{\nu+1} \delta'_\nu \cdots \delta'_1$ where $\delta'_\nu \circ \cdots \circ \delta'_1 = 1$ and $\delta'_\mu \circ \cdots \circ \delta'_{\nu+1} = 1$.

Problem 3.3. For a given surface X and integer $\mu > 0$, are there only finitely many (up to equivalence) irreducible Lefschetz fibrations $f: M \rightarrow S^2$ with μ vanishing cycles.

REMARK 3.4. The result of Moishezon (Example 2.5) states that if genus $X = 1$, then there is only one irreducible Lefschetz fibration $f: M \rightarrow S^2$ where the number μ of critical points, is 12. In this case, every element of the mapping class group of X is realized by a geometric monodromy, and therefore every Lefschetz fibration of genus 1 is equivalent to $M \# \cdots \# M$.

If the answer to problem 3.1 is "yes", we would like to use Lefschetz pencils to study the diffeomorphism types of algebraic

surfaces. Specifically, let M be an algebraic surface, minimal, not rational or ruled, and let $f: \tilde{M} \rightarrow P^1$ be a Lefschetz fibration arising from a Lefschetz pencil on M . We have given a procedure for obtaining a handlebody decomposition of $\tilde{M}$. We would like to have such a decomposition for M . This could be done if it were possible to rearrange our handlebody decomposition, by sliding handles, so that the sections of $f: \tilde{M} \rightarrow P^1$ of self intersection -1 would occur as the cores of 2-handles.

Problem 3.4. Can the handlebody decomposition of $f: \tilde{M} \rightarrow P^1$ be altered so as to obtain a handlebody decomposition of M ?

REFERENCES

1. A. Andreotti and T. Frankel, *The Second Lefschetz Theorem on Hyperplane Section*, Global Analysis, Princeton University Press (1969), 1-20.
2. A. Douady, *Arrondissement des Arrêtes*, Sem. H. Cartan, **14** (1961/62), No. 3.
3. Deligne and Katz, *Groupes de Monodromie en Géométrie Algébrique*, Lecture Notes in Math. 340, Springer-Verlag.
4. C. J. Earle and J. Eells, *The diffeomorphism group of a compact Riemann surface*, Bull. Amer. Math. Soc., **73** No. 4 (1967), 557-560.
5. Harer, Kas and Kirby to appear.
6. R. Mandelbaum and B. Moishezon, *On the Topological Structure of Non-Singular Algebraic Surfaces in CP^3* , Topology, **15** (1976), 23-40.
7. R. Mandelbaum, *Special handlebody decompositions of simply connected algebraic surfaces*, to appear.
8. J. Milnor, *Morse theory*, Annals of Math., **51**, Princeton Univ. Press, (1963).
9. ———, *Lectures on the h-cobordism theorem*, Princeton Mathematical Notes, (1965).
10. B. Moishezon, *Complex Surfaces and Connected Sums of Complex Projective Planes*, Lecture Notes in Math., 603 Springer-Verlag.
11. S. Smale, *Generalized Poincaré's conjecture in dimensions greater than Four*, Ann. of Math., **74** No. 2 (1961).

Received November 1, 1978. During the preparation of this work, the author was partly supported by N.S.F., Grant MCS77-05583.

OREGON STATE UNIVERSITY
CORVALLIS, OR 97331