

The Euler Characteristic is the Unique Locally Determined Numerical Homotopy Invariant of Finite Complexes

Norman Levitt

Department of Mathematics, Rutgers University,
New Brunswick, NJ 08903, USA

Abstract. If a numerical homotopy invariant of finite simplicial complexes has a local formula, then, up to multiplication by an obvious constant, the invariant is the Euler characteristic. Moreover, the Euler characteristic itself has a unique local formula.

1. Introduction

The Euler characteristic χ is the best known as well as the most ancient topological invariant. For a finite simplicial complex K (or, more generally, a C-W complex) there is the familiar definition

$$\chi(K) = \sum_{i=0}^{\dim K} (-1)^i c_i,$$

where c_i = number of i -simplices (or i -cells if K is a C-W complex.) That $\chi(K)$ is an invariant of homotopy type follows from the alternative definition

$$\chi(K) = \sum_{i=0}^{\dim K} (-1)^i \operatorname{rank} H_i(K; \mathbb{Z}).$$

It is well known and easily verified that $\chi(K)$ is *locally determined* in the sense that given K , we may assign to each vertex $v \in K$ a rational number $e_1(v)$ such that $\chi(V) = \sum_v e_1(v)$. Here, $e_1(v)$ depends only on the simplicial structure of star

$v = \bigcup_{v \in \sigma} \sigma$ (σ a simplex of K) and is given by

$$e_1(v) = \sum_i \frac{(-1)^i}{i+1} \cdot s_i(v),$$

where $s_i(v)$ is the number of i -simplices of K containing v . Since $\text{star } v$ is, simplicially, the cone $c(\text{link } v)$, we may think of $e_1(v)$ as an invariant of the simplicial isomorphism type of $\text{link } v = \bigcup_{v \notin \sigma \subset \text{star } v} \sigma$, i.e.,

$$e_1(v) = e(\text{link } v) = 1 + \sum_{i=0}^{\dim \text{link } v} \frac{(-1)^{i+1}}{i+2} \cdot (\text{number of } i\text{-simplices of link } v).$$

Of course, there are countless other $\mathbb{Z}$ -valued (or $\mathbb{R}$ -valued) invariants of finite complexes of finite complexes. It seems natural to ask whether any of these, other than the Euler characteristic, is locally determined in this sense. Specifically, let ρ denote any $\mathbb{R}$ -valued homotopy invariant of finite complexes. We always assume, by way of normalization, that $\rho(\emptyset) = 0$. Consider a real-valued function $d(L)$ defined on the set of finite simplicial complexes and depending only on the simplicial isomorphism type of L . We say that ρ is *locally determined* by d if and only if given any finite simplicial complex K we have $\rho(K) = \sum_{v \in K} d(\text{link } v)$, where the sum is taken over the vertices v of K . Clearly, the example we have in mind is the Euler characteristic χ , locally determined by e as above, and our question is whether there are any other numerical homotopy invariants (in a nontrivial sense) which are locally determined. The answer turns out to be negative.

Theorem A. *Let ρ be any $\mathbb{R}$ -valued homotopy invariant of finite complexes locally determined by some function d on simplicial-isomorphism classes of finite complexes. Then $\rho = \rho(\text{pt.}) \cdot \chi$.*

In other words, up to multiplication by a constant, χ is the unique locally determined homotopy invariant.

We prove Theorem A in the following form:

Theorem A'. *If ρ is an $\mathbb{R}$ -valued homotopy invariant of finite complexes locally determined by d and such that $\rho(\text{pt.}) = 1$, then $\rho \equiv \chi$.*

Theorem A obviously implies Theorem A' and is, in turn, implied by it for the following reason: Let ρ be as in the statement of Theorem A. If $\rho(\text{pt.}) \neq 0$, replace ρ by $\rho' = \rho/\rho(\text{pt.})$ and apply Theorem A' to conclude $\rho' = \chi$, hence $\rho = \rho(\text{pt.}) \cdot \chi$. If, however, $\rho(\text{pt.}) = 0$ let $\rho' = \rho + \chi$. Applying Theorem A' to ρ' , we have $\rho' = \chi$ hence $\rho = 0 = \rho(\text{pt.}) \cdot \chi$.

The author is indebted to the referee for pointing out that the techniques below will, in fact, lead to a somewhat stronger result.

Consider compact PL n -manifolds (not necessarily closed). Let ρ now denote a real-valued PL-homeomorphism invariant of such manifolds. Let d be a

real-valued function defined on triangulations of S^{n-1} and D^{n-1} . Then the notion of ρ being locally determined by d transcribes, in an obvious way, to this context from the definition given above. Corresponding to Theorem A' we have

Theorem A". *If ρ is an $\mathbb{R}$ -valued invariant of compact PL n -manifolds with $\rho(\emptyset) = 0$, $\rho(D^n) = 1$ and ρ is locally determined by some function d , then $\rho = \chi$.*

Note that Theorem A" does, in fact, imply Theorem A'. Let ρ be a numerical homotopy invariant of finite complexes with $\rho(\emptyset) = 0$, $\rho(\text{pt.}) = 1$. Then, for any n , ρ is, *a fortiori*, a PL-homeomorphism invariant of compact PL n -manifolds with $\rho(D^n) = 1$. If ρ is locally determined, then Theorem A" tells us that $\rho(M^n) = \chi(M^n)$ for compact PL manifolds M^n (n arbitrary). However, given a finite complex K , there exists a compact manifold M^n with K homotopically equivalent to M^n (see, e.g., [W1]). Hence $\rho(K) = \rho(M^n) = \chi(M^n) = \chi(K)$.

The observation above notwithstanding, we shall, in the interest of simplicity of exposition, prove Theorem A' directly first and then show how Theorem A" follows by a straightforward modification of the proof.

If we now go on to ask how many functions d , in addition to the e given above, locally determine χ , we find, in fact, that an even greater degree of rigidity prevails than is asserted by Theorem A. Not only is χ the only locally determined homotopy invariant which evaluates to 1 on a point but, as well, there is only one function, namely $e(L)$, which determines it. We rephrase this:

Theorem B. *If χ is locally determined by d , then $d = e$.*

Some remarks before we proceed to the proofs: If we examine more restricted classes of finite complexes, Theorem A no longer holds. For instance, if we look at the class of triangulated, oriented closed $4k$ -manifolds M , then the signature of M , certainly an invariant of orientation-preserving homotopy type within this class of spaces, is locally determined by a function defined on simplicial-isomorphism classes of triangulated, oriented $(4k - 1)$ -spheres. This is a special case of the fact that rational Pontrjagin classes (as well as other PL characteristic classes) are locally determined. See [C], [L], [LR], and [GM] for details.

Moreover, as the referee has astutely pointed out, Theorem B fails as well in the context of closed PL n -manifolds. Given a combinatorial triangulation of such a manifold M , let c_i denote the number of i -simplices. Klee [K] discovered a family of algebraic relations among the c_i valid for all M and Wall [W2] showed this list to be exhaustive. In consequence, there are nonstandard formulae for $\chi(M)$ in terms of c_i differing from the classical formula cited in the first paragraph. For example (and, once more, the author is indebted to the referee for the observation),

$$\chi(M^n) = c_0 + \sum_{j=1}^{\lfloor (n+1)/2 \rfloor} (-1)^j 2B_{2j} c_{2j-1},$$

where B_{2j} is the $2j$ th Bernoulli number. From this it immediately follows that if we set

$$d(L) = 1 + \sum_{j=0}^{\lfloor (n-1)/2 \rfloor} \frac{(-1)^{j+1}}{j+1} B_{2j+2} \text{ (number of } 2j \text{ simplices of } L),$$

then d locally determines χ on closed n -manifolds.

A word concerning notation: We abbreviate star v by $\text{st } v$ and link v by $\text{lk } v$. If there is any ambiguity as to which ambient complex K is under consideration, we resolve this by recourse to subscripts, e.g. $\text{lk}_K v$, $\text{st}_K v$, etc.

Finally, we note that our proof applies to $\mathbb{C}$ -valued or $\mathbb{Q}$ -valued invariants as well.

2. Proof of Theorem A'

Theorem A' follows immediately from:

Lemma 1. *Let ρ be a numerical homotopy invariant of finite complexes locally determined by some function d . Let K be a finite simplicial complex with $K = K_0 \cup K_1$, $K_0 \cap K_1 = K_2$ where K_0, K_1, K_2 are subcomplexes of K . Then $\rho(K) = \rho(K_0) + \rho(K_1) - \rho(K_2)$.*

The derivation of Theorem A' from Lemma 1 comes via a straightforward induction. Given Lemma 1 and the hypothesis that $\rho(\text{pt.}) = 1$, it is immediate that for a 0-complex (i.e., discrete finite set) K , $\rho(K) = \text{number of points of } K = \chi(K)$. So assume, inductively, that $\rho = \chi$ holds for complexes of dimension $\leq k$ and for $(k+1)$ -complexes having $\leq j$ $(k+1)$ -simplices. Let K be a $(k+1)$ -dimensional complex with exactly $(j+1)$ $(k+1)$ -simplices. Choose a $(k+1)$ -simplex σ . Let $K_0 = \sigma$, $K_1 = K - \text{int } \sigma$, so that $K_2 = K_0 \cap K_1$ is a k -sphere. Then

$$\rho(K) = \rho(K_0) + \rho(K_1) - \rho(K_2) = 1 + \chi(K_1) - \chi(S^k) = \chi(K_1) + (-1)^{k+1} = \chi(K).$$

To prove Lemma 1, in turn, it is technically convenient to consider finite *regular cell complexes* in addition to the more special category of finite simplicial complexes. (See [SCF] for definitions and basic properties of regular cell complexes.) Let d_1 be a function defined on pairs (J, p) where J is a regular cell complex which is the union of cells all containing the vertex p . It is understood that d_1 depends only on the isomorphism class of (J, p) as a regular cell-complex pair. Thus in the case when J happens to be simplicial, we see that d_1 depends only on the simplicial isomorphism class of (J, p) and thus, since J will in this instance be $c(\text{lk } p)$, only on the simplicial isomorphism class of $\text{lk } p$. Consequently, d_1 may be viewed as an extension of a function $d(L)$, defined on simplicial complexes of the sort we have heretofore been considering (that is, $d(L) = d_1(cL, *)$). We say that d_1 determines an $\mathbb{R}$ -valued homotopy invariant ρ of finite regular cell-complexes

if and only if $\rho(K) = \sum_v d_1(\text{st } v, v)$ where the sum is taken over the vertices v of K and where $\text{st}(v)$ is now understood to mean the union of all those cells of K containing v .

Lemma 2. *If d (defined on simplicial complexes) locally determines ρ on simplicial complexes, then there is an extension of d to regular cell-complex pairs (J, p) as above which locally determines ρ on regular cell-complexes.*

(Of course, it is understood that since regular cell-complexes are triangulable—in fact the first barycentric subdivision is a simplicial complex—the invariant ρ automatically extends to regular cell-complexes.)

The proof of Lemma 2 is quite straightforward. Given (J, p) let K be the first barycentric subdivision of J , and hence a simplicial complex. Define $d_1(J, p)$ as follows: let e be a cell of J , b_e its barycenter, hence a vertex of K . Let $V(e)$ denote the number of vertices of the regular cell e . Then set $d_1(J, p) = \sum_e (1/V(e)) d(\text{lk}_K b_e)$ where the sum is taken over all cells of J . The assertion that d_1 must locally determine ρ on regular cell complexes is an immediate consequence of this definition.

We now proceed to the proof of Lemma 1. Assume that d , which locally determines ρ on simplicial complexes, has been extended to d_1 , which locally determines ρ on regular cell-complexes. Let M be a finite regular cell-complex and let I denote, as usual, the unit interval as a simplicial complex with one 1-simplex $[0, 1]$ and two vertices 0 and 1. $M \times I$ is then well defined as a regular cell-complex without need of further subdivision. If M has vertices $v_1, \dots, v_k$, then $M \times I$ has vertices $u_1, \dots, u_k, w_1, \dots, w_k$ where $u_i = (v_i, 0)$, $w_i = (v_i, 1)$:

Lemma 3.

$$\begin{aligned} \sum_{i=1}^k d_1(\text{st}_{M \times I} u_i, u_i) &= \sum_{i=1}^k d_1(\text{st}_{M \times I} w_i, w_i) \\ &= \frac{1}{2} \sum_{i=1}^k d_1(\text{st}_M v_i, v_i) = \frac{1}{2} \rho(M). \end{aligned}$$

Proof. $(\text{st}_{M \times I} u_i, u_i)$ is isomorphic as a regular cell-complex pair to $(\text{st}_{M \times I} w_i, w_i)$, thus it is immediate that

$$\sum_{i=1}^k d_1(\text{st}_{M \times I} u_i, u_i) = \sum_{i=1}^k d_1(\text{st}_{M \times I} w_i, w_i).$$

But

$$\sum_{i=1}^k d_1(\text{st}_{M \times I} u_i, u_i) + \sum_{i=1}^k d_1(\text{st}_{M \times I} w_i, w_i) = \rho(M \times I) = \rho(M) = \sum_{i=1}^k d_1(\text{st}_M v_i, v_i),$$

which yields the remainder of the lemma. $\square$

Now let I' denote the first subdivision of I with two 1-simplices $[0, \frac{1}{2}]$ and $[\frac{1}{2}, 1]$ and three vertices $0, \frac{1}{2}, 1$. With M, v_i as above, $M \times I'$ is a regular cell-complex with vertices $u_i = (v_i, 0), w_i = (v_i, 1), x_i = (v_i, \frac{1}{2})$. Clearly, $(st_{M \times I'} u_i, u_i)$ is isomorphic as a regular cell-complex pair to $(st_{M \times I} u_i, u_i)$ and similarly for w_i . This observation leads to:

Lemma 4. $\sum_{i=1}^k d_1(st_{M \times I'} x_i, x_i) = 0$.

Proof.

$$\rho(M \times I') = \sum_{i=1}^k d_1(st_{M \times I'} u_i, u_i) + \sum_{i=1}^k d_1(st_{M \times I'} w_i, w_i) + \sum_{i=1}^k d_1(st_{M \times I'} x_i, x_i).$$

Also

$$\rho(M \times I') = \rho(M \times I) = \sum_{i=1}^k d_1(st_{M \times I} u_i, u_i) + \sum_{i=1}^k d_1(st_{M \times I} w_i, w_i).$$

By the remarks immediately preceding the statement of the lemma, the two summands on the right-hand side of the second equation are respectively equal to the first two summands in the right-hand side of the second. Hence the remaining summand, namely $\sum_{i=1}^k d_1(st_{M \times I'} x_i, x_i)$, must vanish. $\square$

Now we complete the proof of Lemma 1. Let $K = K_0 \cup K_1$ be a simplicial complex with $K_0 \cap K_1 = K_2$. We construct a homotopy-equivalent regular cell-complex $K_0 \cup (K_2 \times I') \cup K_1 = B$ where K_0, K_1 are now disjoint and $K_2 \times 0$ is identified with the copy of K_2 in K_0 and $K_2 \times 1$ with the copy of K_2 in K_1 . Let $B_0 = K_0 \cup (K_2 \times [0, \frac{1}{2}])$, $B_1 = K_2 \times [\frac{1}{2}, 1] \cup K_1$. Thus $B_0 \cap B_1 = K_2 \times \frac{1}{2}$. We denote the vertices of K_2 by $v_1, \dots, v_k$. Thus $K_2 \times \frac{1}{2}$ has vertices $x_i = (v_i, \frac{1}{2})$, $i = 1, \dots, k$. Two elementary observations:

- (i) $(st_{B_1} x_i, x_i)$ is isomorphic (as a regular cell-complex pair) to $(st_{K_2 \times I}(v_i, 0), (v_i, 0))$ and likewise for $(st_{B_2} x_i, x_i)$.
- (ii) $(st_B x_i, x_i)$ is identical with $(st_{K_2 \times I'} x_i, x_i)$.

Now, since d_1 locally determines ρ , we have

$$\rho(K_0) = \rho(B_0) = \sum_{i=1}^k d_1(st_{B_0} x_i, x_i) + Y,$$

where Y involves only the stars of vertices of B_0 not in $K_2 \times \frac{1}{2}$. Likewise,

$$\rho(K_1) = \sum_{i=1}^k d_1(st_{B_1} x_i, x_i) + Z,$$

where Z involves only the stars of vertices of B_1 not in $K_2 \times \frac{1}{2}$. Thus, by Lemma 3 and observation (i) above, we see immediately that

$$\sum_{i=1}^k d_1(\text{st}_{B_0} x_i, x_i) = \sum_{i=1}^k d_1(\text{st}_{B_1} x_i, x_i) = \frac{1}{2}\rho(K_2).$$

On the other hand, it is directly seen that $\rho(K) = \rho(B) = \sum_{i=1}^k d_1(\text{st}_{B_1} x_i, x_i) + Y + Z$. However, $\sum_{i=1}^k d_1(\text{st}_B x_i, x_i)$ vanishes by virtue of observation (ii) and Lemma 4. So $\rho(K) = Y + Z = [\rho(K_0) - \frac{1}{2}\rho(K_2)] + [\rho(K_1) - \frac{1}{2}\rho(K_2)] = \rho(K_0) + \rho(K_1) - \rho(K_2)$. The proof of Lemma 1, and hence of Theorems A and A', is thus complete.

The kindred result Theorem A'' is easily established by a slightly modified version of this reasoning. The key point is the following lemma, analogous to Lemma 1.

Lemma 5. *Let ρ be a locally determined numerical PL-homeomorphism invariant for compact PL n -manifolds. Let K be a compact PL n -manifold of the form $K = K_0 \cup K_1$ where K_0, K_1 are themselves compact n -manifolds and where $K_2 = K_0 \cap K_1$ is a codimension 0 submanifold of both ∂K_0 and ∂K_1 . Then $\rho(K) = \rho(K_0) + \rho(K_1) - \rho(K_2 \times I)$.*

First, we see quite easily that Lemma 5 implies Theorem A''. For a compact PL n -manifold M , let $h(M)$ denote the dimension of the highest dimensional handle in a handlebody-decomposition of M where this highest dimension is minimal (with respect to all possible handlebody structures). Our proof runs by induction on $h(M)$.

If $h(M) = 0$, then M is the disjoint union of some finite number m of n -disks, whence $\rho(M) = m = \chi(M)$.

Now suppose A'' holds for all compact manifolds M with $h(M) \leq j < n$. Consider M_1 with $h(M_1) = j + 1$. Then $M_1 = M \cup ((j + 1)\text{-handles})$ where $h(M) \leq j$, whence $\rho(M) = \chi(M)$ by inductive assumption. Let N denote the union of all the $(j + 1)$ -handles of M_1 , i.e., if there are m such handles, N is the disjoint union of m n -disks, and so $\rho(N) = m = \chi(N)$. Let $L = M \cap N \subseteq \partial M, \partial N$. L is the disjoint union of m copies of $S^j \times D^{n-j-1}$, hence $h(L \times I) = j$ and $\rho(L \times I) = \chi(L \times I) = m(1 + (-1)^j)$. By lemma 5,

$$\begin{aligned} \rho(M_1) &= \rho(M) + \rho(N) - \rho(L \times I) = \chi(M) + \chi(N) - \chi(L \times I) \\ &= \chi(M) + \chi(N) - \chi(L) = \chi(M_1). \end{aligned}$$

The induction is thus complete.

As for Lemma 5 itself, we note that the argument for Lemma 1 goes through almost word for word. Note that lemma 2 holds in the context of regular cell-complex decompositions of PL manifolds. The analogue of Lemma 3 holds where M is now a compact triangulated $(n - 1)$ -manifold. The modified result

reads

$$\sum_{i=1}^k d_1(\text{st}_{M \times I} u_i, u_i) = \sum_{i=1}^k d_1(\text{st}_{M \times I} w_i, w_i) = \frac{1}{2} \rho(M \times I).$$

Lemma 4 holds as well in this context.

The computations leading to Lemma 1 now serve equally well for Lemma 5. Here it need only be observed that if K is a compact PL n -manifold decomposed as $K_0 \cup K_1$ (with $K_2 = K_0 \cap K_1$ a codimension-0 submanifold of ∂K_0 and ∂K_1), then B (as defined in the proof of Lemma 1) is now a compact PL n -manifold PL homeomorphic to K while B_0, B_1 are PL manifolds homeomorphic, respectively, to K_0 and K_1 . The proof then goes through substituting $\rho(K_2 \times I)$ for $\rho(K_2)$ as appropriate.

3. Proof of Theorem B

As noted in the introduction, the Euler characteristic χ is locally determined by

$$e(L) = 1 + \sum_{i=0}^{\dim L} \frac{(-1)^{i+1}}{i+2} (\text{number of } i\text{-simplices of } L).$$

We now show that no other function on simplicial isomorphism classes of finite complexes can locally determine χ .

To this end, let d be some other function for which $\chi(K) = \sum_v d(\text{lk } V)$ for all finite simplicial complexes K . Given any simplicial complex J with σ a simplex, we call σ a *maximal* simplex of J if and only if σ is not a face of any larger simplex. Note that any finite complex is the union of its maximal simplices.

Our proof that $d(L) \equiv e(L)$ proceeds via induction on the number of maximal simplices of L . In the case where L has but one maximal simplex, it is clear that L must be isomorphic to a standard simplex, say Δ^k for some $k \geq 0$. Consider $K = v * L$ for some disjoint vertex v . $v * L$ is of course isomorphic to Δ^{k+1} so $1 = \chi(v * L) = d(L) + \sum_{i=0}^k d(\text{lk}_K v_i)$ (where $v_0, \dots, v_k$ are the vertices of L) since $L = \text{lk}_K v$. But $\text{lk}_K v_i$ is obviously a k -simplex for $i = 0, \dots, k$. So we have $d(\text{lk}_K v_i) = d(L)$, whence $1 = (k+2)d(L)$, $d(L) = 1/(k+2) = e(L)$.

Now suppose $d(L) = e(L)$ for all L having $\leq j$ maximal simplices. Let L have $j+1$ maximal simplices and let $v_1, \dots, v_r$ denote the vertices of L . Consider $K = v * L \cong cL$ where v is a vertex distinct from $v_1, \dots, v_r$. $L = \text{lk}_K v$. Consider $\text{lk}_K v_i$. This is clearly $v * \text{lk}_L v_i \cong c \text{lk}_L v_i$. Thus $\text{lk}_K v_i$ is isomorphic to $\text{st}_L v_i$. However, $\text{st}_L v_i$ is clearly the union of maximal simplices of L . Hence $\text{lk}_K v_i$ has $\leq j+1$ maximal simplices.

Now because K is contractible, $1 = \chi(K) = d(L) + \sum_{i=1}^r d(\text{lk}_K v_i)$. Segregate the v_i into two classes, u_i , $i = 1, \dots, s$, and w_i , $i = 1, \dots, t$, with $s+t=r$, by the criterion that $\text{lk}_K u_i$ has $\leq j$ maximal simplices whereas $\text{lk}_K w_i$ has $j+1$ maximal

simplices. Thus $d(\text{lk}_K u_i) = e(\text{lk}_K u_i)$ for $i = 1, \dots, s$. Thus since

$$\begin{aligned}\chi(K) &= d(L) + \sum_{i=1}^s d(\text{lk}_K u_i) + \sum_{i=1}^t d(\text{lk}_K w_i) \\ &= e(L) + \sum_{i=1}^s e(\text{lk}_K u_i) + \sum_{i=1}^t e(\text{lk}_K w_i),\end{aligned}$$

we have

$$d(L) + \sum_{i=1}^t d(\text{lk}_K w_i) = e(L) + \sum_{i=1}^t e(\text{lk}_K w_i).$$

Note that v and w_i , $i = 1, \dots, t$, may be characterized as those vertices of K which are common to all the maximal simplices of K . It follows that $v, w_1, \dots, w_t$ are the vertices of $\tau = \bigcap_{\alpha} \sigma_{\alpha}$, where $\{\sigma_{\alpha}\}$ is the set of maximal simplices of K . For any pair of these vertices, it is clear that there is thus a simplicial automorphism of K carrying the first into the second. In other words, $L = \text{lk}_K v \cong \text{lk}_K w_i$, $i = 1, \dots, t$. Hence we see that $(t+1)d(L) = (t+1)e(L)$ whence $d(L) = e(L)$. This completes the proof.

References

- [C] Cheeger, J., Spectral geometry of singular Riemannian spaces, *J. Differential Geom.* **18** (1983), 575–657.
- [GM] Gelfand, I. M., and MacPherson, R., A combinatorial formula for the Pontrjagin classes (pre-print).
- [K] Klee, V., A combinatorial analogue of Poincaré's duality theorem, *Canad. J. Math.* **16** (1964), 517–531.
- [L] Levitt, N., *Grassmannians and Gauss Maps in Piecewise-Linear Topology*, Lecture Notes in Mathematics, Vol. 1366, Springer-Verlag, Berlin, 1989.
- [LR] Levitt, N., and Rourke, C., The existence of combinatorial formulae for characteristic classes, *Trans. Amer. Math. Soc.* **239** (1978), 391–397.
- [SCF] Steenrod, M., Cooke, G., and Finney R., *Homology of Cell Complexes*, Princeton University Press, Princeton, NJ, 1967.
- [W1] Wall, C. T. C., Classification problems in differential topology, IV—Thickenings, *Topology* **5** (1966), 73–94.
- [W2] Wall, C. T. C., Arithmetic invariants of subdivisions of complexes, *Canad. J. Math.* **18** (1966), 72–96.

Received September 13, 1989, and in revised form July 3, 1990.