

## SOME CONSEQUENCES OF A THEOREM OF BOTT

BY JOHN MILNOR<sup>1</sup>

(Received February 11, 1958)

It will be shown that the following theorem, due to R. Bott [3], can be used to solve several well known problems ; including the problem of the existence of division algebras, and the parallelizability of spheres<sup>2</sup>. (Independent solutions of these problems, also based on Bott's work, have been given by Kervaire and Hirzebruch.)

**THEOREM OF BOTT.** *For any  $O_m$ -bundle  $\xi$  over the sphere  $S^{4k}$ , the Pontrjagin class  $p_k(\xi) \in H^{4k}(S^{4k}; Z)$  is divisible by  $(2k - 1)!$ . (This result was conjectured, and proved up to powers of 2, by Borel and Hirzebruch [1]).*

The following result, which follows from Wu Wen-Tsün [19], will also be needed. Since Wu's paper is in Chinese a proof is included in the appendix. The epimorphism  $Z \rightarrow Z_4$  induces a homomorphism  $H^*(K; Z) \rightarrow H^*(K; Z_4)$  which will be denoted by  $\alpha \rightarrow (\alpha)_4$ . Let  $i: Z_2 \rightarrow Z_4$  denote the inclusion homomorphism.

**THEOREM OF WU.** *For any  $O_m$ -bundle  $\xi$  over a complex  $K$ , the class  $(p_k(\xi))_4 \in H^{4k}(K; Z_4)$  is determined by the Stiefel-Whitney classes  $w_i(\xi) \in H^i(K; Z_2)$ . In particular if the Stiefel-Whitney classes  $w_1(\xi), \dots, w_{4k-1}(\xi)$  are zero then  $(p_k(\xi))_4 = i_* w_{4k}(\xi)$ .*

Combining these two results, the following is obtained.

**THEOREM 1.** *There exists an  $O_m$ -bundle  $\xi$  over the sphere  $S^n$  with  $w_n(\xi) \neq 0$  only for  $n = 1, 2, 4$  or  $8$ .*

(Examples of such bundles can be given as follows : for  $n = 1$  the 2-fold covering of the circle, and for  $n = 2, 4$  or  $8$  the  $O_n$ -bundle over  $S^n$  associated with the Hopf fibering  $S^{2n-1} \rightarrow S^n$ .)

**PROOF.** According to Wu [16] such a bundle can exist only if  $n$  is a power of 2. Hence it is certainly sufficient to consider the case  $n = 4k$ ,  $k > 2$ . The identity

$$(p_k(\xi))_4 = i_* w_{4k}(\xi) \in H^{4k}(S^{4k}; Z_4) = Z_4$$

is valid, since the lower Stiefel-Whitney classes must be zero. In other words the class  $w_{4k}(\xi)$  is zero if and only if  $p_k(\xi)$  is divisible by 4. But

<sup>1</sup> The author holds a Sloan fellowship.

<sup>2</sup> A preliminary account of this work has been given in [20].

$p_k(\xi)$  is known to be divisible by  $(2k-1)!$ . For  $k > 2$  this proves that  $w_{4k}(\xi) = 0$ .

**THEOREM 2.** *The sphere  $S^r$  is parallelizable only for  $r = 1, 3, 7$ . (Compare Steenrod and Whitehead [10].)*

**PROOF.** The fibering  $SO_r \longrightarrow SO_{r+1} \xrightarrow{f} S^r$  associated with the tangent bundle of  $S^r$  has the following homotopy sequence:

$$\longrightarrow \pi_r(SO_{r+1}) \xrightarrow{f_*} \pi_r(S^r) \xrightarrow{\partial} \pi_{r-1}(SO_r) \longrightarrow \pi_{r-1}(SO_{r+1}) \longrightarrow 0.$$

The group  $\pi_r(S^r)$  will be identified with the integers. Then  $\partial(1) \in \pi_{r-1}(SO_r)$  is the element which corresponds to the tangent bundle of  $S^r$ . (See Steenrod [9, §18]).

For each  $\lambda \in \pi_r(SO_{r+1})$  let  $\xi$  denote the corresponding  $SO_{r+1}$ -bundle over  $S^{r+1}$ , and let  $X(\xi)$  denote its Euler class (=top Stiefel-Whitney class with integer coefficients). Let  $\mu$  be the standard generator of  $H_{r+1}(S^{r+1}; Z)$ . Then  $f_*(\lambda)$  is equal to the negative of the "Euler number"  $\langle X(\xi), \mu \rangle$ . [Proof. Let  $\nu(\xi) \in H^{r+1}(S^{r+1}; \pi_r(SO_{r+1}))$  denote the obstruction to the existence of a cross-section of  $\xi$ . Then  $X(\xi)$ , the obstruction to the existence of a cross-section in the associated sphere bundle, is equal to  $f_*(\nu(\xi))$ . According to Steenrod [9, p. 180] the identity  $\langle \nu(\xi), \mu \rangle = -\lambda$  is satisfied. Therefore  $\langle X(\xi), \mu \rangle = -f_*(\lambda)$ ].

Now if  $S^r$  is parallelizable then  $\partial(1) = 0$ , hence there exists  $\lambda \in \pi_r(SO_{r+1})$  with  $f_*(\lambda) = 1$ . For the corresponding bundle  $\xi$ , the class  $X(\xi)$  generates  $H^{r+1}(S^{r+1}; Z)$ ; hence the class  $w_{r+1}(\xi) = (X(\xi))_2$  is non-zero. Together with Theorem 1 this completes the proof.

It follows immediately that the real projective space  $P^r$  is parallelizable only for  $r = 1, 3, 7$ . (For consequences concerning the immersion of  $P^r$  in euclidean space see Milnor, Comm. Math. Helv. 30 (1956), p. 284).

**COROLLARY 1.** *There exists a division algebra of rank  $n$  over the real numbers only for  $n = 1, 2, 4, 8$ .*

**PROOF.** The existence of a bilinear product operation without zero-divisors in the vector space  $R^n$  implies that the projective space  $P^{n-1}$  is parallelizable. (See Stiefel [11, p. 216]). [ALTERNATIVE PROOF. Suppose that such a product operation in  $R^n$  is given. Then the correspondence  $S^{n-1} \rightarrow GL_n$  defined by  $x \rightarrow$  (left multiplication by  $x$ ) gives rise to a  $GL_n$ -bundle  $\xi$  over  $S^n$ . It is not hard to verify that  $w_n(\xi) \neq 0$ ].

**COROLLARY 2.** *For  $r \geq 8$  the groups  $\pi_{r-1}(SO_r)$  are as follows:*

$r$ modulo 8:	0	1	2	3	4	5	6	7
$\pi_{r-1}(SO_r)$ :	$Z + Z$	group of order 4	$Z + Z_2$	$Z_2$	$Z + Z$	$Z_2$	$Z$	$Z_2$

PROOF. This follows from Bott's computation [2] of the stable groups  $\pi_{r-1}(SO_{r+1})$ , together with the exact homotopy sequence used to prove Theorem 2.

THEOREM 3. *Let  $M^{2n}$  be a simply-connected differentiable manifold such that the cohomology group  $H^i(M^{2n}; Z)$  is infinite cyclic for  $i = 0, n, 2n$ , and zero otherwise. Then  $n$  must be 2, 4, or 8.*

(Examples are provided by the complex, quaternion, and Cayley projective planes. It will be shown in a later paper [7] that the condition of simple-connectivity can be eliminated. This will give an answer to Problem 5 of [5]).

PROOF. If  $\alpha$  generates  $H^n(M^{2n}; Z)$ , then the Poincaré duality theorem implies that  $\alpha \smile \alpha$  generates  $H^{2n}(M^{2n}; Z)$ . Hence

$$Sq^n : H^n(M^{2n}; Z_2) \longrightarrow H^{2n}(M^{2n}; Z_2)$$

is non-zero. The formulas of Wu [15] now imply that the Stiefel-Whitney class  $w_n$  of the tangent bundle  $\theta$  is non-zero. Choose a map  $g : S^n \rightarrow M^{2n}$  which, under the Hurewicz homomorphism, corresponds to a generator of  $H_n(M^{2n}; Z)$ . Then the bundle  $\theta'$  over  $S^n$  induced from  $\theta$  by  $g$  will satisfy  $w_n(\theta') \neq 0$ . Therefore  $n$  must be 1, 2, 4 or 8. Since the case  $n = 1$  is easily excluded, this completes the proof.

Bott's theorem is related to the question of the existence of maps with Hopf invariant 1 as follows. Let  $J : \pi_{n-1}(SO_m) \rightarrow \pi_{m+n-1}(S^m)$  be the homomorphism of G. W. Whitehead [13], and let  $\gamma_n : \pi_{m+n-1}(S^m) \rightarrow Z_2$  be the generalized Hopf invariant of Steenrod [8], which is defined using the functional  $Sq^n$  operation. For each odd prime  $q$  let

$$\gamma_{q,i} : \pi_{m+2i(q-1)-1}(S^m) \longrightarrow Z_q$$

denote the corresponding homomorphism based on the reduced  $q^{\text{th}}$  power  $\mathcal{P}^i$ .

THEOREM 4a. *The image  $J\pi_{n-1}(SO_m)$ ,  $m \geq n$ , contains an element  $J\lambda$  with generalized Hopf invariant  $\gamma_n(J\lambda)$  different from zero only if  $n$  equals 2, 4, or 8.*

THEOREM 4b. *The image  $J\pi_{2i(q-1)-1}(SO_m)$ ,  $m \geq 2i(q-1)$ , contains an element  $J\lambda$  with  $\gamma_{q,i}(J\lambda)$  different from zero only if  $i = 1$ .*

PROOF OF 4b. Let  $\xi$  be the  $SO_m$ -bundle over  $S^n$  associated with  $\lambda$ , where  $n = 2i(q-1)$ . Let  $E$  be the total space of the associated bundle having the unit ball  $B^m$  as fibre, so that the boundary  $\dot{E}$  is the total space of the associated sphere bundle. According to [7, Theorem 3, Corollary 1],

the collapsed space  $E/\dot{E}$  can be obtained from the sphere  $S^m$  by attaching an  $(m+n)$ -cell, using on attaching map in the homotopy class  $J\lambda$ . Thus the generalized Hopf invariant  $\gamma_{q,i}(J\lambda)$  is non-zero if and only if the homomorphism

$$\mathcal{P}^i : H^m(E, \dot{E}; Z_q) \longrightarrow H^{m+n}(E, \dot{E}; Z_q)$$

is non-zero.

Let  $\phi : H^j(S^n; Z_q) \rightarrow H^{j+m}(E, \dot{E}; Z_q)$  denote the isomorphism of Thom [12]. According to Wu [18, § IV] the class  $\phi^{-1}\mathcal{P}^i\phi(1) \in H^n(S^n; Z_q)$  can be expressed as a polynomial in the Pontrjagin classes of  $\xi$ , reduced modulo  $q$ . But these Pontrjagin classes are zero, except for  $p_{i(q-1)/2}(\xi)$  which is divisible by  $(i(q-1)-1)!$ . For  $i > 1$ , since the number  $(i(q-1)-1)!$  is divisible by  $q$ , it follows that the operation  $\mathcal{P}^i$  must be zero.

Theorem 4a is proved in a similar way, using Theorem 1 together with Thom's definition of the Stiefel-Whitney classes. (See [12]).

### Appendix

PROOF OF THE THEOREM OF WU. Following Hirzebruch [6] define the Pontrjagin class  $p_k$  of an  $O_m$ -bundle as  $(-1)^k$  times the Chern class  $c_{2k}$  of the  $U_m$ -bundle induced by the inclusion  $O_m \rightarrow U_m$ . This is slightly different from the Pontrjagin class as defined by Pontrjagin and Wu. (Compare [17, Theorem 4]).

Consider the exact sequence of cohomology group corresponding to the coefficient sequence

$$0 \longrightarrow Z_2 \xrightarrow{i} Z_4 \xrightarrow{j} Z_2 \longrightarrow 0;$$

as well as the Pontrjagin squaring operation

$$\mathfrak{P} : H^{2k}(K; Z_2) \longrightarrow H^{4k}(K; Z_4).$$

(See for example Whitehead [14]).

LEMMA 1. *The Pontrjagin class  $p_k$  of any  $O_m$ -bundle is related to the Stiefel-Whitney classes  $w_1, \dots, w_{4k}$  by an identity*

$$(p_k)_4 = \mathfrak{P}(w_{2k}) + i_* f_k(w_1; \dots, w_{4k})$$

where  $f_k$  is a polynomial with coefficients in  $Z_2$ .

PROOF. It is clearly sufficient to consider the case of the universal bundle over the Grassmann space  $G_m(R)$ , with  $m$  large. The identity  $j_* \mathfrak{P}(w) = w \smile w$  holds for any cohomology class  $w$ . Comparing this with the relation<sup>3</sup>

<sup>3</sup> See Wu [17, Theorem 3].

$$j_*((p_k)_i) = (p_k)_2 = w_{2k} \smile w_{2k}$$

it follows that

$$(p_k)_i - \mathfrak{P}w_{2k} \in (\text{kernel } j_*) = i_* H^{4k}(G_m(R); Z_2) .$$

Since the cohomology ring  $H^*(G_m(R); Z_2)$  is generated by the Stiefel-Whitney classes, this proves Lemma 1.

To prove the theorem it is only necessary to show that the coefficient of  $w_{4k}$  in  $f_k$  is non-zero. Let  $\mathfrak{T}$  denote the universal  $U_m$ -bundle over the complex Grassmann space  $G_m(C)$ . Recall that the cohomology ring  $H^*(G_m(C); Z)$  is a polynomial ring<sup>4</sup> generated by the Chern classes of  $\mathfrak{T}$ . The inclusion  $U_m \rightarrow O_{2m}$  induces an  $O_{2m}$ -bundle over  $G_m(C)$  which will be denoted by  $\mathfrak{T}_R$ . Applying Lemma 1 to this bundle  $\mathfrak{T}_R$ , the relations<sup>5</sup>

$$p_k(\mathfrak{T}_R) = c_k(\mathfrak{T})^2 - 2c_{k-1}(\mathfrak{T})c_{k+1}(\mathfrak{T}) + \cdots \pm 2c_0(\mathfrak{T})c_{2k}(\mathfrak{T})$$

and

$$w_{2r+1}(\mathfrak{T}_R) = 0, \quad w_{2r}(\mathfrak{T}_R) = (c_r(\mathfrak{T}))_2$$

show that the polynomial  $f_k$  must satisfy

$$f_k(0, w_2, 0, w_4, \dots, w_{4k}) = w_{2k-2}w_{2k+2} + w_{2k-4}w_{2k+4} + \cdots + w_0w_{4k} .$$

Therefore  $f_k(0, 0, \dots, 0, w_{4k}) = w_{4k}$ ; which completes the proof.

PRINCETON UNIVERSITY

#### REFERENCES

1. A. BOREL and F. HIRZEBRUCH, *Characteristic classes and homogeneous spaces*, to appear (Amer. J. Math.).
2. R. BOTT, *On the stable homotopy of the classical groups*, Proc. Nat. Acad. Sci. U.S.A. 43 (1957), 933-935.
3. ———, *The space of loops on a Lie-group*, to appear (Mich. J. Math.).
4. S. S. CHERN, *Characteristic classes of Hermitian manifolds*, Ann. of Math. 47 (1946), 85-121.
5. F. HIRZEBRUCH, *Some problems on differentiable and complex manifolds*, Ann. of Math. 60 (1954), 213-236.
6. ———, *Neue topologische Methoden in der algebraischen Geometrie*, Springer, 1956.
7. J. MILNOR, *On spaces with a gap in cohomology*, to appear.
8. N. E. STEENROD, *Cohomology invariants of mappings*, Ann. of Math. 50 (1949), 954-988.
9. ———, *The topology of fibre bundles*, Princeton, 1951.
10. ——— and J.H.C. WHITEHEAD, *Vector fields on the  $n$ -sphere*, Proc. Nat. Acad. Sci. U.S.A. 37 (1951), 58-63.
11. E. STIEFEL, *Über Richtungsfelder in den projektiven Räumen und einen Satz aus der reellen Algebra*, Comm. Math. Helv. 13 (1940), 201-218.
12. R. THOM, *Espaces fibrés en sphères et carrés de Steenrod*, Ann. sci. Ecole norm. sup 69 (1952), 109-182.

<sup>4</sup> Chern [4].

<sup>5</sup> See Hirzebruch [6, p. 68] and Steenrod [9, p. 212].

13. G. W. WHITEHEAD, *On the homotopy groups of spheres and rotation groups*, Ann. of Math. 43 (1942), 634–640.
14. J.H.C. WHITEHEAD, *On simply connected 4-dimensional polyhedra*, Comm. Math. Helv. 22 (1949), 48–92.
15. WU WEN-TSÜN, *Classes caractéristiques et  $i$ -carrés d'une variété*, C. R. Acad. Sci. Paris 230 (1950), 508–511.
16. ———, *Les  $i$ -carrés dans une variété grassmannienne*, C. R. Acad. Sci. Paris 230 (1950), 918–920.
17. ———, *On Pontrjagin classes I*, Scientia Sinica 3 (1954), 353–367.
18. ———, *On Pontrjagin classes II*, Scientia Sinica 4 (1955), 455–490.
19. ———, *On Pontrjagin classes III*, Acta Math. Sinica 4 (1954), 323–346.
20. R. BOTT and J. MILNOR, *On the parallelizability of the spheres*, Bull. Amer. Math. Soc. 64 (1958), 87–89.