



On the Stiefel-Whitney Classes of a Manifold. II

W. S. Massey

Proceedings of the American Mathematical Society, Vol. 13, No. 6 (Dec., 1962),
938-942.

Stable URL:

<http://links.jstor.org/sici?sici=0002-9939%28196212%2913%3A6%3C938%3AOTSCOA%3E2.0.CO%3B2-P>

Proceedings of the American Mathematical Society is currently published by American Mathematical Society.

Your use of the JSTOR archive indicates your acceptance of JSTOR's Terms and Conditions of Use, available at <http://uk.jstor.org/about/terms.html>. JSTOR's Terms and Conditions of Use provides, in part, that unless you have obtained prior permission, you may not download an entire issue of a journal or multiple copies of articles, and you may use content in the JSTOR archive only for your personal, non-commercial use.

Please contact the publisher regarding any further use of this work. Publisher contact information may be obtained at <http://uk.jstor.org/journals/ams.html>.

Each copy of any part of a JSTOR transmission must contain the same copyright notice that appears on the screen or printed page of such transmission.

JSTOR is an independent not-for-profit organization dedicated to creating and preserving a digital archive of scholarly journals. For more information regarding JSTOR, please contact support@jstor.org.

ON THE STIEFEL-WHITNEY CLASSES OF A MANIFOLD. II

W. S. MASSEY¹

1. Introduction. This note is a complement to a previous paper of the same title [1]. In that paper it was proved that certain Stiefel-Whitney classes or dual Stiefel-Whitney classes modulo 2 of a differentiable manifold always vanished. In the present note, analogous theorems are proved about the *integral* Stiefel-Whitney classes of a manifold.

The following convention regarding notation will be followed consistently. The mod 2 Stiefel-Whitney classes are denoted by lower-case letters, w_i , while the integral Stiefel-Whitney classes are denoted by capital letters, W_i ; the subscript denotes the dimension. A bar over the appropriate symbol denotes the *dual* Stiefel-Whitney class, integral or mod 2, thus: $\overline{W}_i, \overline{w}_i$. Of course the integral classes are only defined in odd dimensions.

Let M^n denote a compact, connected, orientable, differentiable n -manifold. We will prove the following three theorems about its integral Stiefel-Whitney classes:

THEOREM 1. *If n is even, $\overline{W}_{n-1}=0$.*

THEOREM 2. *If n is even, $W_{n-1}=0$.*

THEOREM 3. *If $n \equiv 3 \pmod{4}$, $W_{n-2}=0$.*

Note that Theorem 1 implies Corollary 1 to Theorem I of [1] in the orientable case. Similarly, Theorem 2 implies Theorem II of [1], and Theorem 3 represents a strengthening of part of the conclusion of Theorem III of [1].

The main interest in Theorem 1 stems from the fact that A. Haefliger and M. Hirsch have recently proved [3] that any (compact, orientable, differentiable) M^n , $n > 4$, is differentially imbeddable in R^{2n-1} if and only if $\overline{W}_{n-1}=0$ for n even and $\overline{w}_{n-1}=0$ for n odd. Thus it follows from Theorem 1 above and Corollary 1 to Theorem I of [1] that such an M^n is always differentially imbeddable in R^{2n-1} for $n > 4$.

An interesting application of Theorem 2 is to 8-dimensional manifolds. According to [2, p. 170], for any compact, orientable M^8 , $W_5=0$; by Theorem 2, $W_7=0$. Thus W_3 is the only integral Stiefel-

Received by the editors November 16, 1961.

¹ During the preparation of this note the author was partially supported by N.S.F. Grant G-18895.

Whitney class of an 8-dimensional manifold which can be nonzero. It is known that W_3 is the first obstruction to the existence of an almost complex structure on M^8 , and if $W_3=0$, then W_7 is the second obstruction. Therefore if we remove a single point from any compact, orientable M^8 for which $W_3=0$, the resulting noncompact manifold admits an almost-complex structure.

Note also that Theorem 2 above and Theorem III of [1] imply that the first obstruction to defining a field of tangent 2-frames on a compact, orientable n -manifold always vanishes provided $n \not\equiv 1 \pmod{4}$. This raises the question of determining the second obstruction to such a field.

2. **A lemma.** Assume that M^n is a compact, connected orientable n -manifold. Let T^q denote the torsion subgroup of $H^q(M^n, \mathbf{Z})$. Let p be a prime number, and

$$(S) \quad H^q(M, \mathbf{Z}) \xrightarrow{p} H^q(M, \mathbf{Z}) \xrightarrow{r} H^q(M, \mathbf{Z}_p) \xrightarrow{\delta^*} H^{q+1}(M, \mathbf{Z})$$

be the exact sequence associated with the coefficient sequence

$$0 \rightarrow \mathbf{Z} \xrightarrow{p} \mathbf{Z} \rightarrow \mathbf{Z}_p \rightarrow 0.$$

The cup product is a bilinear form

$$U: H^q(M^n, \mathbf{Z}_p) \times H^{n-q}(M^n, \mathbf{Z}_p) \rightarrow H^n(M^n, \mathbf{Z}_p) \approx \mathbf{Z}_p.$$

According to the Poincaré Duality Theorem, this bilinear form is nondegenerate.

LEMMA 1. *For any q , $r(T^{n-q})$ is the annihilator of $r[H^q(M, \mathbf{Z})]$.*

PROOF. It is clear that $r(T^{n-q})$ is contained in the annihilator of $r[H^q(M, \mathbf{Z})]$. We will complete the proof by showing that $r(T^q)$ has the same rank (as a vector space over \mathbf{Z}_p) as the annihilator of $r(H^q)$. For this purpose we introduce the following notation:

b_i = i th Betti no. of M = rank of $H^i(M, \mathbf{Z})$.

c_i = number of cyclic summands in the p -primary component of T^i (i.e., the p -primary component of T^i is the direct sum of c_i cyclic subgroups).

According to the Poincaré Duality Theorem, $b_i = b_{n-i}$ and $c_i = c_{n-i+1}$. Consideration of the exact sequence (S) shows that the rank of the vector space $H^i(M, \mathbf{Z}_p)$ is $b_i + c_i + c_{i+1}$ while the ranks of the subspaces $r(H^i)$ and $r(T^i)$ are $b_i + c_i$ and c_i , respectively. Therefore the rank of the annihilator of $r(H^q)$ is

$$\begin{aligned} & (b_{n-q} + c_{n-q} + c_{n-q+1}) - (b_q + c_q) \\ &= (b_q + c_{n-q} + c_q) - (b_q + c_q) = c_{n-q} \end{aligned}$$

which is precisely the rank of the subspace $r(T^{n-q})$, as was to be proved.

REMARK. This lemma is apparently well known; see [2, p. 169].

3. **Proof of Theorem 1.** We will use the exact sequence (S) and the preceding lemma for the case $p=2$. It is well known that

$$\overline{W}_{i+1} = \delta^*(\tilde{w}_i), \quad (i \text{ even});$$

in fact, this equation may be taken as the definition of \overline{W}_{i+1} . Hence by exactness of (S), to prove $\overline{W}_{n-1}=0$, it suffices to prove that $\tilde{w}_{n-2} \in r(H^{n-2})$. By the preceding lemma, this is equivalent to proving that \tilde{w}_{n-2} annihilates the subspace $r(T^2) \subset H^2(M, \mathbf{Z}_2)$.

By Lemma 7 of [1], the homomorphism $H^2(M^n, \mathbf{Z}_2) \rightarrow H^n(M^n, \mathbf{Z}_2)$ defined by $x \rightarrow x \cdot \tilde{w}_{n-2}$ is a sum of iterated Steenrod squares, which we may assume to be admissible on account of Adem's relations. By Lemma 4 of [1] we may assume that the excess of any such admissible iterated Steenrod square is 1 or 2. We will complete the proof by showing that for $x \in r(T^2)$,

$$Sq^I(x) = 0,$$

where $Sq^I; H^2(M^n, \mathbf{Z}_2) \rightarrow H^n(M^n, \mathbf{Z}_2)$ is any admissible iterated Steenrod square of excess 1 or 2 and degree $n-2$.

In case the excess is 1, then we must have $I = (2^j, 2^{j-1}, \dots, 2, 1)$ for some integer $j \geq 0$. But in this case it is clear that $Sq^I x = 0$, for $x \in r(T^2)$, because

$$Sq^1 x = r\delta^*(x) = 0$$

by exactness of (S).

In case the excess is 2, by Lemma 5 of [1] there exists an admissible iterated Steenrod square, Sq^J , and a power of 2, $m=2^k$, such that

$$Sq^I x = (Sq^J x)^m$$

and J has excess 0 or 1. In case J has excess 0, then

$$Sq^I x = x^m$$

and it is obvious that $Sq^I x = 0$ for $x \in r(T^2)$. In case J has excess 1, then $J = (2^j, 2^{j-1}, \dots, 2, 1)$ for some integer $j \geq 0$ exactly as before, and $Sq^J(x) = 0$ for $x \in r(T^2)$ for the same reason as before. Thus in either case, $Sq^I(x) = 0$ for $x \in r(T^2)$, as was to be proved.

4. Proof of Theorem 2. We will divide the proof into two cases, according as $n=4k+2$ or $n=4k$. In both cases, use will be made of the following lemma.

LEMMA 2. *For any integer i , U_i^2 is the reduction mod 2 of an integral cohomology class.*

PROOF. Here U_i denotes the class of W_i ; the Stiefel-Whitney classes are defined in terms of the U_i by the formula

$$w_j = \sum_i Sq^{j-i} U_i.$$

It is readily seen by induction on j that the U_i may be expressed as polynomials in the Steenrod squares of the w_j , and hence as polynomials in the Stiefel-Whitney classes w_j (since any Steenrod square of a Stiefel-Whitney class may be expressed as a polynomial in the Stiefel-Whitney classes). Now it is well known that the square of any Stiefel-Whitney class, w_j^2 , is the reduction mod 2 of an integral class; for j even, it is the reduction of a Pontrjagin class, while for j odd it is the reduction of W_j^2 . Hence the square of any polynomial in the w_j is the reduction mod 2 of an integral class. In particular, U_i^2 is the reduction of an integral class.

First we will prove Theorem 2 for the case $n=4k+2$, the easiest case. In this case

$$w_{n-2} = w_{4k} = Sq^{2k} U_{2k} = U_{2k}^2$$

which is the reduction mod 2 of an integral class by the lemma. Hence

$$W_{n-1} = \delta^*(w_{n-2}) = 0$$

by the exactness of (S).

Next, we will prove Theorem 2 for the case $n=4k$. In this case

$$w_{n-2} = w_{4k-2} = Sq^{2k-2} U_{2k}.$$

To prove $W_{n-1}=0$, it suffices to prove that $x \cdot w_{n-2}=0$ for any $x \in r(T^2)$, as in the proof of Theorem 1. To achieve this, it obviously suffices to prove that for $m=2^q$, $q \geq 0$, and $x \in r(T^2)$,

$$(1) \quad x^m Sq^{2k-2m} U_{2k} = x^{2m} Sq^{2k-4m} U_{2k}.$$

For, successive application of (1) with $m=1, 2, 4, 8, \dots$ shows that

$$\begin{aligned} x \cdot w_{n-2} &= x \cdot Sq^{2k-2} U_{2k} = x^2 \cdot Sq^{2k-4} U_{2k} \\ &= x^4 \cdot Sq^{2k-8} U_{2k} = \dots = 0. \end{aligned}$$

To prove (1), note first that since $Sq^1x=0$, the only nonzero Steenrod squares of x^m are

$$\begin{aligned} Sq^0x^m &= x^m, \\ Sq^{2m}x^m &= x^{2m}. \end{aligned}$$

Therefore by Cartan's formula,

$$(2) \quad Sq^{2k-2m}(x^mU_{2k}) = x^mSq^{2k-2m}U_{2k} + x^{2m}Sq^{2k-4m}U_{2k}.$$

But by the definition of the U_i ,

$$\begin{aligned} (3) \quad Sq^{2k-2m}(x^mU_{2k}) &= U_{2k-2m}(x^mU_{2k}) \\ &= (U_{2k-2m}x^m)U_{2k} = Sq^{2k}(U_{2k-2m}x^m) \\ &= (U_{2k-2m}x^m)^2 = U_{2k-2m}^2x^{2m}. \end{aligned}$$

By Lemma 2, U_{2k-2m}^2 is the reduction mod 2 of an integral class; since $x \in r(T^2)$, $U_{2k-2m}^2x^{2m} \in r(T^n)$, i.e.,

$$(4) \quad U_{2k-2m}^2x^{2m} = 0.$$

Combining (2), (3), and (4) gives (1) as desired.

5. Proof of Theorem 3. Let $n=4k+3$; then

$$w_{n-3} = w_{4k} = Sq^{2k}U_{2k} = U_{2k}^2$$

which is the reduction mod 2 of an integral class by Lemma 2, and

$$W_{n-2} = \delta^*(w_{n-3}) = 0$$

by exactness of (S).

BIBLIOGRAPHY

1. W. S. Massey, *On the Stiefel-Whitney classes of a manifold*, Amer. J. Math. **82** (1960), 92-102.
2. F. Hirzebruch and H. Hopf, *Felder von Flächenelementen in 4-dimensionalen Mannigfaltigkeiten*, Math. Ann. **136** (1958), 156-172.
3. A. Haefliger and M. Hirsch, *On the existence and classification of differentiable embeddings* (to appear).

YALE UNIVERSITY